

Climate Change Vulnerability Index Report
Salix pseudomonticola (False mountain willow)

Date: 3 March 2020

Assessor: Walter Fertig, WA Natural Heritage Program

Geographic Area: Washington

Heritage Rank: G5/S1

Index Result: Moderately Vulnerable

Confidence: Very High

Climate Change Vulnerability Index Scores

Section A	Severity	Scope (% of range)
1. Temperature Severity	>6.0° F (3.3°C) warmer	0
	5.6-6.0° F (3.2-3.3°C) warmer	0
	5.0-5.5° F (2.8-3.1°C) warmer	0
	4.5-5.0° F (2.5-2.7°C) warmer	0
	3.9-4.4° F (2.2-2.4°C) warmer	100
	<3.9° F (2.2°C) warmer	0
2. Hamon AET:PET moisture	< -0.119	0
	-0.097 to -0.119	100
	-0.074 to -0.096	0
	-0.051 to -0.073	0
	-0.028 to -0.050	0
	>-0.028	0
Section B		Effect on Vulnerability
1. Sea level rise		Neutral
2a. Distribution relative to natural barriers		Somewhat Increase
2b. Distribution relative to anthropogenic barriers		Neutral
3. Impacts from climate change mitigation		Neutral
Section C		
1. Dispersal and movements		Neutral
2ai Change in historical thermal niche		Neutral
2aii. Change in physiological thermal niche		Increase
2bi. Changes in historical hydrological niche		Neutral
2bii. Changes in physiological hydrological niche		Somewhat Increase
2c. Dependence on specific disturbance regime		Neutral
2d. Dependence on ice or snow-covered habitats		Increase
3. Restricted to uncommon landscape/geological features		Somewhat Increase
4a. Dependence on others species to generate required habitat		Neutral
4b. Dietary versatility		Not Applicable
4c. Pollinator versatility		Neutral
4d. Dependence on other species for propagule dispersal		Neutral
4e. Sensitivity to pathogens or natural enemies		Neutral
4f. Sensitivity to competition from native or non-native species		Somewhat Increase
4g. Forms part of an interspecific interaction not covered above		Neutral
5a. Measured genetic diversity		Unknown
5b. Genetic bottlenecks		Unknown
5c. Reproductive system		Neutral

6. Phenological response to changing seasonal and precipitation dynamics	Neutral
Section D	
D1. Documented response to recent climate change	Neutral
D2. Modeled future (2050) change in population or range size	Unknown
D3. Overlap of modeled future (2050) range with current range	Unknown
D4. Occurrence of protected areas in modeled future (2050) distribution	Unknown

Section A: Exposure to Local Climate Change

A1. Temperature: Both of the known and reported occurrences of *Salix pseudomonticola* in Washington (100%) occur in areas with a projected temperature increase of 3.9-4.4° F (Figure 1).




Figure 1. Exposure of *Salix pseudomonticola* occurrences in Washington to projected local temperature change. Base map layers from www.natureserve.org/ccvi

A2. Hamon AET:PET Moisture Metric: Both of the confirmed and reported occurrences of *Salix pseudomonticola* (100%) in Washington are found in areas with a projected decrease in available moisture (as measured by the ratio of actual to potential evapotranspiration) in the range of -0.097 to -0.119 (Figure 2).




Figure 2. Exposure of *Salix pseudomonticola* occurrences in Washington to projected moisture availability (based on ratio of actual to predicted evapotranspiration). Base map layers from www.natureserve.org/ccvi

Section B. Indirect Exposure to Climate Change

B1. Exposure to sea level rise: Neutral.

Washington occurrences of *Salix pseudomonticola* are found at 2950-5500 feet (900-1680 m) and would not be inundated by projected sea level rise.

B2a. Natural barriers: Somewhat Increase.

In Washington, one occurrence of *Salix pseudomonticola* has been confirmed and is from a hummocky calcareous fen with marl-like soil in Pend Oreille County (Camp and Gamon 2011, WNHP records). This site is a component of the Rocky Mountain Subalpine-Montane Fen ecological system (Rocchio and Crawford 2015). A second report from Chelan County is found in a wet meadow along a mountain trail and needs to be confirmed (Fertig and Kleinknecht 2020). The habitat description is too vague to classify further. The Pend Oreille County occurrence is disjunct from other populations and embedded within unsuitable habitat, which creates a barrier for dispersal.

B2b. Anthropogenic barriers: Neutral.

The range of *Salix pseudomonticola* is naturally somewhat fragmented. Human impacts on the landscape of northeastern Washington have little effect on this condition.

B3. Predicted impacts of land use changes from climate change mitigation: Neutral.

Section C: Sensitive and Adaptive Capacity

C1. Dispersal and movements: Neutral.

Salix pseudomonticola produces numerous, many-seeded dry capsules. Seeds are small and have a tuft of wavy hairs to assist in dispersal by wind. Although average dispersal distance may be short, some seeds are capable of moving over 1 km, and so the species is not dispersal limited.

C2ai. Historical thermal niche: Neutral.

Figure 3 depicts the distribution of *Salix pseudomonticola* in Washington relative to mean seasonal temperature variation for the period from 1951-2006 (“historical thermal niche”). The one confirmed occurrence from Pend Oreille County is from an area with average (57.1-77°F/31.8-43.0°C) temperature variation during the past 50 years and is considered at neutral risk from climate change. The unconfirmed population from Chelan County is from an area with slightly lower than average (47.1-57°F/26.3-31.8°C) temperature variation during the same period and would be at somewhat increased risk from climate change (Young et al. 2006).





Figure 3. Historical thermal niche (exposure to past temperature variations) of *Salix pseudomonticola* occurrences in Washington. Base map layers from www.natureserve.org/ccvi

C2aii. Physiological thermal niche: Increase.

The montane fen habitat of *Salix pseudomonticola* in Pend Oreille County, Washington is associated with cold air drainage during the growing season and would have increased vulnerability to temperature changes associated with global warming.

C2bi. Historical hydrological niche: Neutral.

Both populations of *Salix pseudomonticola* confirmed and reported from Washington (100%) are found in areas that have experienced average or greater than average (>20 inches/508 mm) precipitation variation in the past 50 years (Figure 4). According to Young et al. (2016), this occurrence is at neutral vulnerability to climate change.



C2bii. Physiological hydrological niche: Somewhat Increase.

The fen habitat of *Salix pseudomonticola* in Pend Oreille County, Washington is largely dependent on groundwater, and thus more reliant on snowpack for regeneration than rainfall (Rocchio and Ramm-Granberg 2017). Reduction in the timing and amount of precipitation and

increased drought would make these fens and wet meadow habitat in Chelan County more vulnerable to climate change (Rocchio and Ramm-Granberg 2017).

C2c. Dependence on a specific disturbance regime: Neutral.

Salix pseudomonticola is not dependent on periodic disturbances to maintain its fen or wet meadow habitat in the state. The species could, however, be detrimentally affected by increased summer temperatures, drought, or decreased snowpack that might favor conversion of this habitat to forest or drier meadows, or increase fire frequency (Rocchio and Ramm-Granberg 2017).

C2d. Dependence on ice or snow-cover habitats: Increase.

The populations of *Salix pseudomonticola* in Washington are found in the mountainous areas with high snowfall. Its fen and wet meadow habitat is dependent on late-lying snowbanks for recharging groundwater (Rocchio and Ramm-Granberg 2017). Changes in the amount of snow or when the snow melts could lead to shifts in the dominance of herbaceous species or invasion of trees or shrubs.

C3. Restricted to uncommon landscape/geological features: Somewhat Increase.

In Pend Oreille County, *Salix pseudomonticola* is found on glacial drift material associated with the Metaline limestone and dolomite, a formation found sporadically in northeastern Washington. The reported occurrence from Chelan County is found on the Chiwaukum Schist, a formation of limited extent in the Wenatchee Range.

C4a. Dependence on other species to generate required habitat: Neutral.

The fen and wet meadow habitat occupied by *Salix pseudomonticola* is maintained primarily by natural abiotic processes.

C4b. Dietary versatility: Not applicable for plants

C4c. Pollinator versatility: Neutral.

Salix inflorescences lack showy petals or sepals and are capable of wind pollination. Flowers also produce nectar and floral scents to attract small insect pollinators, especially flies, bees, and butterflies.

C4d. Dependence on other species for propagule dispersal: Neutral.

Willow seeds have a tuft of wavy, silky hairs and are dispersed passively by wind.

C4e. Sensitivity to pathogens or natural enemies: Neutral.

Although willows are susceptible to rust fungi, no impacts to *Salix pseudomonticola* are known. This species may be vulnerable to browsing.

C4f. Sensitivity to competition from native or non-native species: Somewhat Increase.

Salix pseudomonticola could be sensitive to competition from other plant species if its wetland habitat becomes drier due to drought or reduced snowpack and water recharge under future climate change (Rocchio and Ramm-Granberg 2017).

C4g. Forms part of an interspecific interaction not covered above: Neutral.

Does not require an interspecific interaction.

C5a. Measured genetic variation: Unknown.
Data are not available on the genetic diversity of this species in Washington.

C5b. Genetic bottlenecks: Unknown.

C5c. Reproductive System: Neutral

Salix pseudomonticola is dioecious (with separate staminate and pistillate individuals) and is thus an obligate outcrosser. Pollination can occur by insects or long-distance dispersal by wind. Seed dispersal occurs by wind. The life history of this species suggests that it should have average genetic diversity across populations. The occurrences in Washington are near the edge of the species range and are likely to have slightly lower genetic diversity due to founder effects or inbreeding depression.

C6. Phenological response to changing seasonal and precipitation dynamics: Neutral.
No changes have been detected in phenology in recent years.

Section D: Documented or Modeled Response to Climate Change

D1. Documented response to recent climate change: Neutral.
No recent changes in the distribution of this species in Washington have been detected.

D2. Modeled future (2050) change in population or range size: Unknown

D3. Overlap of modeled future (2050) range with current range: Unknown

D4. Occurrence of protected areas in modeled future (2050) distribution: Unknown

References

Camp, P. and J.G. Gamon, eds. 2011. Field Guide to the Rare Plants of Washington. University of Washington Press, Seattle. 392 pp.

Fertig, W. and J. Kleinknecht. 2020. Conservation status and protection needs of priority plant species in the Columbia Plateau and East Cascades ecoregions. Natural Heritage Report 2020-02. Washington natural Heritage Program, WA Department of Natural Resources, Olympia, WA. 173 pp.

Rocchio, F.J. and R.C. Crawford. 2015. Ecological systems of Washington State. A guide to identification. Natural Heritage Report 2015-04. Washington Natural Heritage Program, WA Department of Natural Resources, Olympia, WA. 384 pp.

Rocchio F.J. and T. Ramm-Granberg. 2017. Ecological System Climate Change Vulnerability Assessment. Unpublished Report to the Washington Department of Fish and Wildlife. Washington Natural Heritage Program, Department of Natural Resources, Olympia, WA.

Young, B.E., E. Byers, G. Hammerson, A. Frances, L. Oliver, and A. Treher. 2016. Guidelines for using the NatureServe Climate Change Vulnerability Index. Release 3.02. NatureServe, Arlington, VA. 48 pp. + app.