

# METAL MINES OF WASHINGTON — PRELIMINARY REPORT

by  
ROBERT E. DERKEY  
NANCY L. JOSEPH  
and  
RAYMOND LASMANIS

WASHINGTON DIVISION OF GEOLOGY AND EARTH RESOURCES  
OPEN FILE REPORT 90-18

NOVEMBER 1990

---

This report has not been edited or reviewed for conformity with  
Division of Geology and Earth Resources standards and nomenclature.

---



WASHINGTON STATE DEPARTMENT OF  
**Natural Resources**

Brian Boyle - Commissioner of Public Lands  
Art Stearns - Supervisor

Division of Geology and Earth Resources  
Raymond Lasmanis, State Geologist



# **METAL MINES OF WASHINGTON — PRELIMINARY REPORT**

by

**ROBERT E. DERKEY**

**NANCY L. JOSEPH**

and

**RAYMOND LASMANIS**

**WASHINGTON DIVISION OF GEOLOGY AND EARTH RESOURCES**

**OPEN FILE REPORT 90-18**

**NOVEMBER 1990**

---

This report has not been edited or reviewed for conformity with  
Division of Geology and Earth Resources standards and nomenclature.

---





# CONTENTS

	Page
Introduction . . . . .	1

County and site name	Site number	
<b>BENTON</b>		
Berrian Island placer	303	5
Gone Busted placer	304	6
<b>CHELAN</b>		
Blewett Iron	562	7
Blewett camp	557	8
Cannon	558	9
Crown Point	564	11
Dick	565	12
Holden	555	13
Lovitt	566	14
Negro Creek Iron	563	16
Negro Creek placers	650	17
Peshastin Creek placers	561	18
Red Mountain	556	19
Rex	559	20
<b>CLALLAM</b>		
Cedar Creek placer	615	21
Crescent	607	22
Helen	610	24
Hurricane	608	25
Littleton	611	26
Ozette Beach placer	612	27
Shi Shi Beach placer	613	28
Sunset Creek placer	614	29
Victor	609	30
<b>FERRY</b>		
Republic district	- 2	31
Addison	308	32
Alva Stout placer	344	33
Belcher	353	34
Ben Hur	314	35
Black Tail	315	36
Blue Horse	359	37
Boston & New York	360	38
California	316	39
Dan Patch	361	40
El Caliph	317	41
Flag Hill	318	42
Golden Cord	362	43
Golden Harvest	319	44
Gray	363	45
Gwin	364	46
Hellgate Bar placer	345	47
Ida May	320	48
Insurgent	321	49
Jennie	322	50
Johnson placer	346	51

	Page
Kelly Camp	368 . . . . .52
Kettle	339 . . . . .53
Key East	309 . . . . .54
Knob Hill	323 . . . . .55
Lancaster	310 . . . . .57
Last Chance	324 . . . . .58
Little Cove	325 . . . . .59
Lone Pine	326 . . . . .60
Lone Star and Washington	311 . . . . .61
Longstreet	305 . . . . .62
Messenger	356 . . . . .63
Meteor	365 . . . . .64
Minnehaha	312 . . . . .65
Morning Glor	327 . . . . .66
Morning Star	328 . . . . .67
Mount Tolman	307 . . . . .68
Mountain Lion	329 . . . . .70
North San Poil Fraction	330 . . . . .71
Old Hickory	331 . . . . .72
Overlook	355 . . . . .73
Oversight	354 . . . . .74
Pearl	332 . . . . .75
Plum Bar placer	347 . . . . .76
Princess Maude	333 . . . . .77
Quilp	334 . . . . .78
Republic	335 . . . . .79
Rogers Bar placer	348 . . . . .80
San Poil	336 . . . . .81
Seattle	337 . . . . .82
Shamrock	358 . . . . .83
Silver Bell	357 . . . . .84
Silver Leaf	306 . . . . .85
Singer placer	349 . . . . .86
South Penn	338 . . . . .87
Stray Dog	366 . . . . .88
Surprise	340 . . . . .89
Talisman	313 . . . . .90
Threemile placer	350 . . . . .91
Tom Thumb	341 . . . . .92
Trade Dollar	342 . . . . .93
Valley	343 . . . . .94
Whitestone placer	351 . . . . .95
Wilmont Bar placer	352 . . . . .96
Zalla M	367 . . . . .97
<b>GRANT</b>	
Chinaman Bar placer	302 . . . . .98
<b>GRAYS HARBOR</b>	
Burnt Peak	624 . . . . .99
Egge	620 . . . . .100
Elma	623 . . . . .101
Esther-Irene	621 . . . . .102
Skunk Creek No. 19	622 . . . . .103

		Page
<b>JEFFERSON</b>		
Elkhorn	616	104
<b>KING</b>		
Anderson	600	105
Apex	595	106
Carmack	597	107
Cleopatra group	602	108
Clipper	605	109
Condor-Hemlock	606	111
Coney Basin	598	113
Damon and Pythias	596	114
Dutch Miller	592	115
Grand Central	590	116
Great Republic	591	117
Guye	601	118
Quartz Creek	599	119
Royal Reward	604	120
Seattle-Cascade	603	121
Snoqualmie	593	122
Una	594	123
<b>KITTITAS</b>		
Boulder Creek	567	124
Boulder Creek placer	584	125
Cascade Chief	572	126
Clarence Jordin	573	127
Cle Elum River Iron	581	128
Dolphin	570	129
Esther and Louisa	583	130
Flodine	576	131
Francis Virdin Park	588	132
Golden Fleece	577	133
H-O-M-E	582	134
Huckleberry	569	135
Liberty	575	136
Mineral Creek	571	137
Mount Hawkins	568	138
Mountain Daisy	579	139
Old Bigney placer	585	140
Ollie Jordan	574	141
Silver Creek	580	142
Silver Tip	589	143
Swauk Creek placers	586	144
Wall Street	578	145
Williams Creek placers	587	146
<b>LEWIS</b>		
Barnum-McDonnell	632	147
Eagle Peak	631	148
Lytle-Lynch	634	149
Mineral Creek	635	150
Roy	633	151
<b>LINCOLN</b>		
Barnell placer	298	152
China Bar placer	299	153
Clark placer	300	154
Keller Ferry placer	301	155
Spokane Molybdenum	297	156

		Page
<b>MASON</b>		
Black Hump	619	157
Black and White	617	158
Triple Trip	618	160
<b>OKANOGAN</b>		
49th Parallel	369	161
Adelia	375	162
Alder	452	163
American Flag	376	164
American Flag	446	165
American Rand	407	166
Andy O'Neal	473	167
Antimony Bell	441	168
Antimony Queen	442	169
Apache	474	170
Arlington	420	171
Bales	443	172
Bellevue	377	173
Bi-Metallic	472	174
Black Bear	378	175
Blue Lake	447	176
Bodie	379	177
Bullfrog	380	179
Bunker Hill	381	180
Butcher Boy	382	181
Caribou	383	182
Casimer Bar placer	487	183
Castle Creek	466	184
Central	475	185
Chicago	384	186
Chloride Queen	385	187
Copper Glance	448	188
Copper World Extension	449	189
Crounse placer	488	190
Crystal Butte	386	191
First Thought	421	192
Four Metals	467	193
Fourth of July	476	194
Friday	453	195
Fuller	465	196
Gold Axe	387	197
Gold Crown	388	198
Gold Key	454	199
Golden Chariot	370	200
Golden Zinc	389	201
Gray Eagle	390	202
Gray Eagle no. 2	391	203
Hiawatha	392	204
Hidden Treasure	455	205
Highland	393	206
Holden-Campbell	394	207
Homestake	414	208
Horn Silver	422	209
Independence	456	210
Ivanhoe	423	211
John Judge	396	212
Johnson Creek	445	213

		Page
<b>OKANOGAN (continued)</b>		
Kaaba	415	214
Kankakee	450	215
Kelsey	438	216
Key	424	218
Kimberly	416	219
King Solomon	435	220
Lakeview	395	221
Last Chance	477	222
Leuena	425	223
Lilman	478	224
Little Chief	479	225
Lone Star	468	226
Lucky Knock	444	227
Magnetic	412	228
Mammoth	426	230
Mary Ann Creek placer	489	231
Mazama	439	232
Methow	397	233
Mid Range	457	234
Minnie	398	235
Mohawk	436	236
Montana	451	237
Mountain Beaver	458	238
Mountain Boy	469	239
Mountain Sheep	428	240
Murray placer	490	241
Nevada	429	242
Nighthawk	417	243
O.K.	371	244
Okanogan Free Gold	399	245
Palmer Summit	400	246
Panama	480	247
Paymaster	459	248
Peacock	481	249
Pinnacle	401	250
Pittsburg	482	251
Pogue Flat	471	252
Poland China	402	253
Poorman	485	254
Prize	418	255
Rainbow	403	256
Ramore	470	257
Reco	404	258
Red Shirt	460	259
Review	405	260
Roosevelt	413	261
Ruby	430	262
Security	372	263
Sheridan	431	264
Shotwell placer	491	265
Silver Bell	406	266
Silver Bluff	432	267
Silver Cliff	483	268
Silver Ledge	462	269
Silver Mountain	433	270
Similkameen placers	437	271
Spokane	463	272
St. Anthony	461	273

		Page
Starr	440	274
Sullivan	464	276
Summit	419	277
Tough Nut	434	278
Triune	408	279
Twin Pine	486	280
Walker placer	492	281
War Eagle	409	282
Western Star	373	283
Wheeler	427	284
Whitestone	410	285
Wolframite	484	286
Wyandotte	411	287
Yakima	374	288
<b>PEND OREILLE</b>		
Metaline district	1	289
Bella May	120	291
Blue Bucket	121	292
Calispell	101	293
Cliff	122	294
Comstock	108	295
Diamond R.	109	296
Grandview	123	297
Hanley	124	298
Harvey Bar placer	102	299
Highnoon	135	300
Hoage	125	301
Josephine	126	302
King Tut	110	303
Kootenai Conquest	111	304
Lead Hill	127	305
Lead King	112	306
Lead Queen	113	307
Lehigh No. 1	106	308
Lehigh No. 2	107	309
Lucky Strike	128	310
Metaline	129	311
Molybdenite Mountain	119	312
Oriole	130	313
Pend Oreille	131	314
Red Top	114	316
Ries	115	317
Schierding placer	103	318
Schultz placer	104	319
Sterling	132	320
Sullivan	116	321
Sullivan Creek		
placer	105	322
Washington	117	323
West Contact	134	324
Yellowhead	133	325
Z Canyon	118	326
<b>PIERCE</b>		
Silver Creek	625	327
Silver Creek Gold		
& Lead	626	328
Silver Creek placer	627	329

		Page
<b>SKAGIT</b>		
Hamilton	523	330
Johnsburg	524	331
Mount Vernon	525	332
Ready Cash	522	333
Stephens	527	334
Willis and Everett	526	335

<b>SKAMANIA</b>		
Black Jack	638	336
Camp Creek placer	637	337
Margaret	639	338
Sweden	636	340

<b>SNOHOMISH</b>		
Bonanza Queen	528	341
Buckeye	529	342
Copper Belle	530	343
Ethel	531	344
Florence Rae	532	345
Forty Five	551	346
Glacier Peak	554	347
Helena	533	349
Horseshoe Bend placer	547	350
Index Gold	543	351
Iowa	534	352
Jefferson	549	353
Justice	544	354
Kromona	535	355
Lake Serene	536	356
Lockwood Pyrtie	550	357
Mackinaw	537	358
Monte Cristo	545	359
O and B	538	360
Rainy	546	361
Silverton mines	539	362
St. Louis and Jackson	552	363
Sultan King	540	364
Sultan placer	548	365
Sunrise	553	366
Sunset	541	368
Wayside	542	369

<b>SPOKANE</b>		
C. I. Smith	296	370
Dahl	295	371
Daybreak	294	372
Herem-Moore	291	373
Huffman	292	374
Silver Hill	293	375

<b>STEVENS</b>		
A and C	182	376
Admiral	144	377
Admiral Consolidated	273	378
Aichan Bee	142	379
Amazon	183	380
Anaconda	150	381
Antelope	207	382
B and B	184	383
Bechtol	149	384
Beecher	208	385

		Page
Big Chief	152	386
Big Iron	232	387
Big Smoke	270	389
Black Rock	275	390
Blue Bar placer	221	391
Bonanza	153	392
Bonanza Copper	185	393
Brooks	241	394
Bullion	289	395
Burrus	276	396
Calhoun	274	397
Chewelah Standard	186	398
Chinto	187	399
Chloride Queen	154	400
Clara	209	401
Cleveland	136	402
Columbia River	188	404
Copper Butte	189	405
Copper King	155	407
Copper King	190	408
Coyote	290	409
Daisy	242	410
Deep Creek	277	411
Deer Trail	137	412
Deer Trail Monitor	239	413
Double Eagle	156	414
Eagle	243	415
Easter Sunday	210	416
Edna	145	417
Electric Point	157	418
Eureka	211	419
Evans placer	222	420
Evergreen	158	421
F. H. and C.	212	422
Farmer	278	423
Finley	159	424
First Thought	213	425
Frisco Standard	244	427
Galena Hill	160	428
Galena Knob	161	429
Gem	214	430
Germania		
Consolidated	138	431
Germania	140	432
Gibson Bar placer	223	433
Gladstone	162	434
Gold Bar	215	435
Gold Ledge	216	436
Gold Reef	217	437
Gray Eagle	202	438
Great Western	163	439
Hidden Treasure	218	440
Homestake	219	441
Hope and Twin Cabins	191	442
Hubbard	164	443
Iroquois	279	444
Jackson	280	445
Jay Dee	246	446

		Page
<b>STEVENS (continued)</b>		
Jay Gould	247	.447
John Day	248	.448
Juno-Echo	192	.449
Keystone	234	.450
Krug	193	.451
Kulzer	235	.452
Lakeview	245	.453
Last Chance	165	.454
Lead Trust	166	.455
Liberty Copper	147	.456
Little Mountain	265	.457
Lone Star	249	.458
Longshot	167	.459
Loon Lake Blue Bird	194	.460
Loon Lake Copper	195	.461
Lotie	196	.462
Lowley	272	.463
Lucile	281	.464
Lucky Four	250	.465
Magma	282	.466
Maki	168	.467
McGrath placer	224	.468
McNally	197	.469
Melrose	251	.470
Middleport	283	.471
Midnite	272	.472
Minorca	198	.474
Morning	252	.475
Mountain View	253	.476
Myerah	169	.477
Napoleon	236	.478
Neglected	170	.479
Negro Creek Bar placer	225	.480
Nevada	171	.481
New Blue Grouse	263	.482
New Leadville	172	.483
Nobles placer	226	.484
O-Lo-Lim	146	.485
Old Blue Grouse	264	.486
Old Dominion	254	.487
Ora	199	.489
Orazada	255	.490
Orchid	256	.491
Ore Cache	173	.492
Plata Rica	257	.493
Pohle placer	227	.494
Providence	258	.495
Queen	141	.496
R. J.	174	.497
Rambler	200	.498
Read	237	.499
Red Cloud	201	.500
Red Top	175	.501
Reed and Roberts placer	228	.502
Roosevelt	259	.503
Sand Creek	266	.504

		Page
Scaman	176	.505
Scandia	284	.506
Scotia	203	.507
Sherwood	148	.508
Shoemaker	285	.510
Sierra Zinc	286	.511
Silver Crown	177	.512
Silver Queen	240	.513
Silver Summit	178	.514
Silver Trail	179	.515
Sterrett	180	.516
Sunday	220	.517
Superior Copper	204	.518
Tempest	260	.519
Tenderfoot	151	.520
Thompson	238	.521
Togo	139	.522
Tungsten King	267	.523
Tungsten Products	268	.524
Turk	205	.525
Tyee	233	.526
U. S. Copper Gold	261	.527
Uncle Sam	181	.528
United Copper	206	.529
United Treasure	262	.530
Valbush Bar placer	229	.531
Van Stone	287	.532
Van Stone placer	230	.534
Washington Metals	269	.535
Wells Fargo	143	.536
Weston placer	231	.537
Young America	288	.538
<b>WHATCOM</b>		
Allen Basin	496	.540
Anacortes	497	.541
Azurite	498	.542
Blonden	505	.543
Boundary Red Mountain	499	.544
Breckenridge Creek	515	.545
Chancellor	500	.546
Danny	493	.547
Evergreen	501	.548
Farrar Placer	518	.549
Fourth of July	502	.550
Gargett	503	.551
Glacier	517	.552
Goat	504	.553
Gold Basin	506	.554
Gold Hill	516	.555
Golden Arrow	513	.556
Great Excelsior	507	.557
Lazy Tar Heel Placer	519	.559
Lone Jack	508	.560
Mammoth	509	.561
Minnesota	512	.562
New Light	510	.563
North American	511	.564
Old Discovery Placer	520	.565

	Page		Page
<b>WHATCOM (continued)</b>		<b>YAKIMA</b>	
Ribbon	494 . . . . . 566	Chinook	630 . . . . . 570
Scougale Placer	521 . . . . . 567	New Find	628 . . . . . 571
Silver Tip	495 . . . . . 568	Red Bird	629 . . . . . 572
Whistler	514 . . . . . 569		

## APPENDIX

Listing of alternate site names with primary site name, site number, and county . . . . .	573
---	-----

# Metal Mines of Washington—Preliminary Report

by Robert E. Derkey, Nancy L. Joseph, and Raymond Lasmanis

This report is a preliminary compilation that summarizes the geology and related features of 541 metal mines in Washington. This volume is an updated revision of producing mining properties described by Huntting (1956) in "Inventory of Washington Minerals—Part II, Metallic Minerals". For a mine to be included in this data set, it should have produced about \$1,000 worth of ore. Much of the production information included here is from Huntting (1956); however, production statistics, which are commonly incomplete in Huntting, are not significantly improved in this report.

In essence, this is a working document; the database will be continuously updated as new information is received. If a mine in which you are interested has produced metal(s) worth at least \$1,000 but is not included here, please let us know. If you can supply appropriate production data for this mine, we will include the mine in the database and future versions of this report. In addition, if you find errors in this report or have new information about any included mine, please contact us. Send your comments or additions to:

Robert E. Derkey  
Washington Division of Geology and Earth Resources  
MS PY-12  
Olympia, WA 98504

or

Nancy L. Joseph  
Washington Division of Geology and Earth Resources  
Spokane County Agricultural Center  
N. 222 Havana  
Spokane, WA 99202

The information in this report can be made available on disc. Contact R. E. Derkey for more information.

## EXPLANATION OF INFORMATION

This inventory of Washington mines is a collection of geologic and mineral deposit information that has been entered into a computer database program called GSMODS (Johnson, 1987). All accumulated information for a single mine constitutes one record in the database. A category within a record, such as "ore controls", is termed a field. Two types of fields make up each record. The first type requires entry of names or abbreviations, such as for "ore minerals" or "commodities", and results in a listing. The other type of field allows insertion of an unlimited amount of text or data, including pertinent information from literature sources, the reference, and pages cited.

Each mine or occurrence (record) in the database is intended to stand alone with respect to its content. Consequently, some fields in a record contain the same data as those fields for other nearby mines. The repeated data normally pertain to or describe the geology of the area as a whole. For example, the text of "tectonic setting" may be the same for all mines in a given mining district.

We have attempted to make individual category/field entries as complete as possible. However, where information is unavailable, the field is left blank. For some mines, information from literature sources is entered as a direct quotation, especially in unlimited-length text fields.

**Site Name** - The first entry of a site's record is the best known or most recent name of the mine, mineral deposit, or mining district. This is referred to as the primary site name. Sites are presented in alphabetical order by primary site name within each county (also presented in alphabetical order). The primary site name is accompanied by a unique site number which will be used to identify mines on map plots of this inventory data set.

**Alternate Names** - All other names by which the mine has been known. See the Appendix at the back of this report for a listing of these alternate names, along with the primary site name, site number and county.

**District** - The mining district(s) in which the mine is situated

**County** - The county in which the mine is located (See Fig. 1.)

**Primary Quad** - The 7.5- and/or 15-minute quadrangle(s) in which the mine is located

**Scale** - Map scales for the primary quadrangles, presented in the same order as the primary quadrangles, that is, the first map scale listed is that for the first primary quadrangle in the list

**1/2° x 1° Quad** - The 1/2 x 1 degree quadrangle in which the mine is located

**1° x 2° Quad** - The 1 x 2 degree quadrangle in which the mine is located

**Latitude** - The latitude of the mine location. This information is generally obtained by digitizing from the primary quadrangle(s).

**Longitude** - The longitude of the mine location. This information is generally obtained by digitizing from the primary quadrangle(s).




Figure 1. The counties of Washington.

**Section, Township, and Range** - The section, township, and range location for each mine. For many mines, the point selected is the center of a cluster of adits and/or shafts. If the location of a principal adit or shaft is known, that point is entered. If a mine portal is distant from an underground deposit, then a point directly over the deposit is selected.

**Location** - Any information that will assist in location of the mine. Some of this information may refer to features that are not shown of a topographic map but that may be helpful.

**Host Rock: Name** - The geological units, sequences, or formations in which the mineralization occurs. This information is generally taken from available geologic maps.

**Lithology** - The host rock lithology. This is given in the same order as that for the host rock names. The lithology, if taken from the explanation of a geologic map, may differ from lithologies given in the Ore Controls field for a mine. The lithology as recorded in Ore Controls is commonly from an

individual description of the mine, whereas the description of the host formation lithology is commonly a general description from a geologic map.

**Age** - Host rock age. Entries here correspond sequentially to the listing of host rock names.

**Associated Igneous Rock: Description** - A brief description of igneous rocks genetically associated with the mineralization. Because the age of relation of igneous rock to mineralization at a mine is not always known, all igneous rock near a mine site may be included in this entry. This entry is a list if two or more igneous units are associated with the mineralization.

**Age** - The age of the associated igneous rock(s). This may be a listing, corresponding sequentially to the listing of the associated igneous rocks.

**Commodities** - The known commodities at the mine. The chemical symbols for elements and acronyms for commodities are listed at the top of the facing page.

**Ore Minerals** - All ore minerals known to occur at the mine. The minerals are listed in generally decreasing order of



## Commodity symbols and acronym

Ag - silver	Ni - nickel
As - arsenic	Os - osmium
Au - gold	P - phosphorus
Be - beryllium	Pb - lead
Bi - bismuth	PGE - platinum group elements
Cd - cadmium	Pt - platinum
Ce - cerium	S - sulfur
Co - cobalt	Sb - antimony
Cr - chromium	Se - selenium
Cu - copper	Sn - tin
Fe - iron	Te - tellurium
Ga - gallium	Th - thorium
Ge - germanium	Ti - titanium
Ir - iridium	U - uranium
Hg - mercury	V - vanadium
Mg - magnesium	W - tungsten
Mn - manganese	Zn - zinc
Mo - molybdenum	

abundance. This listing may be incomplete even though the commodities are known.

**Non-Ore Minerals** - A brief listing of the types of hydrothermal alteration, gangue minerals, and other non-ore minerals associated with mineralization at the mine.

**Deposit Type** - Concise labels that describe the type(s) of mineral deposit(s) at the site

**Mineralization Age** - The age of the mineralization

**Production** - The production history of the mine from various published sources, including Huntting (1956). This information is incomplete, but for many deposits, it is the best currently available.

**Tectonic Setting** - A brief description of the tectonic setting or geologic conditions at the time mineral deposition took place. In many instances, a general tectonic history of an area is included because the age of mineralization is unknown.

**Ore Controls** - A description of the factors controlling mineralization, including structural, stratigraphic, chemical, or other controls.

**Geologic Setting** - The general geology of the mine. This information is expanded from the host rock description and is commonly taken from geologic maps. Map scale and detail vary, and the data in this entry for some mines is very general.

**Comments** - A field for entry of miscellaneous information that does not conveniently fit another field. Some development history and property characteristics are included here.

**Unpublished Information** - A notation of informal, unpublished information in the Division of Geology and Earth Resources (DGER) files or library is given here.

**References** - Selected literature used as sources from which to compile the data for the mine, as well as other information that may be of interest to the user. Acronyms used in some entries for "Production" and references: USGS, U.S. Geological Survey; MRDS, Mineral Resource Data System, a national database of the USGS.

## ACKNOWLEDGMENTS

The authors appreciate the numerous discussions with colleagues concerning the geology of areas surrounding the mineral deposits. We also gratefully acknowledge reviews by William Phillips, Hank Schasse, Eric Schuster, and Kieth Stoffel of DGER. The illustration was prepared by David Clark.

## REFERENCES CITED

- Huntting, M. T., 1956, Inventory of Washington Minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v: 1, 428 p.; v. 2, 67 p.
- Johnson, B. R., 1987, GSMODS - A personal mineral occurrence database system; Reference manual, Version 1.01: U.S. Geological Survey Open-File Report 87-636, 225 p.



***Berrian Island placer*** (303)

ALTERNATE NAMES		DISTRICT	COUNTY
Goody placer			Benton
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Umatilla	1:24,000		Pendleton
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
45° 56' 49.00" N	119° 15' 21.53" W	sec. 1, 5N, 28E, and sec. 6, 5N, 29E	
LOCATION: on the north bank of the Columbia River, elev. 300 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: The plant operated for 6 weeks, and concentrate was shipped to the Tacoma smelter in 1949 (Hunting, 1956, p. 181).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

***Gone Busted placer*** (304)

ALTERNATE NAMES		DISTRICT	COUNTY
			Benton
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Blalock Island	1:24,000		Pendleton
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
45° 53' 29.19" N	119° 39' 16.29" W		
LOCATION: on Blalock Island in the Columbia River near Patterson			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: A dry-land washing plant operated from 1938 to 1940 (Hunting, 1956, p. 181).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Blewett Iron (562)**

ALTERNATE NAMES		DISTRICT	COUNTY
Blewett Washington Nickel		Blewett	Chelan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Blewett	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 23' 32.52" N	120° 39' 29.46" W	secs. 13 and 14, 22N, 17E	
LOCATION: on a prominent ridge north of Shaser Creek at its junction with Peshastin Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	serpentine	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Ni Cr	hematite magnetite chromite nickel silicates	serpentine, clay-rich residuum of laterites	

<b>DEPOSIT TYPE</b>	<b>MINERALIZATION AGE</b>
laterite residual concentration	
<b>PRODUCTION:</b> The Blewett Iron deposit has not produced but has been extensively evaluated as an iron resource (Hunting, 1956, p. 194).	
<b>TECTONIC SETTING:</b> Several iron-nickel-chromite lateritic deposits overlie serpentinized ultramafic rocks (Broughton, 1943, p. 9-13) of the Ingalls Complex, a Jurassic ophiolite complex (Miller, 1985, p. 27; Tabor and others, 1982, p. 5-6).	
<b>ORE CONTROLS:</b> The Blewett iron deposits are laterite deposits. Many of the laterite deposits were eroded from their nearby source area and deposited to form a sequence of conglomerates and fine-grained, iron-rich beds. The iron and chromium were from iron in silicate minerals, magnetite-hematite, and chromite weathered from the serpentine. Nickel is from silicates released during laterization (deep weathering) of the serpentine (Broughton, 1943, p. 10).	
<b>GEOLOGIC SETTING:</b> The laterites were transported and deposited beneath or as basal beds of the Eocene Swauk Formation. Lamey (1950, p. 1) tentatively interprets the conglomeratic iron beds as landslide debris or mudflow deposits.	
<b>COMMENTS:</b> Iron deposits of the Blewett area were evaluated in the 1940s by the Washington Division of Geology. This was followed by extensive drilling by the U.S. Bureau of Mines (10 holes, 2,395 ft) and metallurgical testing. (See references for specific reports.) This evaluation showed reserves of 46,000 tons at a grade of 0.88% Ni, 2.5% Cr <sub>2</sub> O <sub>3</sub> , 32% Fe, and 8,000,000 tons at a grade of 0.39% Ni, 0.85% Cr <sub>2</sub> O <sub>3</sub> , and 11.53% Fe (Hunting, 1956, p. 194).	

**REFERENCES**

- Broughton, W. A., 1943, The Blewett iron deposit, Chelan County, Washington (with preliminary tonnage estimates): Washington Division of Geology Report of Investigations 10, 17 p., 1 pl.
- Broughton, W. A., 1944, Economic aspects of the Blewett-Cle Elum iron ore zone, Chelan and Kittitas Counties, Washington: Washington Division of Geology Report of Investigations 12, 42 p., 7 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Lamey, C. A., 1950, The Blewett iron-nickel deposit, Chelan County, Washington: U.S. Geological Survey Bulletin 969-D, p. 87-103, 2 pl.
- Lupher, R. L., 1944, Stratigraphic aspects of the Blewett-Cle Elum iron ore zone, Chelan and Kittitas Counties, Washington: Washington Division of Geology Report of Investigations 11, 63 p., 2 pl.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Blewett camp** (557)

ALTERNATE NAMES		DISTRICT	COUNTY
Blewett, Peshastin, Culver, Alta Vista, Black Jack, La Rica, Manistee, Phipps, Pole Pick #1, Prospect, Wilder, Blue Bell, Eureka, Fraction, Lucky Queen, Golden Eagle, Hummingbird, North Star, Olden, Sandell, Tip Top		Blewett	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Blewett	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 25' 29.46" N	120° 39' 35.02" W	SW1/4 sec. 1, 22N, 17E	
LOCATION: near Peshastin Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	serpentine	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	native gold	pyrite, arsenopyrite, quartz, calcite	
Ag	native copper		
Hg	chalcopryite		
Cu	malachite		
	galena		
	stibnite		
	cinnabar		
	magnetite		
	hematite		
	chromite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Production records for the various mines of the Blewett camp are incomplete; however, the value of production totals several hundred thousand dollars. Many of the mines have some recorded production, but a production figure for the entire camp is not available (Hunting, 1943, p. 9-14; 1956, p. 109-117).

**TECTONIC SETTING:** The Jurassic Ingalls Complex, the host rocks of the Blewett Camp deposits, is a disrupted, Late Jurassic ophiolite complex. Igneous activity in the area includes the Late Cretaceous Mount Stuart batholith and intrusive rocks of early magmatic activity of the Cascades magmatic arc (Miller, 1985, p. 27; Tabor and others, 1982, p. 5-6).

**ORE CONTROLS:** Where the veins cut gneissic rocks, the gangue is chiefly quartz with minor amounts of calcite, but where the veins cut serpentinite, the amount of calcite increases relative to quartz. The most productive veins are in Culver Gulch and trend N75W. The ore is generally of low grade. Most production was from high-grade ore shoots (Hunting, 1943, p. 8).

**GEOLOGIC SETTING:** Most of the mineralization of the Blewett Camp is in veins cutting serpentinite of the Ingalls Complex (Hawkins Formation in Weaver, 1911, p. 31-34). Other veins are in metasedimentary and metavolcanic rocks of the Complex; however, veins in the serpentinite are generally more productive than those in the other rocks (Weaver, 1911, p. 29-33).

**COMMENTS:** Mines of the Culver Gulch area are here combined into a single entry called the Blewett Camp deposit.

**REFERENCES**

- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.
- Weaver, C. E., 1911, Geology and ore deposits of the Blewett mining district: Washington Geological Survey Bulletin 6, 104 p.

**Cannon (558)**

ALTERNATE NAMES		DISTRICT Wenatchee	COUNTY Chelan
PRIMARY QUADRANGLE Wenatchee	SCALE 1:24,000	1/2° x 1° QUAD Wenatchee	1° x 2° QUAD Wenatchee
LATITUDE 47° 23' 44.24" N	LONGITUDE 120° 19' 30.55" W	SECTION, TOWNSHIP, AND RANGE NE1/4SW1/4 sec. 22, 22N, 20E	
LOCATION: on the west side of Squillchuck Creek.			
HOST ROCK: NAME Chumstick Formation unnamed sedimentary rocks	LITHOLOGY conglomerate, sandstone, siltstone sandstone, mudstone, siltstone, conglomerate	AGE middle Eocene Eocene?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION porphyry of Wenatchee dome and Rooster Comb, a flow-layered rhyodacite dome Saddle Rock andesite		AGE Eocene Eocene	
COMMODITIES Au Ag	ORE MINERALS electrum gold pyrargyrite naumannite acanthite agularite chalcopryrite stibnite sphalerite galena hessite	NON-ORE MINERALS pyrite, arsenopyrite, marcasite, quartz, chalcedony, adularia, calcite. Hydrothermal alteration, particularly silicification is important in ore localization. Other alteration minerals include sericite, argillic minerals, potassium silicates, and chlorite. Mineralization is found in zones of pervasive and vein-controlled silicification and potassium silicate alteration. Intermediate argillic alteration zones cap the mineralization. Mixed layer illite/smectite grades outward into smectite.	
DEPOSIT TYPE vein stockwork disseminated		MINERALIZATION AGE Tertiary	

**PRODUCTION:** The Cannon mine began production in the mid 1980s and is producing at the present time (1990). Production for 1989 was 150,420 oz of Au and 302,731 oz of Ag (Joseph, 1990, p. 19).

**TECTONIC SETTING:** The Cannon mine is situated in the Chiwakum graben (Eocene), a north-northwest-trending right-lateral, strike-slip graben bounded by the Entiat fault zone on the east and the Leavenworth fault zone on the west (Ott and others, 1986, p. 426).

**ORE CONTROLS:** Mineralization consists of widely spaced veins, veinlets (stockworks), and disseminated minerals in brecciated rocks. By far the most abundant ore mineral is electrum. The highest ore grades are in quartz-chalcedony-adularia-calcite veins. Two sets of fractures control vein mineralization: early veins in fractures radial to fold hinges, and veins in faults that cut the earlier radial faults-veins (Ott and others, 1986, p. 425, 432). Electrum in banded veins is found near the vein margin, and pyrite, where present, is found near the center of the vein. The mineralization occurs in sandstone and siltstone beds that were silicified prior to brecciation and deposition of quartz veins and veinlets. The quartz veins range to as much as 4 ft wide and are irregular and discontinuous. Most veinlets are less than 1 in. wide.

**GEOLOGIC SETTING:** An unnamed sandstone unit and the overlying Eocene Chumstick Formation host mineralization at the Cannon mine. Mineralization occurs in favorably altered (commonly silicified) horizons of these Tertiary (Eocene?) arkosic sandstone units (Ott, 1988, p. 20-23). Ore bodies extend from the surface to depths of more than 900 ft.

**COMMENTS:** This deposit is currently (1990) being mined as a joint venture project of Asamera Minerals Inc. and Breakwater Resources Ltd.

**UNPUBLISHED INFORMATION:** Information gained from a tour of the Cannon mine lead by mine geologist M. Mehlhorn on July 25, 1990, is supplemented the published accounts of the Cannon mine geology.

**REFERENCES**

Gresens, R. L., 1983, Geology of the Wenatchee and Monitor quadrangles, Chelan and Douglas Counties, Washington: Washington Division of Geology and Earth Resources Bulletin 75, 75 p., 3 pl.

## Chelan

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., 1990, Mineral industry news notes: Washington Geologic Newsletter, v. 18, no. 2, p. 19-20.
- Ott, L. E., 1988, Economic geology of the Wenatchee mining district, Chelan County, Washington: University of Idaho Doctor of Philosophy thesis, 270 p., 2 pl.
- Ott, L. E.; Groody, Diane; Follis, E. L.; Siems, P. L., 1986, Stratigraphy, structural geology, ore mineralogy and hydrothermal alteration at the Cannon mine, Chelan County, Washington, U.S.A. *In* McDonald, A. J., editor, Gold '86—An international symposium on the geology of gold deposits—Proceedings volume: Gold '86 [Toronto, Ont.], p. 425-435.
- Patton, T. C.; Cheney, E. S., 1971, L-D gold mine, Wenatchee, Washington—New structural interpretation and its utilization in future exploration: Society of Mining Engineers of AIME Transactions, v. 250, no. 1, p. 6-11.



**Crown Point (564)**

ALTERNATE NAMES		DISTRICT	COUNTY
Aurelia Crown Crown Power		Railroad Creek	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Suiattle Pass	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 12' 6.97" N	120° 53' 46.03" W	NE¼ sec. 8, 31N, 16E	
LOCATION: at the head of the cirque basin southwest of Hart Lake at the head of Railroad Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Cloudy Pass pluton		laboradorite granodiorite	early Miocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mo Cu	molybdenite chalcopyrite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced from 1897 to 1902, 10 tons in 1901 and 12 tons in 1902 (Hunting, 1956, p. 268).

TECTONIC SETTING: The Cloudy Pass pluton was emplaced during early Miocene igneous activity of the Cascades magmatic arc.

ORE CONTROLS: Molybdenite occurs as large crystals in a flat-lying quartz vein which ranges from 3 in. to 3 ft thick (Hunting, 1956, p. 268).

GEOLOGIC SETTING: The quartz vein occurs in laboradorite granodiorite of the early Miocene Cloudy Pass Pluton (Cater and Crowder, 1967, geol. map).

## REFERENCES

- Cater, F. W.; Crowder, D. F., 1967, Geologic map of the Holden quadrangle, Snohomish and Chelan Counties, Washington: U.S. Geological Survey Geologic Quadrangle Map GQ-646, 1 sheet, scale 1:62,500.
- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

# Chelan

## Dick (565)

ALTERNATE NAMES		DISTRICT	COUNTY
Chelan Winesap		Entiat	Chelan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Winesap	1:24,000	Chelan	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 46' 11.86" N	120° 12' 23.36" W	S½NE¼ sec. 9 26N, 21E	
LOCATION: Oklahoma Gulch area			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed peridotite	peridotite	pre-Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ni Cu	pyrrhotite pentlandite chalcopyrite malachite nickel sulfate	peridotite	
DEPOSIT TYPE		MINERALIZATION AGE	
magmatic segregation		pre-Jurassic	

PRODUCTION: Four short adits on the property and seven drill holes totaling 1,016 ft were completed by the USBM (Hunting, 1956, p. 275).

TECTONIC SETTING: The area is underlain by migmatite of the Chelan Complex (Tabor and others, 1987, geol. map).

ORE CONTROLS: Disseminated and massive sulfide segregations in peridotite (Hunting, 1956, p. 275; 1943, p. 25-26). Oxidation extends to about 40 ft depth. The peridotite body is about 400 ft long and 100 ft wide at the surface (Hunting, 1956, p. 275).

GEOLOGIC SETTING: Creasey and Storch (1945, geol. map) show the Dick deposit in a peridotite body between a quartz diorite dike and granite gneiss complex.

### REFERENCES

- Creasey, S. C.; Storch, H. H., 1945, Winesap nickel prospect, Chelan County, Washington: U.S. Geological Survey Open-File Report, 10 sheets.
- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Whetten, J. T.; Waitt, R. B.; Swanson, D. A.; Byerly, G. R.; Booth, D. B.; Hetherington, M. J.; Zartman, R. E., 1987, Geologic map of the Chelan 30-minute by 60-minute quadrangle, Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1661, 1 sheet, scale 1:100,000, with 29 p. text.

**Holden (555)**

ALTERNATE NAMES		DISTRICT	COUNTY
Howe Sound Irene			Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Holden	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 11' 47.50" N	120° 46' 48.58" W	secs. 18 and 19, 31N, 17E; secs. 12 and 13, 31N, 16E	
LOCATION: near the town of Holden on Railroad Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
rocks of the Napeequa River area	hornblende schist and gneiss, sericite-quartz schist	Late Paleozoic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au Zn Ag	chalcopyrite sphalerite gold galena molybdenite	pyrite, pyrrhotite, magnetite, arsenopyrite, quartz, sericite	
DEPOSIT TYPE		MINERALIZATION AGE	
metamorphosed massive sulfide		Permian	

**PRODUCTION:** From 1938 through 1957 a total of more than 212 million lb of Cu, 40 million lb of Zn, 2 million oz of Ag, and 600,000 oz of Au were obtained from 10 million tons of ore (McWilliams, 1958, p. 3).

**TECTONIC SETTING:** The deposit formed in an island arc to back arc basin environment with deposition of both mafic and siliceous volcanic rocks. Nonvolcanic rocks, including marble, are also prevalent near the mine (Nold, 1983, p. 945-948).

**ORE CONTROLS:** The orebody is lens shaped and has a length downdip of about 2,000 ft and a thickness of as much as 100 ft. The zoned orebody ranges from a zinc-rich footwall to a copper-rich hanging wall, indicating the orebody is overturned. Foliation of the schist parallels bedding (Nold, 1983, p. 945-948).

**GEOLOGIC SETTING:** The mine occurs within a series of schists, gneisses, and amphibolites, which are part of Cater and Crowder's (1967, geol. map; Cater, 1982, p. 6-7) Late Paleozoic "rocks of the Napeequa River area". The orebody is surrounded by sericite-quartz schist occurring within the schists, gneisses, and amphibolite sequence (Nold, 1983, p. 944-948). The Holden mine is believed to be Kuroko-type, volcanogenic massive sulfide deposit (R. E. Derkey). Kuroko-type features include the zoned, massive-sulfide orebody (zoned with copper-rich base and zinc-rich top) conformable with bedding, enclosure in sericite schist (altered felsic tuffs), and presence of anhydrite.

**REFERENCES**

- Cater, F. W., 1982, Intrusive rocks of the Holden and Lucerne quadrangles, Washington—The relation of depth zones, composition, textures, and emplacement of plutons: U.S. Geological Survey Professional Paper 1220, 108 p.
- Cater, F. W.; Crowder, D. F., 1967, Geologic map of the Holden quadrangle, Snohomish and Chelan Counties, Washington: U.S. Geological Survey Geologic Quadrangle Map GQ-646, 1 sheet, scale 1:62,500.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- McWilliams, J. R., 1958, Mining methods and costs at the Holden mine, Chelan Division, Howe Sound Co., Chelan County, Wash.: U.S. Bureau of Mines Information Circular 7870, 44 p.
- Nold, J. L., 1983, The Holden mine, a metamorphosed volcanogenic deposit in the Cascade Range of Washington: Economic Geology, v. 78, no. 5, p. 944-953.
- Youngberg, E. A.; Wilson, T. L., 1952, The geology of the Holden mine: Economic Geology, v. 47, no. 1, p. 1-12.

**Lovitt (566)**

ALTERNATE NAMES		DISTRICT	COUNTY
Golden King Wenatchee Squillchuck Gold King L-D		Wenatchee	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Wenatchee	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 22' 55.43" N	120° 18' 54.75" W	NE1/4SW1/4 sec. 22, 22N, 20E	
LOCATION: on the west side of Squillchuck Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Chumstick Formation unnamed unit	conglomerate, sandstone, and siltstone sandstone, mudstone, siltstone, conglomerate	middle Eocene Eocene?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
porphyry of Wenatchee dome and Rooster Comb, a flow-layered rhyodacite dome		Eocene	
Saddle Rock andesite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	electrum native gold pyrargyrite naumannite acanthite aguilarite chalcopryrite stibnite sphalerite galena hessite	pyrite, arsenopyrite, marcasite, quartz, chalcedony, adularia, calcite, siderite. Hydrothermal alteration, particularly silicifi- cation, is important in ore localization. Other alteration minerals include sericite, argillic minerals, potassium silicates, and chlorite.	
DEPOSIT TYPE		MINERALIZATION AGE	
vein stockwork disseminated		Tertiary	

**PRODUCTION:** Production in 1894, 1910, 1938-39 and 1944-46 (Hunting, 1956, p. 113). Continuous production from 1949 to 1967, when 1,036,572 tons of ore yielded 410,482 oz Au (0.396 oz/ton) and 625,849 oz Ag (0.60 oz/ton) (Ott, 1988, p. 10).

**TECTONIC SETTING:** The Lovitt mine is in the Chiwakum graben (Eocene), a north-northwest-trending right-lateral, strike-slip graben bounded by the Entiat fault zone on the east and the Leavenworth fault zone on the west (Ott and others, 1986, p. 426).

**ORE CONTROLS:** Mineralization is disseminated and in stockworks in pervasively altered and silicified sandstone and siltstone. Two sets of fractures control vein mineralization; early veins in fractures radial to fold hinges, and veins in faults that cut the earlier radial faults-veins (Ott and others, 1986, p. 425, 432).

**GEOLOGIC SETTING:** An unnamed sandstone unit and the overlying Eocene Chumstick Formation host mineralization at the Lovitt and nearby Cannon mines. Mineralization occurs in favorably altered (commonly silicified) horizons of these extensively deformed and brecciated arkosic sandstone units (Ott, 1988, p. 20-23).

**COMMENTS:** The Lovitt is part of the property controlled by the Cannon mine, which is currently (1990) being mined as a joint venture project of Asamera Minerals Inc. and Breakwater Resources Ltd.

**REFERENCES**

- Gresens, R. L., 1983, Geology of the Wenatchee and Monitor quadrangles, Chelan and Douglas Counties, Washington: Washington Division of Geology and Earth Resources Bulletin 75, 75 p., 3 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Lovitt, E. H.; Skerl, A. C., 1958, Geology of the Lovitt gold mine, Wenatchee, Wash.: Mining Engineering, v. 10, no. 9, p. 963-966.

- Ott, L. E., 1988, Economic geology of the Wenatchee mining district, Chelan County, Washington: University of Idaho Doctor of Philosophy thesis, 270 p., 2 pl.
- Ott, L. E.; Groody, Diane; Follis, E. L.; Siems, P. L., 1986, Stratigraphy, structural geology, ore mineralogy and hydrothermal alteration at the Cannon mine, Chelan County, Washington, U.S.A. *In* McDonald, A. J., editor, Gold '86—An international symposium on the geology of gold deposits—Proceedings volume: Gold '86 [Toronto, Ont.], p. 425-435.
- Patton, T. C.; Cheney, E. S., 1971, L-D gold mine, Wenatchee, Washington—New structural interpretation and its utilization in future exploration: Society of Mining Engineers of AIME Transactions, v. 250, no. 1, p. 6-11.

**Negro Creek Iron (563)**

ALTERNATE NAMES		DISTRICT	COUNTY
Davenport		Blewett	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Blewett	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 24' 27.60" N	120° 44' 20.76" W	S1/2SW1/4 and NE1/2SE1/4 and NW1/4 SW1/4 sec. 8, 22N, 17E and SE1/4SE1/4 sec. 12, 22N, 16E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	serpentine	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Ni Cr	magnetite chromite	serpentine, clay-rich residuum of laterites	
DEPOSIT TYPE		MINERALIZATION AGE	
laterite residual concentration		Eocene	

**PRODUCTION:** Broughton (1943, p. 16-19) reports 59,850 tons of exposed, 45,720 tons of probable, and 435,850 tons of possible ore reserves in three areas.

**TECTONIC SETTING:** Iron-nickel-chromium deposits (residuum of weathering products) overlie serpentinized ultramafic rocks (Broughton, 1943, p. 9-13) of the Ingalls Complex, a Jurassic ophiolite complex (Miller, 1985, p. 27; Tabor and others, 1982, p. 5-6).

**ORE CONTROLS:** The Negro Creek Iron deposit is one of several Blewett-area iron deposits. These deposits are a residuum of weathering products developed over iron-, chromium-, and nickel-bearing serpentinized peridotite. Iron and chromium are found in the minerals magnetite, hematite, and chromite weathered from the serpentinite. Nickel silicates develop during laterization (extended weathering) of the serpentinite. The iron beds occur in a series of thick conglomerates and associated rocks (Broughton, 1943, p. 10).

**GEOLOGIC SETTING:** Some of the laterites were reworked prior to deposition of the Eocene Swauk Formation. Lamey (1950, p. 1) tentatively interprets the conglomeratic iron beds as landslide debris or a mudflow deposit.

**REFERENCES**

- Broughton, W. A., 1943, The Blewett iron deposit, Chelan County, Washington (with preliminary tonnage estimates): Washington Division of Geology Report of Investigations 10, 17 p., 1 pl.
- Broughton, W. A., 1944, Economic aspects of the Blewett-Cle Elum iron ore zone, Chelan and Kittitas Counties, Washington: Washington Division of Geology Report of Investigations 12, 42 p., 7 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Lamey, C. A., 1950, The Blewett iron-nickel deposit, Chelan County, Washington: U.S. Geological Survey Bulletin 969-D, p. 87-103, 2 pl.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Negro Creek placers (560)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Blewett	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Blewett	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 26' 34.77" N	120° 39' 55.74" W	secs. 2 and 3, 22N, 17E	
LOCATION: from the mouth of Negro Creek upstream for several miles			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cr Pt	native gold chromite	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced \$100 prior to 1897 (Hunting, 1956, p. 181). Platinum was also found about 6 mi upstream from the mouth of Negro Creek. Only a small amount (reportedly) was produced (Hunting, 1956, p. 281).

TECTONIC SETTING: Alluvial deposits containing heavy minerals.

ORE CONTROLS: Heavy minerals were concentrated by stream action. Several pieces of nearly pure chromite weighing about 3 lb each were reported (Hunting, 1956, p. 37) from Negro Creek in the SW 1/4 sec. 12, 22N, 16E.

GEOLOGIC SETTING: Negro Creek drains an area containing several gold and chromite deposits of the Blewett Camp. Bedrock of the area includes ultramafic rocks of the Ingalls Complex (Tabor and others, 1982).

## REFERENCES

- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

***Peshastin Creek placers*** (561)

ALTERNATE NAMES		DISTRICT	COUNTY
		Blewett	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Blewett	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 24' 20.72" N	120° 39' 24.98" W	22N, 17E	
LOCATION: on the upper reaches of Peshastin Creek.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	native gold	sand and gravel	
DEPOSIT TYPE	MINERALIZATION AGE		
placer	Quaternary		

PRODUCTION: Huntting (1956, p. 181) reports the area has produced.

TECTONIC SETTING: Alluvial deposits containing heavy minerals.

ORE CONTROLS: Heavy-mineral concentration by stream action. Huntting (1956, p. 181) reports the gold is coarse.

GEOLOGIC SETTING: Division files contain a mineral examination report prepared for an application to patent placer claims in sec. 13, 22N, 17E. This property is near the mouth of Allen Creek, which appears to be the southern limit of placer gold on Peshastin Creek. The creek drains northward past several mines between Allen Creek and Negro Creek. This is the apparent extent of favorable ground for placer deposits on Peshastin Creek.

UNPUBLISHED INFORMATION: U.S. Forest Service, 1964, Report of Mineral Examination for claims on Peshastin Creek. This report is in DGER file - "Peshastin".

## REFERENCES

- Huntting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.



**Red Mountain (556)**

ALTERNATE NAMES		DISTRICT	COUNTY
Royal		Chiwawa	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Trinity	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 4' 31.83" N	120° 50' 55.75" W	secs. 15 and 22, 30N, 16E	
LOCATION: at the south end of Phelps Ridge			
HOST ROCK: NAME		LITHOLOGY	AGE
Swakane Biotite Gneiss unnamed diorite		biotite gneiss diorite	Precambrian - Paleozoic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au Zn Pb W	chalcopyrite sphalerite galena scheelite	pyrrhotite, pyrite, arsenopyrite, chlorite, quartz, calcite, biotite, sericite	
DEPOSIT TYPE		MINERALIZATION AGE	
breccia zone			

**PRODUCTION:** Produced more than 15,000 tons of ore during intermittent periods from 1930 to 1940 (Hunting, 1956, p. 50).

**TECTONIC SETTING:** Ore at the Red Mountain mine was emplaced during or following major deformation and metamorphism of the host rocks. The extent of metamorphism of the orebody, if any, is not reported (Cater and Crowder, 1967, geol. map).

**ORE CONTROLS:** Mineralization is most extensive in the brecciated zone between Swakane Biotite Gneiss and diorite. Sulfide minerals are chiefly pyrrhotite and chalcopyrite. Scheelite occurs as disseminated grains, veinlets, and small masses in the sulfides (Culver and Broughton, 1945, p. 14-15).

**GEOLOGIC SETTING:** Radiometric ages on zircons from the Swakane Biotite Gneiss indicate an age for metamorphism of 415 m.y. and formation ages of 1,650 m.y. or more (Mattinson, 1972, p. 3773). The age and nature of the diorite are unknown.

**REFERENCES**

- Cater, F. W., 1982, Intrusive rocks of the Holden and Lucerne quadrangles, Washington—The relation of depth zones, composition, textures, and emplacement of plutons: U.S. Geological Survey Professional Paper 1220, 108 p.
- Cater, F. W.; Crowder, D. F., 1967, Geologic map of the Holden quadrangle, Snohomish and Chelan Counties, Washington: U.S. Geological Survey Geologic Quadrangle Map GQ-646, 1 sheet, scale 1:62,500.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mattinson, J. M., 1972, Ages of zircons from the northern Cascade Mountains, Washington: Geological Society of America Bulletin, v. 83, no. 12, p. 3769-3783.

# Chelan

## Rex (559)

ALTERNATE NAMES		DISTRICT	COUNTY
Rogers		Entiat	Chelan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Ardenvoir	1:24,000	Chelan	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 42' 53.05" N	120° 16' 19.86" W	N1/2 sec. 36, 26N, 20E	
LOCATION: in the drainage of Crum Canyon, a tributary to the Entiat River			
HOST ROCK: NAME		LITHOLOGY	AGE
Chelan Complex		tonalite and tonalite gneiss	Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag		quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: More than \$170,000 by 1930 and small amounts in 1933, 1934, and 1940 (Hunting, 1956, p. 116).

TECTONIC SETTING: The Chelan Complex formed in Late Cretaceous and consists of high-grade metamorphic rocks that were intruded by numerous tonalitic plutons. These massive tonalite plutons were again metamorphosed in latest Cretaceous and earliest Tertiary time (Tabor and others, 1987, p. 5-6). The relation between metamorphism and emplacement of the quartz veins is unknown.

ORE CONTROLS: Two oxidized quartz veins, 3 to 12 in. in width, in decomposed gneiss (Hunting, 1956, p. 116).

GEOLOGIC SETTING: The rocks of the Chelan Complex (Hopson and Mattinson, 1971, p. 13; see also map of Tabor and others, 1987, p. 5-6) consist of massive and gneissic tonalite and migmatite near Chelan and along Lake Chelan and banded gneiss and banded migmatite gneiss along the Entiat River.

### REFERENCES

- Hopson, C. A.; Mattinson, J. M., 1971, Metamorphism and plutonism, Lake Chelan region, northern Cascades, Washington [abstract]: Geological Association of Canada, Cordilleran Section, Programme and Abstracts, p. 13.
- Hunting, M. T., 1943, Inventory of mineral properties in Chelan County, Washington: Washington Division of Geology Report of Investigations 9, 63 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Whetten, J. T.; Waitt, R. B.; Swanson, D. A.; Byerly, G. R.; Booth, D. B.; Hetherington, M. J.; Zartman, R. E., 1987, Geologic map of the Chelan 30-minute by 60-minute quadrangle, Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1661, 1 sheet, scale 1:100,000, with 29 p. text.

***Cedar Creek placer*** (615)

ALTERNATE NAMES		DISTRICT	COUNTY
Starbuck Placer			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Allens Bay	1:24,000	Cape Flattery	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 00' 53" N	124° 40' 37" W	E 1/2 sec. 18, 29N, 15W	
LOCATION: 10 mi north of La Push, near the mouth of Cedar Creek by the beach			
HOST ROCK: NAME		LITHOLOGY	AGE
beach sands		sand and gravel	Pleistocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pt	gold platinum ilmenite magnetite chromite zircon	sand	
DEPOSIT TYPE		MINERALIZATION AGE	
marine placer		Pleistocene	

PRODUCTION: Produced a reported \$5,000 in gold and 5 oz of platinum prior to 1917 (Hunting, 1956, p. 182).

TECTONIC SETTING: Uplifted beach terrace.

ORE CONTROLS: Heavy-mineral concentration by wave action. Gold and platinum occur in a 2-15-in.-thick layer of sand on the surface of a wave-cut bench in clay (Hunting, 1956, p. 182).

## REFERENCES

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

# Clallam

## Crescent (607)

ALTERNATE NAMES		DISTRICT	COUNTY
			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Muller	1:24,000	Port Angeles	Victoria
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 04' 38" N	123° 56' 33" W	near the line between secs. 23 and 24, 30N, 10W	
LOCATION: elev. 1,250 ft, about 1.25 mi west of Lake Crescent. The railroad crosses the property. Access to the property, about 0.5 mi north of US Highway 101, is by way of road beginning at the highway 1 mi east of Fairholm.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	basalt	Eocene	
Crescent Formation	red limestone	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
spilitic pillow basalts and associated pyroclastic rocks and volcanic breccias.		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	hausmannite	hematite	
Fe	bementite		
Hg	neotocite		
S	braunite		
Cu	cinnabar		
Zn	native copper		
P			
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

**PRODUCTION:** Produced more than 50,000 tons between 1923 and 1954 (Hunting, 1956, p. 254).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** Mineralization occurs in three lenses, and several reverse faults offset the ore bodies (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may have formed as disseminated bodies of replacement origin or as volcanogenic exhalative bodies (Sorem and Gunn, 1967).

**GEOLOGIC SETTING:** The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated limestones were deposited in deep water (Snively, 1987, p. 306-308).

### REFERENCES

- Brown, R. D., Jr.; Gower, H. D.; Snively, P. D., Jr., 1960, Geology of the Port Angeles-Lake Crescent area, Clallam County, Washington: U.S. Geological Survey Oil and Gas Investigations Map OM-203, 1 sheet, scale 1:62,500.
- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Pardee, J. T., 1927, Manganese-bearing deposits near Lake Crescent and Humptulips, Washington. In Contributions to economic geology (short papers and preliminary reports), 1927; Part I-Metals and nonmetals except fuels: U.S. Geological Survey Bulletin 795-A, p. 1-24.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.

- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. *In* Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Helen** (610)

ALTERNATE NAMES		DISTRICT	COUNTY
			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Muller	1:24,000	Port Angeles	Victoria
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48 05 35 N	123 59 56 W	near center, N1/2 sec. 21, 30N, 10W	
LOCATION: elev. 1905 ft, about 8.2 mi west-northwest of Fairholm, about 0.8 mi north of US Highway 101.			
HOST ROCK: NAME		LITHOLOGY	AGE
Crescent Formation		basalt	Eocene
Crescent Formation		limestone	Eocene
ASSOCIATED IGNEOUS ROCK: DESCRIPTION			AGE
Crescent Formation volcanic rocks			Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

**PRODUCTION:** Produced a 10-ton shipment in 1952 and some production in 1954 (data from U.S.G.S. MRDS, 1990).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** Mineralization occurs as lenses near the contact between basalt and limestone (data from U.S.G.S. MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

**GEOLOGIC SETTING:** Red argillaceous limestone lenses occur enclosed in basalt of the Crescent Formation. Within the limestone lenses, boulder-like masses of brown to black manganiferous minerals are found associated with nodules of ferruginous quartz. Two bementite ore bodies are 30 ft x 6 ft and 23 ft x 6 ft (data from U.S.G.S. MRDS, 1990). The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. The basalts are associated with pelagic limestones, indicating deposition in deep water (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Purdy, C. P., Jr., 1954, Directory of Washington mining operations, 1954: Washington Division of Mines and Geology Information Circular 23, 73 p.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Hurricane (608)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Elwha	1:24,000	Port Angeles	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 00' 23" N	123° 30' 12" W		
LOCATION: Unsurveyed, elev. 4,000 ft, about 2.7 mi east-southeast of Elwha ranger station and 0.3 mi east-northeast of Griff Peak			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	limestone	Eocene	
Crescent Formation	basalt (altered)	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
basalt (altered)		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn Fe	hausmannite bementite neotocite rhodonite	hematite	
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic			

PRODUCTION: Produced about 1,000 tons of ore (Hunting, 1956, p. 255).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Mineralization at the Hurricane deposit is found in lens-shaped bodies (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: Area is underlain by highly altered basalt containing intercalated pods or lenses of red limestone (data from USGS MRDS, 1990). The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones indicate deposition in deep water (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Littleton (611)**

ALTERNATE NAMES		DISTRICT	COUNTY
Peacock Johnnie M			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Snider Peak	1:24,000	Cape Flattery	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48 04 52 N	124 00 35 W	sec. 20, 30N, 10W	
LOCATION: elev. 1,400 ft, 5.1 mi east-northeast of Snider ranger station, 500 ft west of Littleton Creek; part of the Littleton group of claims, probably on the border of sec. 20 and 21.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	basalt	Eocene	
Crescent Formation	red limestone	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite neotocite hausmannite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic			

PRODUCTION: Produced in 1952 and 1953 (Huntting, 1956, p. 256).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Ore bodies are lens shaped. Small faults offset the ore bodies. Manganese minerals occur as oxides and silicates in red limestone which is interbedded in basalt (data from USGS MRDS, 1990).

GEOLOGIC SETTING: Lenses of manganese oxides and silicates occur in red limestone, which is interbedded in basalt. Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and occur in reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.



**Ozette Beach placer (612)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Cape Flattery	Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Ozette	1:24,000	Cape Flattery	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 12' 10" N	124° 41' 20" W	sec. 12, 31N, 16W	
LOCATION: 2 mi north of the mouth of the Ozette River			
HOST ROCK: NAME	LITHOLOGY	AGE	
beach sands	sand	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pt PGE	gold platinum	sand	
DEPOSIT TYPE		MINERALIZATION AGE	
beach placer		Quaternary	

PRODUCTION: Produced a small amount in the early 1900s (Pardee, 1929; Huntting, 1956, p. 183). Produced \$15,000 from Ozette and Shi Shi placers prior to 1904 (Huntting, 1956, p. 182).

TECTONIC SETTING: Uplifted beach terraces.

ORE CONTROLS: Concentration of heavy minerals by wave action.

GEOLOGIC SETTING: Beach deposit of gold and platinum concentrated along the surface of a wave-cut terrace in sandstone (data from USGS MRDS, 1990).

## REFERENCES

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mertie, J. B., Jr., 1969, Economic geology of the platinum metals: U.S. Geological Survey Professional Paper 630, 120 p., 1 pl.
- Pardee, J. T., 1929, Platinum and black sand in Washington. In Contributions to economic geology (short papers and preliminary reports) 1928; Part I—Metals and nonmetals except fuels: U.S. Geological Survey Bulletin 805, p. 1-15.

***Shi Shi Beach placer*** (613)

ALTERNATE NAMES		DISTRICT	COUNTY
Lovelace			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Makah Bay	1:24,000	Cape Flattery	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 15' 25" N	124° 41' 03" W	secs. 18, 19, and 30, 32N, 15W	
LOCATION: approximately 2.5 mi south of Portage Head, near mouth of Petroleum Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
bBeach sands		sand	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pt Ir Os	gold platinum magnetite ilmenite iridosmine zircon	sand	
DEPOSIT TYPE		MINERALIZATION AGE	
marine placer		Quaternary	

PRODUCTION: Produced before 1904 and, together with the Ozette placer, produced more than \$15,000. Concentrate ran 1,120 lb of ilmenite, 96 lb of zircon, \$558.09 per ton in gold, and \$20.45 per ton in platinum (Huntting, 1956, p. 182).

TECTONIC SETTING: Uplifted beach terraces.

ORE CONTROLS: Heavy-mineral concentration by wave action. Grains of platinum only about a quarter the size of gold grains; about 1% of the platinum grains are ferromagnetic (data from USGS MRDS, 1990).

GEOLOGIC SETTING: Sand deposit on a wave-cut terrace in sandstone is covered by a thin layer of heavy mineral-rich black sand and a 1-3-ft-thick layer of gravel and sand. Gold and platinum are found in the thin heavy-mineral sand layer and in cracks in the underlying rock.

## REFERENCES

- Day, D. T.; Richards, R. H., 1906, Useful minerals in the black sands of the Pacific slope. *In* U.S. Geological Survey, Mineral resources of the United States, calendar year 1905: U.S. Geological Survey, p. 1175-1258.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mertie, J. B., Jr., 1969, Economic geology of the platinum metals: U.S. Geological Survey Professional Paper 630, 120 p., 1 pl.
- Pardee, J. T., 1929, Platinum and black sand in Washington. *In* Contributions to economic geology (short papers and preliminary reports) 1928; Part I—Metals and nonmetals except fuels: U.S. Geological Survey Bulletin 805, p. 1-15.

**Sunset Creek placer (614)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
La Push	1:24,000	Forks	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 59' 20" N	124° 40' 17" W	sec. 19, 29N, 15W	
LOCATION: about 6 mi north of the mouth of the Quillayute River, near the mouth of Sunset Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
beach sands	sand	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pt PGE	gold platinum	sand	
DEPOSIT TYPE		MINERALIZATION AGE	
marine placer		Pleistocene	

PRODUCTION: Produced prior to 1917 (data from USGS MRDS, 1990).

TECTONIC SETTING: Uplifted beach terraces.

ORE CONTROLS: Heavy-mineral concentration by stream action. The metal-bearing part of the deposit is a thin layer of fine heavy-mineral sand composed chiefly of pink garnet and grains of ilmenite and magnetite. Minor amounts of chromite and some very small, clear crystals of zircon are also present (data from USGS MRDS, 1990).

GEOLOGIC SETTING: These Pleistocene deposits consist of stream gravel and drift. At this site the wave terrace is cut in a layer of indurated clay in the drift (data from USGS MRDS, 1990).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mertie, J. B., Jr., 1969, Economic geology of the platinum metals: U.S. Geological Survey Professional Paper 630, 120 p., 1 pl.
- Pardee, J. T., 1929, Platinum and black sand in Washington. *In* Contributions to economic geology (short papers and preliminary reports) 1928; Part I—Metals and nonmetals except fuels: U.S. Geological Survey Bulletin 805, p. 1-15.

**Victor (609)**

ALTERNATE NAMES		DISTRICT	COUNTY
Bear Creek Victory Black Bear			Clallam
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Deadman Hill	1:24,000	Cape Flattery	Cape Flattery
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 05' 36" N	124° 11' 31" W	NW1/4SW1/4 sec. 24, 30N, 12W	
LOCATION: elev. 1,800 ft, 2.25 mi northeast of Bear Creek (village). The Victor mine is on the steep southeast slope of Mount Muller Ridge; a 3.5-mi access road leads directly northward to the claims from Bear Creek on U.S. Highway 101.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation volcanic rocks	altered basalt	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
altered basalt		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	hausmannite bementite neotocite secondary manganese oxides		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic			

**PRODUCTION:** Produced 50 tons or more in 1953 (Hunting, 1956, p. 257). A 102-ton shipment of ore from the property assayed 48% Mn and 15% combined SiO<sub>2</sub> and Al<sub>2</sub>O<sub>3</sub> (data from USGS MRDS, 1990).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** The ore body occurs in lenses in a shear zone (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

**GEOLOGIC SETTING:** The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Republic district (2)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain Republic	1:24,000 1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
LOCATION: at the town of Republic			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, chalcedony	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: See description for individual mines.

TECTONIC SETTING: The Republic district is in the Republic graben, a major intracratonic, volcano-tectonic depression in northeastern Washington. The graben contains a thick sequence of Eocene volcanic and volcanoclastic rocks.

ORE CONTROLS: A major hydrothermal system in the Republic district produced a series of epithermal veins capped by hot-spring (sinter) deposits. The hot-spring deposits formed at paleosurfaces in the Sanpoil Volcanics prior to deposition of the overlying Klondike Mountain Formation. The majority of the recovered gold and silver was from the epithermal veins. The veins exhibit features indicative of episodic vein sealing and overpressuring followed by flashing and precipitation of precious metals, thus forming the characteristic coloform bands. Geometrically, ore deposits in the Republic district extend from the paleosurface mineralization in hydrothermally altered rocks and sinter downward into epithermal stockworks and hydrothermal breccias, and farther downward into tightly confined, commonly coloform-banded veins (Muessig, 1967; Tschauder, 1989).

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Full, R. P.; Grantham, R. M., 1968, Ore deposits of the Republic mining district, Ferry County, Washington. *In* Ridge, J. D., editor, Ore Deposits of the United States, 1933-1967; the Graton-Sales Volume: American Institute of Mining, Metallurgical, and Petroleum Engineers, v. 2, p. 1481-1494.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. *In* Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Addison (308)**

ALTERNATE NAMES		DISTRICT	COUNTY
Pacific Mutual			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Friedlander Meadows	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 3' 2.34" N	118° 34' 51.73" W	SE1/4 sec. 36, 30N, 33E, SW1/4 sec. 31, 30N, 34E, and NW1/4 sec. 6, 29N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	sandstone, greenstone, carbonate	Ordovician?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
porphyritic granodiorite of Manila Creek		Eocene - Paleocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Pb Zn Ag Au W	galena chalcopryrite chalcocite sphalerite malachite azurite scheelite wolframite lead carbonate	pyrite, quartz	

<b>DEPOSIT TYPE</b>	<b>MINERALIZATION AGE</b>
vein	
<b>PRODUCTION:</b> Produced a small amount of lead, copper, zinc, silver, and gold in 1923 (Hunting, 1956, p. 51). Produced 1,500 tons of ore in 1973 (Moen, 1976, p. 100).	
<b>TECTONIC SETTING:</b> Paleocene to Eocene granitic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map).	
<b>ORE CONTROLS:</b> Mineralization is in 4-6-ft-wide quartz veins in schist. The ore occurs as lenses in three of the exposed veins. One ore shoot is 50 ft long; another is 100 ft long (Hunting, 1956, p. 51).	
<b>GEOLOGIC SETTING:</b> Ordovician? sedimentary rocks of the Covada Group host quartz veins at the Addison mine. The sedimentary rocks are near Paleocene to Eocene granodiorite. Numerous Eocene hypabyssal dikes are found in the area (Joseph, 1990, geol. map, p. 6-7).	

**REFERENCES**

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.

***Alva Stout placer*** (344)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 38' 55.70" N	118° 46' 29.58" W	NW¼ sec. 2, 36N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaterna	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE	MINERALIZATION AGE		
placer	Quaternary		

PRODUCTION: Produced in 1934 (Huntting, 1956, p. 183).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Belcher (353)**

ALTERNATE NAMES		DISTRICT	COUNTY
Blue Bell-Belcher		Belcher	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooke Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 43' 26.36" N	118° 32' 46.02" W	NW1/4 sec. 8, 37N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	graywacke, argillite, chert, limestone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Cu Au	pyrite magnetite pyrrhotite chalcopyrite	garnet, tremolite, epidote	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic			

**PRODUCTION:** Considerable production in 1909-1911 and 1913-1917 (Hunting, 1956, p. 195).

**TECTONIC SETTING:** Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the Republic graben (Muessig, 1967, geol. map).

**ORE CONTROLS:** Two or three irregular replacement bodies are present in limestone in a sequence of contact-metamorphosed limestone and argillite cut by monzonite dikes. Iron sulfides are more abundant than magnetite (Hunting, 1956, p. 195).

**GEOLOGIC SETTING:** The Permian-Triassic rocks of the Belcher mine area are contact-metamorphosed limestone, dolomitic limestone, argillite, and graywacke. The ore bodies lie next to a network of dikes, sills, and intrusive bodies of Scatter Creek Rhyodacite (Muessig, 1967, p. 113).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.



**Ben Hur (314)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' 6.69" N	118° 45' 29.42" W	center of line between secs. 34 and 27, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, calcite, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** Produced \$65,000 up to 1910; produced during 1909-1915, 1918, 1933, 1949, and 1950 (Hunting, 1956, p. 118).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** Mineralization consists of a 4-ft-wide quartz vein in propylitic latite porphyry. The vein is composed of fine-grained banded quartz and 10%-30% calcite. The vein is said to extend the length of the claim (Hunting, 1956, p. 118). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Black Tail** (315)

ALTERNATE NAMES		DISTRICT	COUNTY
Hope		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 48.02".02 N	118° 44' 51.36" W	near E1/4 corner, sec. 34, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** Produced 300 tons prior to 1902; produced in 1909-1910 and 1912-1920 (Hunting, 1956, p. 118).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** Mineralization occurs in several 2-6-ft-wide quartz veins in quartz latite porphyry (dacite and andesite of recent authors) and propylitic andesite (Hunting, 1956, p. 118). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Blue Horse** (359)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Edds Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 33' 25.95" N	118° 35' 50.45" W	sec. 6, 35N, 34E	
LOCATION: on Iron Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	argillite, metawacke, greenstone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
diorite dike		pre-Tertiary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Au	arsenopyrite galena	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced prior to 1913 and in 1934 (Hunting, 1956, p. 288).

TECTONIC SETTING: Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the Republic graben (Muessig, 1967, geol. map).

ORE CONTROLS: A quartz vein roughly follows the contact between limestone-limy argillite and diorite (Hunting, 1956, p. 288).

GEOLOGIC SETTING: The Blue Horse mine is in a sequence of Permian-Triassic metasedimentary and metavolcanic rocks. Diorite adjacent to the vein contains thompsonite, which means it is probably pre-Tertiary in age (Muessig, 1967, geol. map, p. 17; Hunting, 1956, p. 288).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Boston & New York (360)**

ALTERNATE NAMES		DISTRICT	COUNTY
Welcome		Curlew	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Malo	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 51' 50.56" N	118° 34' 52.28" W	sec. 24, 39N, 33E	
LOCATION: The default location is to the center of the section. The mine may be the shaft and prospects shown in the northeast corner of the section on the 7.5' map.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Scatter Creek Rhyodacite	rhyodacite	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
diorite			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Pb	galena chalcopyrite		
DEPOSIT TYPE		MINERALIZATION AGE	

PRODUCTION: Produced in 1916 and 1928 (Hunting, 1956, p. 288).

TECTONIC SETTING: Tertiary intrusive rocks of the Republic graben (Parker and Calkins, 1964, geol. map).

ORE CONTROLS: No deposit data are available for this mine. It has produced Ag, Cu, and Pb, and galena and chalcopyrite are assumed.

GEOLOGIC SETTING: The mine is in the Scatter Creek Rhyodacite (Parker and Calkins, 1964, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Parker, R. L.; Calkins, J. A., 1964, Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1169, 95 p., 4 pl.

**California** (316)

ALTERNATE NAMES		DISTRICT	COUNTY
Apollo			Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Edds Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 36′ 9.61″ N	118° 34″ 54.48″ W	NW¼SW¼ sec. 20, 36N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	argillite, metawacke, greenstone	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	galena chalcopyrite sphalerite malachite azurite	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced in 1901-1902, 1908, 1914-1916, 1927-1929, and 1938-1939 (Hunting, 1956, p. 119).

**TECTONIC SETTING:** Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the area (Muessig, 1967, geol. map).

**ORE CONTROLS:** Mineralization is in a quartz vein along a fracture zone in greenstone and argillite (Hunting, 1956, p. 119).

**GEOLOGIC SETTING:** The California mine is in a sequence of Permian-Triassic rocks which are metasedimentary rocks near the deposit (Muessig, 1967, geol. map; Hunting, 1956, p. 119).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Dan Patch** (361)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Inchelium	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 16' 20.07" N	118° 12' 44.26" W	near center, SW¼ sec. 13, 32N, 36E	
LOCATION: near center, 3½ mi north of the Covada post office			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	wacke and quartzite	Ordovician?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Zn Au	galena sphalerite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: The mine produced in 1909.

TECTONIC SETTING: Metamorphosed Ordovician? rocks were intruded by Jurassic to Tertiary rocks (Joseph, 1990, geol. map, p. 21).

ORE CONTROLS: The deposit consists of three quartz veins cutting quartzite and slate. One vein is 16 in. wide and another 2 to 8 in. wide (Hunting, 1956, p. 289).

GEOLOGIC SETTING: The Dan Patch is in metamorphosed wacke and quartzite of the Ordovician? Covada Group (Joseph, 1990, geol. map, p. 21).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Weaver, C. E., 1913, Geology and ore deposits of the Covada mining district: Washington Geological Survey Bulletin 16, 87 p.

**El Caliph (317)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 16.52" N	118° 45' 36.44" W	S½ sec. 34, 37N, 32E	
LOCATION: just east of the Morning Glory property			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	pyrite, quartz, calcite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Estimated production of \$15,000-\$20,000 by the end of 1936; produced in 1916, 1933, 1934, and 1937-1939 (Hunting, 1956, p. 120).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Mineralization occurs in a 0.5-18-in.-wide vein cutting quartz latite and shale. The vein is displaced by minor faults (Hunting, 1956, p. 120). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Flag Hill (318)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 7.59" N	118° 45' 16.89" W	NW1/4NW1/4 sec. 1, 36N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Se	gold	quartz, calcite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Said to have produced 400 tons prior to 1940 (Hunting, 1956, p. 120).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Vein reported to be 5 ft wide and have an estimated length of 1,500 ft (Hunting, 1956, p. 120). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.



**Golden Cord** (362)

ALTERNATE NAMES		DISTRICT	COUNTY
		Keller	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Friedlander Meadows	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 3' 15.87" N	118° 35' 59.09" W	NW1/4SE1/4 sec. 36, 30N, 33E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	sandstone, greenstone, carbonate	Ordovician?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
porphyritic granodiorite of Manila Creek hypabyssal dacite dikes		Paleocene - Eocene Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Pb Zn	chalcopyrite sphalerite galena	pyrite, quartz, epidote, garnet	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Minor shipments were made in the early 1900s (Moen, 1976, p. 101).

TECTONIC SETTING: Paleocene to Eocene granitic rocks intrude metamorphosed Ordovician? rocks in the Keller graben (Joseph, 1990, geol. map, p. 12).

ORE CONTROLS: Mineralization is in silicified schist, which is cut by porphyry. Ore occurs as pockets and sparse disseminations in small quartz veinlets (Hunting, 1956, p. 290).

GEOLOGIC SETTING: Joseph (1990, geol. map) shows the mine is in porphyritic granodiorite of Manila Creek. Because Hunting (1956, p. 290) notes the mine is in schist, the area must be a small roof pendant in the Paleocene to Eocene granitic body. The area also contains numerous dacite dikes (Joseph, 1990, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Golden Harvest** (319)

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Bear Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 34' 39.01" N	118° 44' 54.26" W	near center sec. 36, 36N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	porphyritic dacite, andesite, and trachyte flows	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene	

PRODUCTION: Produced 800 tons of ore in 1937-1938 (Hunting, 1956, p. 120).

TECTONIC SETTING: Epithermal gold deposits are found in Eocene volcanic rocks of the Republic graben (Tschauder, 1989).

ORE CONTROLS: Mineralization is said to be in a 2.5-ft-wide vein that has an estimated length of 150 ft (Hunting, 1956, p. 120). The deposit is similar to epithermal deposits in the nearby Republic district.

GEOLOGIC SETTING: Mineralization at the Golden Harvest deposit is in the Eocene Sanpoil Volcanics (Muessig, 1967, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Gray** (363)

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Friedlander Meadows	1:24,000	Nespelem	Okanōgan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 2' 56.67" N	118° 34 40.31 W	NW1/4 sec. 6, 29N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	sandstone, greenstone, carbonate	Ordovician?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
porphyritic granodiorite of Manila Creek		Paleocene - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	galena		
Au	chalcopyrite		
Cu			
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Reportedly produced 140 tons of ore prior to 1940 (Huntting, 1956, p. 290).

TECTONIC SETTING: Paleocene to Eocene granitic rocks intrude metamorphosed Ordovician? rocks in the Keller graben (Joseph, 1990, geol. map, p. 12).

ORE CONTROLS: The ore is said to be present in an 8-in.-wide vein exposed for 600 ft (Huntting, 1956, p. 290). No ore minerals are reported from the Gray mine; however, it is near the Addison mine, which contains galena and chalcopyrite.

GEOLOGIC SETTING: Ordovician? sedimentary rocks host quartz veins at the Gray deposit. The sedimentary rocks are near Paleocene to Eocene granodiorite. Numerous Eocene hypabyssal dikes are found in the area (Joseph, 1990, geol. map, p. 6-7).

## REFERENCES

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.

**Gwin (364)**

ALTERNATE NAMES		DISTRICT	COUNTY
Guinn Hall Creek			Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Inchelium	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 17' 36.81" N	118° 14' 3.49" W	NW¼ sec. 11, 32N, 36E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	wacke and quartzite	Ordovician?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Au Pb W	tetrahedrite galena wolframite tennantite enargite malachite azurite	pyrite, arsenopyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced silver and some tungsten during the First World War (Hunting, 1956, p. 290).

**TECTONIC SETTING:** Host rocks are metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 21).

**ORE CONTROLS:** Quartz veins are as much as 4 ft wide and are parallel to the bedding in quartzite and argillite (Hunting, 1956, p. 290).

**GEOLOGIC SETTING:** The Gwin deposit is in wacke and sandstone of the Ordovician? Covada Group (Joseph, 1990, Geol. map, p. 21).

## REFERENCES

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Weaver, C. E., 1913, Geology and ore deposits of the Covada mining district: Washington Geological Survey Bulletin 16, 87 p.

***Hellgate Bar placer*** (345)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Whitestone Rock	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 56' .13" N	118° 36' 11.32" W	lot 5, sec. 13, 28N, 33E	
LOCATION: on the north side of the Columbia River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced a small amount in the early 1900s (Hunting, 1956, p. 183).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. A high bar contained gravel 2.5 to 3 ft thick showing values of 21.8 cents per cubic yard. A lower bar had 7 or 8 acres of gravel 2 to 3 ft thick averaging 27 cents per cubic yard. A low bench 2 mi long and 300 ft to 0.5 mi wide contained an iron-stained gold-bearing layer about 6 in. thick, 100 to 600 yd wide, and 2 mi long (Hunting, 1956, p. 183).

## REFERENCES

- Collier, A. J., 1907, Gold-bearing river sands of northeastern Washington. *In* Contributions to economic geology 1906; Part I—Metals and nonmetals, except fuels: U.S. Geological Survey Bulletin 315, p. 56-70.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Ferry****Ida May (320)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 17.27" N	118° 45' 52.72" W	SW¼ SW¼ sec. 34, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1914 (Hunting, 1956, p. 120).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Similar to ore deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Insurgent** (321)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' .82" N	118° 44' 46.16" W	NW1/4NW1/4 sec. 35, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1908-1912 and 1927 (Hunting, 1956, p. 121).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: The vein is thought to be an offshoot of the Lone Pine vein and cuts propylitic andesite. The ore shoot, which was 30 ft long and 2.5 ft wide, is now stoped out (Hunting, 1956, p. 121). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

# Ferry

## Jennie (322)

ALTERNATE NAMES		DISTRICT	COUNTY
Jenny Blue Mountain Patterson		Orient	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Laurier	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 55' 4.61" N	118° 13' 33.53" W	S 1/2 sec. 34, 40N, 36E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metamorphic rocks	schist, marble, quartzite	pre-Tertiary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Pb Zn	galena sphalerite	pyrite, quartz, gypsum	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

PRODUCTION: No large shipments were reported (Hunting, 1956, p. 121).

TECTONIC SETTING: The deposit is in the Kettle metamorphic core complex (Stoffel, 1990, p. 8).

ORE CONTROLS: Schist, marble, and quartzite intruded by lamprophyre dikes are traversed by a 35-ft-wide shear zone filled with brecciated country rock, gouge, quartz, and ore (Hunting, 1956, p. 121).

GEOLOGIC SETTING: The metamorphic rocks are part of the Kettle metamorphic core complex (Stoffel, 1990, p. 8).

### REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.



**Johnson placer** (346)

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Inchelium	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 17' 36.90" N	118° 9' 40.92" W	NE¼ sec. 8, 32N, 37E	
LOCATION: on the bank of the Columbia River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Reportedly produced \$100 (Hunting, 1956, p. 183).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. The paystreak consists of 1-3 ft of medium- to fine-textured gravel beneath 4-8 ft of sand (Hunting, 1956, p. 183).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Ferry**

**Kelly Camp (368)**

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bodie Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 48' 10.99" N	118° 47' 21.73" W	SW1/4 sec. 9, 38N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
metamorphic rocks of Tonata Creek	calc-silicate gneiss, schist, quartzite	pre-Tertiary	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
quartz monzonite, Herron Creek suite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
W Cu Mo	chalcopryrite scheelite magnetite molybdenite	garnet, epidote, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic			

PRODUCTION: Ten tons of ore were shipped for a mill test in 1951 (Hunting, 1956, p. 344).

TECTONIC SETTING: Eocene felsic rocks intruded pre-Tertiary metamorphic rocks adjacent to the Toroda Creek graben (Pearson, 1967, geol. map).

ORE CONTROLS: Mineralization is in a contact-metamorphic zone along the west side of a small roof pendant. The deposit contains a considerable amount of low-grade milling ore (Hunting, 1956, p. 344).

GEOLOGIC SETTING: Pre-Tertiary metamorphic rocks including calc-silicate gneiss, mica schist, and carbonaceous quartzite host contact metamorphic deposits of the Kelly Camp mine. These rocks are a roof pendant in quartz monzonite and monzonite of the Herron Creek suite (Pearson, 1967, geol. map, p. 3; Stoffel, 1990, geol. map)

**REFERENCES**

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pearson, R. C., 1967, Geologic map of the Bodie Mountain quadrangle, Ferry and Okanogan Counties, Washington: U.S. Geological Survey Geologic Quadrangle Map GQ 636, 1 sheet, scale 1:62,500, with 4 p. text.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.

**Kettle (339)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Vulcan Mountain	1:24,000	Republic	Okanogan
Curlew	1:24,000		
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 52' 43.51" N	118° 37' 31.54" W	near center sec. 15, 39N, 33E	
LOCATION: about 1.5 mi south of Curlew			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	pyrargyrite	quartz, calcite, pyrite, chalcedony, sericitic, propylitic, argillic, and quartz alteration	
Ag	electrum		
	gold		
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** The first gold-silver dore was poured (1990) from the jointly operated Overlook and Kettle mines. Anticipated production for the first year is 110,000 oz of Au, with the majority to come from the Overlook deposit (Joseph, 1990, p. 20).

**TECTONIC SETTING:** Epithermal deposits are found in the Sanpoil Volcanics and are similar to deposits of the Republic district.

**ORE CONTROLS:** The deposit exposed in the mine is in the zone just below the sinter zone of a typical hot-spring deposit. Veins are associated with some dikes (Walt Hunt, personal commun., 1990; Tschauder, 1989).

**GEOLOGIC SETTING:** Host rocks for the Kettle mine are the same as for deposits of the Republic district and are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Joseph, N. L., 1990, Mineral industry news notes: Washington Geologic Newsletter, v. 18, no. 2, p. 19-20.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Parker, R. L.; Calkins, J. A., 1964, Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1169, 95 p., 4 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Key East (309)**

ALTERNATE NAMES		DISTRICT	COUNTY
Copper Key		Belcher	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooke Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 42' 36.70" N	118° 33' 5.24" W	NE1/4 sec. 18, 37N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	graywacke, argillite, chert, limestone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Fe	pyrite pyrrhotite magnetite chalcopyrite	chlorite, sericite	
DEPOSIT TYPE		MINERALIZATION AGE	
manto-like massive replacement			

**PRODUCTION:** Shipped 3,249 tons of oxidized ore in 1907 and 7,000 tons of ore prior to 1940 (Hunting, 1956, p. 52).

**TECTONIC SETTING:** Permian-Triassic metasedimentary and metavolcanic rocks are preserved in the Republic graben east and northeast of Republic. They are locally cut by Eocene epizonal plutons and associated sills and dikes (Muessig, 1967, geol. map).

**ORE CONTROLS:** Iron-gold mineralization is manto-like and occurs as replacement bodies in limestone. One of these bodies is 100 ft long, 100 ft wide, and 25 ft thick. In places the ore bodies consist of nearly pure magnetite (Hunting, 1956, p. 52). Currently (1990), some of the iron bodies (magnetite, pyrite, pyrrhotite) are being mined for gold (Tschauder, 1989, p. 245).

**GEOLOGIC SETTING:** The Key East deposit is in Permian-Triassic limestone, dolomitic limestone, argillite, and graywacke. The ore bodies lie next to a network of dikes, sills, and intrusive bodies of the Scatter Creek Rhyodacite (Muessig, 1967, p. 113).

**COMMENTS:** The Key East deposit (formerly the Copper Key) and the nearby Key West deposit are identified areas that contain gold reserves. These deposits are about a mile northeast of the Overlook mine, which is now (1990) being mined by Echo Bay/Crown Resources.

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Knob Hill** (323)

ALTERNATE NAMES		DISTRICT	COUNTY
Golden Promise		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' 24.41" N	118° 45' 28.15" W	W1/2SE1/4 sec. 27, 37N, 32E	
LOCATION: at the head of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold electrum tellurides chalcopryrite stibnite realgar tetrahedrite polybasite pyrargyrite argentite umangite naumannite	quartz, chalcedony, pyrite, arsenopyrite, adularia, sericite, calcite, barite, graphite, marcasite; hydrothermal alteration is low- sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** Produced more than \$10,000,000 by the end of 1951 (Hunting, 1956, p. 121). On June 24, 1989, Hecla Mining Company celebrated the production of the 2 millionth ounce of gold from the Knob Hill No. 2 shaft (Lasmanis, 1989, p. 9).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** The deposit consists of several veins with mining widths of 5-15 ft (Hunting, 1956, p. 121). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

**REFERENCES**

- Full, R. P.; Grantham, R. M., 1968, Ore deposits of the Republic mining district, Ferry County, Washington. *In* Ridge, J. D., editor, Ore Deposits of the United States, 1933-1967; the Graton-Sales Volume: American Institute of Mining, Metallurgical, and Petroleum Engineers, v. 2, p. 148°1-1494.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Lasmanis, Raymond, 1989, Knob Hill No. 2 shaft at the Republic Unit produces 2 millionth ounce of gold: Washington Geologic Newsletter, v. 17, no. 3, p. 9-11.
- Lindgren, Waldemar; Bancroft, Howland, 1914, Republic (Eureka) district. *In* Bancroft, Howland, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, p. 133-166.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

## Ferry

Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. *In* Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Lancaster** (310)

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Curlew	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 54' 51.15" N	118° 33' 19.62" W	secs. 5 and 6, 39N, 34E	
LOCATION: 1 mi north of Curlew on the east bank of the Kettle River			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	argillite, siltstone, limestone	Permian - Triassic	
unnamed quartz monzonite	quartz monzonite	Eocene	
Scatter Creek Rhyodacite	rhyodacite	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
quartz latite dikes		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Pb Zn Au Ag	galena sphalerite chalcopyrite	pyrite, quartz, calc-silicates	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic disseminated			

PRODUCTION: Carload of 29 tons of hand-sorted ore shipped in 1929 (from open cuts) contained 1,651 lb Cu, 4,719 lb Pb, and 395 oz Ag (Parker and Calkins, 1964, p. 89).

TECTONIC SETTING: Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the area (Parker and Calkins, 1964, geol. map).

ORE CONTROLS: Chimney-like zones as much as 3 ft wide are present at the intersections of fractures in limestone at its contact with granite (Hunting, 1956, p. 54).

GEOLOGIC SETTING: A small body of Permian-Triassic, carbonate-bearing sedimentary rocks was intruded first by Eocene Scatter Creek Rhyodacite and then by Eocene quartz monzonite. All these rocks are cut by north-east-trending quartz latite dikes (Parker and Calkins, 1964, p. 89).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Parker, R. L.; Calkins, J. A., 1964, Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1169, 95 p., 4 pl.

**Last Chance** (324)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 56.93" N	118° 44' 46.69" W	W1/2NW1/4 sec. 35, 37N, 32E	
LOCATION: on the east side of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold tetrahedrite	quartz, calcite, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced \$3,000,000 by the end of 1923; produced again in 1940 (Hunting, 1956, p. 122).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: The vein in andesite flow breccia averages 8 ft in width. Vein filling consists of chalcedonic banded quartz, calcite, and fragments of country rock. Ore was largely removed above the 500-ft level (Hunting, 1956, p. 121). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.



**Little Cove (325)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' 4.44" N	118° 45' 16.97" W	NE1/4NW1/4 sec. 34, 37N, 32E	
LOCATION: adjoins the Pearl on the north			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	chalcedonic quartz, calcite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced two carloads of ore valued at \$1,450 prior to 1934; produced in 1934 and 1939-1940 (Hunting, 1956, p. 122).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: The deposit is similar to other deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Lone Pine (326)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 56.00" N	118° 44' 58.46" W	E1/2NE1/4 sec. 34, 37N, 32E	
LOCATION: on the east side of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, calcite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced \$137,000 worth of ore by 1910; produced again in 1935 (Hunting, 1956, p. 122).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Five veins in propylitically altered andesite are from 2 to 14 ft wide and consist of chalcedonic quartz traversed by narrow, black, crenulated ribbons. Most of the ore was above the 500-ft level and is stoped out (Hunting, 1956, p. 122). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Lone Star and Washington (311)**

ALTERNATE NAMES		DISTRICT	COUNTY
Lone Star			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Curlew	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 59' 47.36" N	118° 36' 9.55" W	SW1/4NE1/4 sec. 2, 40N, 33E	
LOCATION: adjacent to the international boundary			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	graywacke, argillite, chert, limestone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite alkalic? dike		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au Ag	chalcopyrite chalcocite malachite gold	pyrite, pyrrhotite, quartz, dolomite, calcite, sericite, chlorite, epidote, clinozoisite, serpentine, talc	
DEPOSIT TYPE		MINERALIZATION AGE	
disseminated and stockwork			

**PRODUCTION:** Produced in 1897 (1,700 tons) and in 1910-1917 (36,000 tons) (Hunting, 1956, p. 54).

**TECTONIC SETTING:** Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the area (Parker and Calkins, 1964, geol. map).

**ORE CONTROLS:** Chalcopyrite is present as disseminations and in veinlets along foliation in schistose, serpentinized dacite in a zone 50 ft wide in the hanging wall of an diabase dike dipping 30-50 degrees to the east and southeast. Company records indicated nearly 250,000 tons of ore containing 1.94% Cu, 0.047 oz/ton Au, 0.204 oz/ton Ag remained in the ground (Hunting, 1956, p. 54).

**GEOLOGIC SETTING:** The Lone Star and Washington mine is covered by a thick overburden of glacial debris, and few outcrops of Permian-Triassic greenstone, graywacke, argillite, and limestone can be found in the area. The diabase dike in the mine may be similar to alkalic rocks exposed on Shasket Creek (Parker and Calkins, 1964, p. 87).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Parker, R. L.; Calkins, J. A., 1964, Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1169, 95 p., 4 pl.

**Longstreet (305)**

ALTERNATE NAMES		DISTRICT Covada Meteor	COUNTY Ferry
PRIMARY QUADRANGLE Cedonia	SCALE 1:24,000	1/2° x 1° QUAD Nespelem	1° x 2° QUAD Okanogan
LATITUDE 48° 14' 16.56" N	LONGITUDE 118° 11' 59.36" W	SECTION, TOWNSHIP, AND RANGE NE1/4NE1/4 sec 36, 32N, 36E	
LOCATION:			
HOST ROCK: NAME granite to granodiorite near Meteor	LITHOLOGY granite and granodiorite	AGE Jurassic - Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION alaskite dike		AGE Cretaceous	
COMMODITIES Ag Sb Pb Zn Cu Au	ORE MINERALS stibnite argentite galena tetrahedrite sphalerite chalcopyrite	NON-ORE MINERALS quartz, pyrite, kaolinite	
DEPOSIT TYPE breccia zone		MINERALIZATION AGE	

**PRODUCTION:** Produced at least 120 tons of ore. A handpicked ore shipment ran 25% Sb. Three carloads shipped ran 53-57 oz/ton Ag and about 0.2 oz/ton Au (Hunting, 1956, p. 291).

**TECTONIC SETTING:** Jurassic to Cretaceous granitic to granodioritic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 34).

**ORE CONTROLS:** The deposit is a mineralized zone of silicified granodiorite breccia, 8 to 30 ft wide, that occurs adjacent to an alaskite dike. The breccia is cemented by quartz and ore minerals. Mineralization in the breccia zone is as much as 150 ft wide (Hunting, 1956, p. 291). Mineralization is accompanied by strong alteration and occurs on both sides of the alaskite dike. On the west side of the dike, mineralization is principally antimony and on the east side principally gold and silver (Purdy, 1951, p. 68-71).

**GEOLOGIC SETTING:** The Longstreet mine is in Jurassic to Cretaceous granite to granodiorite of the Meteor area (Joseph, 1990, geol. map, p. 34).

## REFERENCES

- Campbell, A. B.; Raup, O. B., 1964, Preliminary geologic map of the Hunters quadrangle, Stevens and Ferry Counties, Washington: U.S. Geological Survey Mineral Investigations Field Studies Map MF-276, 1 sheet, scale 1:48,000.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

**Messenger (356)**

ALTERNATE NAMES		DISTRICT	COUNTY
Big Joker		Covada	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cedonia	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 13' 38.73" N	118° 12' 12.72" W	SE¼ sec. 36, 32N, 36E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
granite to granodiorite near Meteor		granite and granodiorite	Jurassic - Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb Ag	galena	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced in 1934 (Hunting, 1956, p. 212).

**TECTONIC SETTING:** Jurassic to Cretaceous granitic to granodioritic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 34).

**ORE CONTROLS:** The ore is in several mineralized quartz veins from 12 to 26 in. wide in granodiorite near its contact with other rock (Hunting, 1956, p. 212).

**GEOLOGIC SETTING:** The granodiorite at the Messenger mine is Jurassic to Cretaceous and intrudes Ordovician? rocks of the Covada Group (Joseph, 1990).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Weaver, C. E., 1913, Geology and ore deposits of the Covada mining district: Washington Geological Survey Bulletin 16, 87 p.

**Meteor** (365)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Kewa	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 13' 43.73" N	118° 16' 45.56" W	north of center SW1/4 sec. 33, 32N, 36E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	wacke and quartzite	Ordovician?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	galena	pyrite, quartz	
Pb	sphalerite		
Au	rhodochrosite		
Cu	chalcopyrite		
Zn	cerargyrite		
	pyrargyrite		
	asrenopyrite		
	argentite		
	tetrahedrite		
	native silver		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced 20 tons of ore with a total gross value of \$1,000 by 1918 (Hunting, 1956, p. 292).

TECTONIC SETTING: Jurassic to Cretaceous intrusive rocks intrude Ordovician? sedimentary rocks of the area (Joseph, 1990, geol map).

ORE CONTROLS: The ore is in a 1 in.-1.5-ft-wide quartz vein in a shear zone in metamorphic rocks (Hunting, 1956, p. 292).

GEOLOGIC SETTING: The host rocks at the Meteor mine are of the Ordovician? Covada Group; they are metamorphosed wacke and quartz sandstone with slaty interbeds (Joseph, 1990, geol. map, p. 21).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Weaver, C. E., 1913, Geology and ore deposits of the Covada mining district: Washington Geological Survey Bulletin 16, 87 p.

**Minnehaha** (312)

ALTERNATE NAMES		DISTRICT	COUNTY
		Danville	Ferry
PRIMARY QUADRANGLE	SCALE	1½° x 1° QUAD	1° x 2° QUAD
Boundary Mtn	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 39.96" N	118° 27' 42.96" W	SE corner sec. 23, 40N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Permian metasedimentary rocks	phyllite and argillite	Permian	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite		
DEPOSIT TYPE		MINERALIZATION AGE	

PRODUCTION: Produced in 1903 and in 1924 (Hunting, 1956, p. 54).

TECTONIC SETTING: Permian metasedimentary rocks adjacent to rocks of the Republic graben (Stoffel, 1990, geol. map, p. 21).

ORE CONTROLS: No deposit data are available. Chalcopryrite is an assumed ore mineral because of the copper production.

GEOLOGIC SETTING: The Minnehaha deposit is in Permian phyllite and argillite. Eocene intrusive rocks crop out nearby (Stoffel, 1990, geol. map, p. 31).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.

**Morning Glory (327)**

ALTERNATE NAMES		DISTRICT	COUNTY
Old Glory		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATTITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 17.99" N	118° 45' 45.65" W	S1/2SW1/4 sec. 34, 37N, 32E	
LOCATION: on Flag Hill			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold tellurides	quartz, calcite, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced \$100,000 worth of ore prior to 1936; produced in 1937-1939 (Hunting, 1956, p. 122).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Vein of drusy banded quartz in quartz latite porphyry ranges from a few inches to 2 or 3 ft wide. Several rich pay shoots have been mined (Hunting, 1956, p. 122). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.



**Morning Star** (328)

ALTERNATE NAMES		DISTRICT	COUNTY
Lucile Dreyfus Faithful Surprise Mineral Hill Virginia			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Curlew	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 39.42" N	118° 30' 52.99" W	W1/2 SW1/4 SE1/4 sec. 16, 40N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metavolcanic rocks	greenstone	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu W	gold scheelite chalcopyrite	pyrite, pyrrhotite, quartz, serpentine	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced \$15,000 by 1910, \$27,000 in 1917, and about \$15,000 in 1935; also produced in 1936-1939 and 790 tons in 1940-1943 (Hunting, 1956, p. 122).

**TECTONIC SETTING:** Permian-Triassic rocks are in contact with serpentine (Parker and Calkins, 1964, geol. map).

**ORE CONTROLS:** The quartz veins cutting serpentine are said to average 2 ft in width (Hunting, 1956, p. 122). Veins are found in and near the sheared contact between greenstone and serpentinite. Gold-pyrite-quartz veins are prevalent in the greenstones, and pyrite-chalcopyrite-quartz veins are commonly found at the contact between serpentine and greenstone (Parker and Calkins, 1964, p. 89).

**GEOLOGIC SETTING:** Permian-Triassic greenstone is in sheared contact with serpentinite (Parker and Calkins, 1964, geol. map, p. 88-89).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Parker, R. L.; Calkins, J. A., 1964, Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1169, 95 p., 4 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Mount Tolman** (307)

ALTERNATE NAMES		DISTRICT	COUNTY
		Sanpoil Keller	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Keller	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 3' 24.74" N	118° 41' 31.53" W	secs. 31 and 32, 30N, 33E; N1/2 sec. 5 and NE1/4 sec. 6, 29N, 33E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Keller Butte pluton border phase	granodiorite and quartz diorite	Paleocene - Eocene Paleocene -	
Keller Butte pluton interior zone	quartz monzonite and granite	Eocene Paleocene - Eocene	
Keller Butte pluton interior zone	quartz porphyry		
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
hypabyssal complex (rhyolite, rhyodacite, andesite, and diorite) post-ore dikes		47-53 m.y.	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mo	molybdenite	pyrite, arsenopyrite, rhodochrosite, calcite,	
Cu	chalcopyrite	stilbite, laumontite, gypsum, limonite, quartz,	
Ti	rutile	sericite, muscovite, kaolinite, chlorite,	
Ag	chalcocite	K-spar, fluorite, topaz, garnet, magnetite,	
	copper	epidote, anhydrite	
	galena		
	sphalerite		
	wolframite		
	scheelite		
	malachite		
	chrysocolla		
DEPOSIT TYPE		MINERALIZATION AGE	
porphyry system fractures veins		50-60 m.y.	

**PRODUCTION:** There was an unknown amount of Pb, Cu, and Ag production from the Meadow Creek (Mount Tolman) mine in 1935 and 1936. The Golden Cord and the Ophir mines also produced, but no records are available.

**TECTONIC SETTING:** The Keller Butte pluton, which hosts the Mount Tolman deposit, is situated on the northwest projection of the N65-80W-trending Lewis and Clark structure (Osborn fault in Idaho). Foliation and mineral grains in the pluton trend N65-75W. Emplacement of the multi-phased pluton in this N65-80W trend resulted in the Cu-Mo deposit being elongate N68W.

**ORE CONTROLS:** The Mount Tolman deposit is 1 mi wide in a north-south direction, 2 mi long, and 1,200 ft thick. During crystallization of the last phase of the interior zone of the host quartz porphyry, a minimum of five periods of fracturing and concurrent mineralization took place. Each period affected the same rocks, and each episode of mineralization added to the distinctive vertical zonation: quartz with molybdenite predominate in the center, while chalcopyrite, pyrite, and sphalerite increase away from the core. Post-stockwork veins surrounding the Mount Tolman deposit cut the molybdenite-bearing fractures. These veins contain galena, sphalerite, and chalcopyrite. Drilling, underground exploration, and bulk sampling have outlined proven and probable reserves of 2.4 billion tons of ore with an average grade of 0.093% MoS<sub>2</sub> and 0.09% Cu; the cutoff grade used was 0.048% MoS<sub>2</sub>. Metallurgical testing resulted in recoveries of 85% of the molybdenite and 80% of the chalcopyrite. The deposit represents a recoverable reserve of 1.9 million tons of MoS<sub>2</sub> and 1.73 million tons of copper. A mine plan designed by AMAX would have recovered 727,260 tons of MoS<sub>2</sub> and 662,400 tons of copper. The AMAX plan was based on mining approximately 900 million tons of ore averaging 0.10% MoS<sub>2</sub> and 0.09% Cu and a waste to ore ratio of 0.83:1. This ranks the Mount Tolman deposit as the third or fourth largest molybdenum deposits in the world after Climax and Quartz Hill. Whole-rock analyses indicate the Mount Tolman rocks contain 0.3-0.5% TiO<sub>2</sub>, and if recoverable, the deposit could produce titanium.

**GEOLOGIC SETTING:** The Mount Tolman deposit is in the Paleocene-Eocene Keller Butte pluton. Post-ore faults displaced the deposit and resulted in ore dilution. These faults strike N65W, N20E, N45E, and N35E. Post-ore dikes strike N60-75E and have steep northwest to vertical dips. Post-ore faulting and dike activity is probably related to development of the Republic graben.

**COMMENTS:** The Mount Tolman deposit is owned (1990) by the Colville Confederated Tribes. Prior to tribal acquisition of the land, numerous mines were developed in search of silver ore. Mine adits were driven by: Consolidated Mines and Smelting Co., Ltd. (adits include the Iconoclast, Golden Cord, Advance, Silver Ridge, California, Umatilla, Walla Walla, Jumper, Handspike, and Dewey); the Meadow Creek mine (adits include the Blue Bird, Abe Lincoln, King Richard, Blevins, and Sanpoil); the Rover-Bonanza group (adits include the Golden Rule, Golden Chariot, and North Star); and also the adits Bobbie Jay, Josie, Ophir, Byrne, Addie B, Humboldt, Pole Pick, and adits by Illinois Copper and Silver Mining and Milling Co. Bear Creek Mining Co. leased the property from the Colville Business Council in July of 1964. From 1964 to 1976 Bear Creek completed 45,000 ft of drilling and drove a 650-ft adit at elev. 3,000 ft. AMAX Exploration, Inc. signed an agreement in 1978 and undertook a major 2-year development program. They completed 386,827 ft of drilling, three adits totaling 2,160 ft, and metallurgical testing. Drilling costs alone from August 18, 1978, to December 1980 were \$10,167,739. AMAX terminated its lease in 1982.

**UNPUBLISHED INFORMATION:** Utterback, W.C., 1983, Geology of the Mount Tolman molybdenum deposit. This private report is in DGER files.

#### REFERENCES

- Colville Confederated Tribes Geology Department, 1984, Revised geology and mineral potential of the Colville Indian Reservation, Washington, 1984: Colville Confederated Tribes [Nespelem, Wash.], 2 v.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Mountain Lion** (329)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' 47.16" N	118° 46' 2.49" W	W½NW¼NW¼ sec. 27, 37N, 32E	
LOCATION: on the east side of the North Fork Granite Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	pyrite, quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced ore worth \$200,000 prior to 1910; produced in 1936-1938. Production after 1938 is included with production for the Knob Hill mine (Hunting, 1956, p. 122).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Mineralization is in three parallel veins of banded quartz in andesite flow breccia. The productive vein is 10-12 ft wide (Hunting, 1956, p. 122). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**North San Poil Fraction (330)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 40' .10" N	118° 45' 31.38" W	near west line NW¼NE¼ sec. 34, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1911 (Hunting, 1956, p. 123).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: This deposit is similar to other deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Old Hickory (331)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 38' 20.83" N	118° 44' 33.79" W	N1/2 NE1/4 sec. 12, 36N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1918 and 1931 (Hunting, 1956, p. 123).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: This deposit is similar to other deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Overlook (355)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Belcher	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooke Mountain	1:24,000	Republic	Okañogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 41' 56.32" N	118° 34' 8.00" W	NW1/4 sec. 18, 37N, 34E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	graywacke, argillite, chert, limestone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	pyrite	chlorite, sericite	
Ag	pyrrhotite		
Cu	magnetite		
Fe	chalcopyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
manto-like massive replacement stockwork and disseminated			

**PRODUCTION:** The first gold-silver dore was poured (1990) from the jointly operated Overlook and Kettle mines. Anticipated production for the first year is 110,000 oz; the majority will be from the Overlook deposit (Joseph, 1990, p. 20).

**TECTONIC SETTING:** Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the Republic graben (Muessig, 1967, geol. map).

**ORE CONTROLS:** Gold and silver mineralization at the Overlook deposit occurs in manto-like replacement bodies of magnetite, pyrite, and pyrrhotite in limestone and as disseminated and stockwork bodies overlying the massive, iron-rich horizon in the limestone. Currently (1990), this deposit is being mined for its gold-silver mineralization (Tschauder, 1989, p. 245).

**GEOLOGIC SETTING:** The Overlook deposit is in Permian-Triassic limestone, dolomitic limestone, argillite, and graywacke. The ore bodies lie next to a network of dikes, sills, and intrusive bodies of the Scatter Creek Rhyodacite (Muessig, 1967, p. 113).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., 1990, Mineral industry news notes: Washington Geologic Newsletter, v. 18, no. 2, p. 19-20.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Oversight (354)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Belcher	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooke Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 42' 27.80" N	118° 33' 13.87" W	near center E1/2 sec. 18, 37N, 34E	
LOCATION: on the southeast side of Cooke Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary and metavolcanic rocks	graywacke, argillite, chert, limestone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Cu Au	pyrite pyrrhotite magnetite chalcopyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
manto-like massive replacement			

PRODUCTION: Produced in 1934 (Hunting, 1956, p. 195).

TECTONIC SETTING: Eocene felsic rocks intruded Permian-Triassic metasedimentary and metavolcanic rocks of the Republic graben (Muessig, 1967, geol. map).

ORE CONTROLS: Ore minerals occur in a lenticular body replacing limestone. Some ore is nearly pure magnetite (Hunting, 1956, p. 195).

GEOLOGIC SETTING: A massive replacement lens in Permian-Triassic limestone (Muessig, 1967, geol. map).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.



**Pearl (332)**

ALTERNATE NAMES		DISTRICT Republic	COUNTY Ferry
PRIMARY QUADRANGLE Storm King Mountain	SCALE 1:24,000	1/2° x 1° QUAD Republic	1° x 2° QUAD Okanogan
LATITUDE 48° 39' 53.89" N	LONGITUDE 118° 45' 9.97" W	SECTION, TOWNSHIP, AND RANGE near center NE¼ sec. 34, 37N, 32E	
LOCATION: adjoining the Surprise claim on the north			
HOST ROCK: NAME Sanpoil Volcanics	LITHOLOGY dacite and andesite flows and flow breccias	AGE Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION Scatter Creek Rhyodacite		AGE Eocene	
COMMODITIES Au Ag	ORE MINERALS gold	NON-ORE MINERALS quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE vein hot springs		MINERALIZATION AGE Eocene	

PRODUCTION: Produced from 1909 to 1922 (Hunting, 1956, p. 123).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: A 12-ft-wide vein (Surprise) continues through the entire length of the claim (1,500 ft). Wall rock is propylitic quartz latite porphyry (Hunting, 1956, p. 123). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Plum Bar placer (347)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Mica Mountain	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 55' 46.71" N	118° 48' 57.81" W	lots 7 and 8, sec. 17, 28N, 32E	
LOCATION: about 6 mi up the Columbia River from Grand Coulee Dam, on the north side of the river			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: J. H. Collins washed 700-800 yd<sup>3</sup>/day in 1934. Also produced in 1938-1939 (Hunting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Princess Maude** (333)

ALTERNATE NAMES		DISTRICT	COUNTY
Southern Republic		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 38' 4.50" N	118° 44' 51.08" W	near center sec. 12, 36N, 32E, near the base of the east slope of Copper Mountain	
LOCATION: near center sec. 12, 36N, 32E, near the base of the east slope of Copper Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, calcite, laumontite, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** The mine has produced (Hunting, 1956, p. 123).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** Mineralization is in a 2-4-ft wide vein in propylitic andesite. The vein consists of vitreous white quartz that shows lines of crustification parallel to the walls (Hunting, 1956, p. 123). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Quilp** (334)

ALTERNATE NAMES		DISTRICT	COUNTY
Imperator Eureka		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 32.76" N	118° 44' 41.07" W	NW¼SW¼ sec. 35, 37N, 32E	
LOCATION: on the east side of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold chalcopyrite silver	quartz, pyrite; hydrothermal alteration is low sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** Total production of \$720,938.70 worth of ore to the end of 1920; produced in 1936, 1937 (22,402 tons), 1938 (9,828 tons), and in 1939-1940 (Hunting, 1956, p. 123).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** The vein of chalcedonic banded quartz is 7-8 ft wide and cuts propylitic andesite (Hunting, 1956, p. 123). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Republic** (335)

ALTERNATE NAMES		DISTRICT	COUNTY
Blaine Republic		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 38' 14.52" N	118° 44' 43.40" W	W1/2NE1/4 sec. 12, 36N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Se	gold	quartz, calcite, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

**PRODUCTION:** Estimated production of \$1,400,000 prior to 1910; shipped 2,757 tons of ore to a smelter in 1937; shipped in 1933-1946 (Hunting, 1956, p. 124).

**TECTONIC SETTING:** Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

**ORE CONTROLS:** The vein is as much as 10 ft wide, but averages about 3 ft. The vein is composed principally of chalcedonic quartz and is concentrically crustified. Crustifications are marked by dark, crenulated bands (Hunting, 1956, p. 124). (See Republic district for additional details.)

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

**REFERENCES**

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Jonte, J. H., 1942, The relationship of selenium and gold in ores from the Republic district: State College of Washington Master of Science thesis, 33 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Rogers Bar placer (348)**

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Hunters	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 5' 19.79" N	118° 14' 11.92" W	sec. 23, 30N, 36E	
LOCATION: on the west bank of the Columbia River, 2 mi downstream of the town of Hunters			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pt	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Four operators were working and apparently producing in 1934 (Hunting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. Deposits consist of three bars, 30, 75, and 100 ft above the river. Property includes 1,500 acres of land. The best pay gravel is found in bars exposed only at low water (Hunting, 1956, p. 184).

## REFERENCES

- Collier, A. J., 1907, Gold-bearing river sands of northeastern Washington. *In* Contributions to economic geology 1906; Part I—Metals and nonmetals, except fuels: U.S. Geological Survey Bulletin 315, p. 56-70.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**San Poil (336)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 53.32" N	118° 45' 20.63" W	W1/2NE1/4 sec. 34, 37N, 32E	
LOCATION: on the west side of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, calcite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1909-1921, 1931, and 1935; 200 tons of ore were produced prior to 1902 (Hunting, 1956, p. 124).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Latite porphyry is cut by a quartz-calcite vein thought to be the southward extension of the Ben Hur vein. The vein is cut by several faults. Ore occurs in lenses as much as 8 ft thick (Hunting, 1956, p. 124). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Ferry****Seattle (337)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 53.64" N	118° 45' 58.26" W	W1/2 NW1/4 sec. 34, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1938 and in 1939 (Hunting, 1956, p. 124).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: The deposit is similar to other deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.



**Shamrock** (358)

ALTERNATE NAMES		DISTRICT	COUNTY
Iron Creek			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Louie Creek	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 8' 46.21" N	118° 37' 34.18" W	sec. 2, 30N, 33E, and secs. 26, 27, 34, and 35, 31N, 33E	
LOCATION: 12.5 mi northeast of Keller			
HOST ROCK: NAME	LITHOLOGY	AGE	
Covada Group	limestone, dolomite, impure marble	Ordovician?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granodiorite		Paleocene - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Au Pb Zn Cu Ni	galena malachite cerussite lead oxide	limestone	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

**PRODUCTION:** Produced in 1914 (50 tons), in 1922 (3,500 oz Ag), and in 1926 (Hunting, 1956, p. 277). Total silver production around 7,000 oz (Moen, 1976, p. 102).

**TECTONIC SETTING:** Paleocene to Eocene granitic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 12).

**ORE CONTROLS:** Mineralization is in silicified limestone block surrounded by granodiorite. Silver-lead ore (vein) is 8 in.-8 ft wide and 700 ft long. Nickel occurs in a mineralized zone 100 ft wide in which there is a high-grade ore zone 55 ft wide (Hunting, 1956, p. 277).

**GEOLOGIC SETTING:** Granodiorite (the Paleocene to Eocene granodiorite of Manila Creek) intrudes carbonate rocks of the Ordovician? Covada Group at the Shamrock mine (Hunting, 1956, p. 277; Joseph, 1990, geol. map, p. 21).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Silver Bell (357)**

ALTERNATE NAMES		DISTRICT Covada	COUNTY Ferry
PRIMARY QUADRANGLE Cedonia	SCALE 1:24,000	1/2° x 1° QUAD Nespelem	1° x 2° QUAD Okanogan
LATITUDE 48° 14' 14.77" N	LONGITUDE 118° 10' 45.47" W	SECTION, TOWNSHIP, AND RANGE near NE corner sec. 31, 32N, 37E	
LOCATION: near the end of a southeast spur of Rattlesnake Mountain			
HOST ROCK: NAME Covada Group	LITHOLOGY argillite and wacke	AGE Ordovician??	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION granite and granodiorite near Meteor		AGE Jurassic - Cretaceous	
COMMODITIES Pb Zn Au Ag Cu	ORE MINERALS galena sphalerite	NON-ORE MINERALS pyrite, hornstone	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: Produced 29 tons in 1940 and 84 tons in 1941 (Hunting, 1956, p. 214).

TECTONIC SETTING: Jurassic to Cretaceous granitic to granodioritic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 34).

ORE CONTROLS: The vein is in hornstone that weathers rusty and is irregularly mineralized. An exposed vein width of at least 1 ft contained a "fair proportion" of galena (Hunting, 1956, p. 214).

GEOLOGIC SETTING: The Silver Bell mine is in hornstone which probably is contact metamorphosed Covada Group (Ordovician?) argillite adjacent to Jurassic to Cretaceous granite to granodiorite of Meteor (Joseph, 1990, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., 1990, Mineral industry news notes: Washington Geologic Newsletter, v. 18, no. 2, p. 19-20.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Silver Leaf** (306)

ALTERNATE NAMES		DISTRICT Covada Meteor	COUNTY Ferry
PRIMARY QUADRANGLE Cedonia	SCALE 1:24,000	1/2° x 1° QUAD Nespelem	1° x 2° QUAD Okanogan
LATITUDE 48° 14' 17.87" N	LONGITUDE 118° 11' 17.29" W	SECTION, TOWNSHIP, AND RANGE center of north line, sec. 31, 32N, 37E	
LOCATION: on the south slope of Rattlesnake Mountain			
HOST ROCK: NAME Covada Group	LITHOLOGY wacke and quartzite	AGE Ordovician?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION granite and granodiorite near Meteor		AGE Jurassic - Cretaceous	
COMMODITIES Ag Pb Zn W Sb	ORE MINERALS tetrahedrite chalcopyrite sphalerite galena stibnite scheelite silver pyrargyrite wolframite	NON-ORE MINERALS quartz, pyrite	

DEPOSIT TYPE vein	MINERALIZATION AGE
----------------------	--------------------

PRODUCTION: Produced in 1911, 1912, 1915, 1916, and 1935. Produced a small amount of tungsten in 1915 (Hunting, 1956, p. 294). According to Moen (1976, p. 98), the mine probably produced about 2,000 oz of Ag from a small total tonnage.

TECTONIC SETTING: Jurassic to Cretaceous granitic to granodioritic rocks intrude metamorphosed Ordovician? rocks (Joseph, 1990, geol. map, p. 34).

ORE CONTROLS: Mineralization is in quartz lenses in a 50-ft-wide shear zone in limestone, schist, and argillite (Moen, 1976, p. 98).

GEOLOGIC SETTING: The Silver Leaf deposit is in limestone, schist, and argillite of the Ordovician? Covada Group (Joseph, 1990, geol. map; Moen, 1976, p. 98).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

***Singer placer*** (349)

ALTERNATE NAMES		DISTRICT	COUNTY
		Danville	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Curlew	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 59' 45.96" N	118° 31' 50.75" W	secs. 4 and 5, 40N, 34E	
LOCATION: on Fourth of July Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE	MINERALIZATION AGE		
placer	Quaternary		

PRODUCTION: Produced in 1934 (Huntting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**South Penn (338)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 41' 24.62" N	118° 45' 20.69" W	sec. 22, 37N, 32E	
LOCATION: A default location is selected at the center of section 22.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced between 1941 and 1949; 172 tons in 1945 yielded \$2,168 (Hunting, 1956, p. 125).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: This deposit is similar to other deposits of the Republic district. (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Stray Dog** (366)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Kewa	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 14' 43.27" N	118° 15' 24.42" W	SW1/4 NW1/4 sec. 27, 32N, 36E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
granite to granodiorite near Meteor		granite and granodiorite	Jurassic - Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	chalcopyrite	pyrite, quartz	
Pb	sphalerite		
Cu	galena		
Zn	tetrahedrite		
Au	pyrargyrite		
	argentite		
	arsenopyrite		
	native silver		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Two shipments of 20 tons each reported prior to 1910 (Hunting, 1956, p. 294).

TECTONIC SETTING: Jurassic to Cretaceous intrusive rocks intrude Ordovician? sedimentary rocks of the area (Joseph, 1990, geol. map).

ORE CONTROLS: The ore is in a sparsely mineralized quartz vein that ranges from a stringer to 2 ft wide and occupies a shear zone in quartz monzonite porphyry (Hunting, 1956, p. 294).

GEOLOGIC SETTING: The Stray Dog deposit is in Jurassic to Cretaceous granitic to granodioritic rocks (Joseph, 1990, geol. map, p. 34).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Weaver, C. E., 1913, Geology and ore deposits of the Covada mining district: Washington Geological Survey Bulletin 16, 87 p.

**Surprise** (340)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 39' 39.00" N	118° 44' 57.13" W	SE¼NE¼ and NE¼SE¼ sec. 34, 37N, 32E	
LOCATION: on the east side of Eureka Gulch			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, pyrite; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Production exceeded \$1,000,000; produced in 1910-1923, 1934, 1938, and 1947-1950 (Hunting, 1956, p. 125).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Mineralization is present in a 4-8-ft-wide vein in propylitic quartz latite. The vein consists of banded quartz and includes fragments of country rock (Hunting, 1956, p. 125). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Talisman (313)**

ALTERNATE NAMES		DISTRICT	COUNTY
Laurier			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Independent Mtn	1:24,000	Republic	Okañogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 59' .70" N	118° 16' 14.13" W	sec. 4, 40N, 36E	
LOCATION: on the east slopes of Owl Mountain, 1 mi south of the international border			
HOST ROCK: NAME		LITHOLOGY	AGE
unnamed paragneiss unnamed amphibolite		gneiss amphibolite	pre-Tertiary pre-Tertiary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au Ag Pb W Cd Bi	chalcopyrite sphalerite galena scheelite	pyrite, magnetite, epidote, garnet, zoisite	

DEPOSIT TYPE	MINERALIZATION AGE
--------------	--------------------

contact metamorphic	
---------------------	--

PRODUCTION: Produced in 1915 (\$15,000), 1916 (\$36,000), 1946, 1947, 1948, (54 tons of concentrates), 1949 (37 tons of concentrates), and 1950 (18 tons of Zn concentrates) (Hunting, 1956, p. 55).

TECTONIC SETTING: Part of the suite of metamorphic rocks of the Kettle metamorphic core complex (Stoffel, 1990, geol. map).

ORE CONTROLS: Shallowly dipping contact-metamorphic deposit in schist. The ore body is 1-12 ft thick (Hunting, 1956, p. 55).

GEOLOGIC SETTING: The deposit is in paragneiss near the contact with amphibolite. The metamorphic rocks are part of the Kettle metamorphic core complex (Stoffel, 1990, geol. map, p. 45-47).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.



**Threemile placer** (350)

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Fort Spokane	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 56' 35.12" N	118° 21' 32.92" W	sec. 11, 28N, 35E, on the Columbia river	
LOCATION: on the Columbia river, 3 mi above the mouth of the Spokane River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Worked by the Chinese prior to 1910 (Hunting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. Gold occurs in large, thin flakes worth about 0.01 cents each and as flour gold. The bench is 20 ft above the river and contains large angular boulders. Pay gravels occupied the spaces between boulders (Hunting, 1956, p. 184).

## REFERENCES

- Collier, A. J., 1907, Gold-bearing river sands of northeastern Washington. *In* Contributions to economic geology 1906; Part I—Metals and nonmetals, except fuels: U.S. Geological Survey Bulletin 315, p. 56-70.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Tom Thumb** (341)

ALTERNATE NAMES		DISTRICT	COUNTY
		Republic	Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Storm King Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 41' 45.87" N	118° 45' 25.51" W	S1/2SE1/4 sec. 15, and N1/2N1/2 sec. 22, 37N, 32E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Sanpoil Volcanics	dacite and andesite flows and flow breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Scatter Creek Rhyodacite		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE		MINERALIZATION AGE	
vein hot springs		Eocene	

PRODUCTION: Produced in 1908-1910, 1915, 1916, 1934, and 1938 (Hunting, 1956, p. 125).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Quartz is spread widely through bedded tuff. Lodes are not well defined. One 8-ft-wide vein is present in andesite (Hunting, 1956, p. 125). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.
- Umpleby, J. B., 1910, Geology and ore deposits of Republic mining district: Washington Geological Survey Bulletin 1, 65 p.

**Trade Dollar (342)**

ALTERNATE NAMES		DISTRICT Republic	COUNTY Ferry
PRIMARY QUADRANGLE Storm King Mountain	SCALE 1:24,000	1/2° x 1° QUAD Republic	1° x 2° QUAD Okanogan
LATITUDE 48° 40' 20.52" N	LONGITUDE 118° 45' 39.07" W	SECTION, TOWNSHIP, AND RANGE E1/2 SW1/4 sec. 27, 37N, 32E	
LOCATION:			
HOST ROCK: NAME Sanpoil Volcanics	LITHOLOGY dacite and andesite flows and flow breccias	AGE Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION Scatter Creek Rhyodacite		AGE Eocene	
COMMODITIES Au Ag	ORE MINERALS gold	NON-ORE MINERALS quartz; hydrothermal alteration is low-sulfur adularia-sericite type	
DEPOSIT TYPE vein hot springs		MINERALIZATION AGE Eocene	

PRODUCTION: Produced about \$25,000 worth of ore by 1934 (Huntting, 1956, p. 125).

TECTONIC SETTING: Epithermal deposits of the Republic district are found in the Eocene Sanpoil Volcanics. (See Republic district for additional details.)

ORE CONTROLS: Quartz vein varies from 20 in. to 13 ft wide (Huntting, 1956, p. 125). (See Republic district for additional details.)

GEOLOGIC SETTING: Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics (Muessig, 1967, p. 50-51; Tschauder, 1989, p. 242).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Valley** (343)

ALTERNATE NAMES		DISTRICT	COUNTY
Golden Valley Lame Foot			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Republic	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 44' 3.76" N	118° 41' 10.06" W	E½ sec. 6, 37N, 33E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Scatter Creek Rhyodacite	porphyritic dacite	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Se	auriferous selenide tetrahedrite	quartz, calcite, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene	

**PRODUCTION:** Produced 1,994 tons prior to 1941, 5,800 tons in 1942, 1,302 tons in 1943, and in 1950 (Hunting, 1956, p. 125).

**TECTONIC SETTING:** Eocene volcanic rocks host mineral deposits in the Republic graben.

**ORE CONTROLS:** The vein is in andesite and is estimated to average 7 ft wide and 1,200 ft long (Hunting, 1956, p. 125).

**GEOLOGIC SETTING:** Host rocks for gold deposits of the Republic district are dacite and andesite flows, flow breccias, tuffs, and tuff breccias of the Eocene Sanpoil Volcanics and subvolcanic intrusions of the Scatter Creek Rhyodacite (Muessig, 1967, p. 54-58).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Jonte, J. H., 1942, The relationship of selenium and gold in ores from the Republic district: State College of Washington Master of Science thesis, 33 p.
- Muessig, Siegfried, 1967, Geology of the Republic quadrangle and a part of the Aeneas quadrangle, Ferry County, Washington: U.S. Geological Survey Bulletin 1216, 135 p., 1 pl.
- Tschauder, R. J., 1989, Gold deposits in northern Ferry County, Washington. In Joseph, N. L. and others; editors, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 241-253.

**Whitestone placer (351)**

ALTERNATE NAMES		DISTRICT	COUNTY Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Whitestone Rock	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 55' 23.25" N	118° 32' 17.77" W	sec. 16, 28N, 34E	
LOCATION: along the Columbia River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced in 1933 (Hunting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Wilmont Bar placer (352)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Ferry
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Miller Mtn	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 2' 55.39" N	118° 16' 17.97" W	lot 6, sec. 4, 29N, 36E	
LOCATION: on the north bank of the Columbia River, opposite the Gerome post office			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ce Th	gold monazite magnetite ilmenite zircon	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced in 1934 (Hunting, 1956, p. 184).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. The deposit consists of two terraces, one 20 ft and the other 100 ft above the Columbia River. In the lower terrace, gold occurs in the uppermost 1-5 ft of material (Hunting, 1956, p. 184).

## REFERENCES

- Collier, A. J., 1907, Gold-bearing river sands of northeastern Washington. *In* Contributions to economic geology 1906; Part I—Metals and nonmetals, except fuels: U.S. Geological Survey Bulletin 315, p. 56-70.
- Day, D. T.; Richards, R. H., 1906, Useful minerals in the black sands of the Pacific slope. *In* U.S. Geological Survey, Mineral resources of the United States, calendar year 1905: U.S. Geological Survey, p. 1175-1258.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1929, Platinum and black sand in Washington. *In* Contributions to economic geology (short papers and preliminary reports) 1928; Part I—Metals and nonmetals except fuels: U.S. Geological Survey Bulletin 805, p. 1-15.

**Zalla M (367)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Sheridan	Ferry
PRIMARY QUADRANGLE	SCALE	1½° x 1° QUAD	1° x 2° QUAD
Bodie Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 45' 40.64" N	118° 50' .37" W	W½ sec. 30, 38N, 32E	
LOCATION: a few hundred yards east of the Ferry-Okanogan County boundary line			
HOST ROCK: NAME	LITHOLOGY	AGE	
Klondike Mountain Formation	rhyolite flows and pyroclastic breccias	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
phonolite			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	chalcopyrite	pyrite, quartz, calcite, chlorite, epidote, zoisite, sericite	
Cu	galena		
Au	fluorite		
	hessite		
	krennerite		
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Produced \$40,000 worth of ore prior to 1903; ore was shipped from the dump in 1935 (Hunting, 1956, p. 295).

TECTONIC SETTING: Sequence of Eocene volcanic rocks in the Toroda Creek graben (Pearson, 1967, geol. map).

ORE CONTROLS: Mineralization is in a well-defined 1-6-ft-wide quartz vein filling an irregular fracture in phonolite. The vein is sparsely mineralized and low grade except in the zone of secondary enrichment, which extends from the surface to a depth of about 150 ft (Hunting, 1956, p. 295). Wall rock adjacent to the vein is brecciated and sheared, and the breccia is cemented by quartz, chlorite, and fluorite. Ore was mined from ore shoots in the zone of secondary enrichment (Moen, 1980, p. 70).

GEOLOGIC SETTING: The Zalla M is in flows and breccias of the Eocene Klondike Mountain Formation. The rocks appear to be rhyolitic in composition, but on close examination, outlines of what appear to be pyroxene phenocrysts are found. This suggests the rocks are probably altered andesite (Stoffel, 1990, geol. map, p. 11; Pearson, 1967, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Pearson, R. C., 1967, Geologic map of the Bodie Mountain quadrangle, Ferry and Okanogan Counties, Washington: U.S. Geological Survey Geologic Quadrangle Map GQ 636, 1 sheet, scale 1:62,500, with 4 p. text.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.

***Chinaman Bar placer*** (302)

ALTERNATE NAMES		DISTRICT	COUNTY
			Grant
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Priest Rapids NE	1:24,000		Walla Walla
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 38' 14.17" N	119° 46' 6.26" W	secs. 10 and 11, 13N, 24E	
LOCATION: at Chinaman Bar on the Columbia River, 4 mi east of Priest Rapids			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE	MINERALIZATION AGE		
placer	Quaternary		

PRODUCTION: Produced from 1939 to 1941 (Hunting, 1956, p. 185).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

- Carithers, Ward, 1943, Directory of Washington mining operations: Washington Division of Mines and Mining Information Circular 8, 36 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.



**Burnt Peak (624)**

ALTERNATE NAMES		DISTRICT	COUNTY
Burnt Mountain Burnt Hill			Grays Harbor
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Quinault Lake	1:62,500	Shelton	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 19' 36" N	123° 52' 22" W	SE1/4 sec. 7, 21N, 9W	
LOCATION: elev. 800 ft, approximately 0.25 mi west of the Burnt Hill lookout station. The property is reached from Amanda Park on Lake Quinault by following US Highway 101 south 10 mi to the Burnt Hill road, thence east 1.2 mi to its junction with the Newbury Creek road, thence by the right branch 0.7 mi uphill to the mine road, then 0.5 mi along the mine road.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	basalt	Eocene	
Crescent Formation	red limy argillite	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite hausmannite	jasper	
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

**PRODUCTION:** Produced approximately 80 tons of ore (Hunting, 1956, p. 258).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** Manganese occurs in discontinuous pods. The mineralization is in a highly siliceous ferruginous rock, which on a fractured surface shows small veinlets of clear to milky, crystalline quartz (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

**GEOLOGIC SETTING:** Altered basalt is the principal country rock for the Burnt Hill deposit. Superficial weathering gives the basalt a yellowish to light-brown color; however, megascopic examination of a newly fractured surface reveals fine-grained, greenish-gray material (data from USGS MRDS, 1990). The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

**REFERENCES**

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

# Grays Harbor

## Egge (620)

ALTERNATE NAMES		DISTRICT	COUNTY
Quinault			Grays Harbor
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Quinault Lake	1:62,500	Shelton	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 22' 31" N	123° 51' 40" W	secs. 19, 20, and 30, 22N, 9W	
LOCATION: about 6.3 mi south of Quinault. Claims cover approximately 3 mi <sup>2</sup> on Quinault Ridge at the headwaters of Cook, Skunk, and Stevens Creeks			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	basalt	Eocene	
Crescent Formation	red limy argillite	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

PRODUCTION: Minor production in 1941 and 1952 (Hunting, 1956, p. 258).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

COMMENTS: Nearby claims include: Black Wonder (produced 5 tons in 1936), Antlers, Pioneer, Lizard (small amount produced in 1916), and Stevens Creek (produced 5 tons in 1941, 1952) (Hunting, 1956, p. 257);

### REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Elma (623)**

ALTERNATE NAMES		DISTRICT	COUNTY
Dennis			Grays Harbor
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
McCleary	1:24,000	Shelton	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 00' 49" N	123° 18' 45" W	S 1/2 SE 1/4 sec. 28, 18N, 5W	
LOCATION: elev. 200 ft, 3.5 mi east of the railroad at Elma. Southeast part of the county; 2.75 mi off the Olympic highway			
HOST ROCK: NAME		LITHOLOGY	AGE
Lincoln Creek Formation		sandstone	Eocene - Oligocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Ti	ilmenite magnetite titaniferous magnetite		
DEPOSIT TYPE		MINERALIZATION AGE	
fossil placer		Eocene - Oligocene	

PRODUCTION: Produced 1 carload and a 1,000-lb shipment for testing purposes (Hunting, 1956, p. 196).

TECTONIC SETTING: Forearc volcanoclastic basin that grades eastward into nonmarine volcanoclastic rocks associated with the Cascade magmatic arc (Logan, 1987, geol. map, p. 8).

ORE CONTROLS: This fossil heavy-mineral deposit occurs as a 1-4-ft-thick bed of stratified, weakly consolidated black sand that covers 2-3 acres (Hunting, 1956, p. 196). A specimen of the ore, studied under a hand lens, appears to be made up of small, well-rounded grains of magnetite and ilmenite, which are firmly cemented together (data from USGS MRDS, 1990).

GEOLOGIC SETTING: The Lincoln Creek Formation is a marine, poorly bedded to massive, dominantly siltstone unit of predominantly basaltic and andesitic source-rock composition (Logan, 1987, p. 8).

## REFERENCES

- Logan, R. L., compiler, 1987, Geologic map of the south half of the Shelton and south half of the Copalis Beach quadrangles, Washington: Washington Division of Geology and Earth Resources Open File Report 87-9, 15 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

**Esther-Irene** (621)

ALTERNATE NAMES		DISTRICT	COUNTY
			Grays Harbor
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Quinault Lake	1:62,500	Shelton	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 22' 55" N	123° 52' 47" W	SW1/4NW1/4 sec. 19, 22N, 9W	
LOCATION: elev. 1,265 ft, about 3.9 mi north-northwest of Burnt Hill lookout, at the head of Phillips Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	altered greenstone	Eocene	
Crescent Formation	red limy argillite	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite neotocite manganite psilomelane		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

PRODUCTION: Produced at least 75 tons of ore (Hunting, 1956, p. 258).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Minor faults and fractures occur in the greenstone. At the mine two small manganiferous ore bodies are present in siliceous zones in altered basalt. In the upper siliceous zone, the manganiferous area is 6 ft by 30 ft, and the lower area is 6 ft wide and 22 ft long. The siliceous zone of the lower manganese area is about 10 ft by 70 ft. Manganese is present as silicates and, in minor amounts, as secondary oxides (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: The host basalt of the Crescent Formation is somewhat bleached at the Esther-Irene deposit (data from USGS MRDS, 1990). Paleocene and Eocene pillow basalts are compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Green, S. H., 1945, Manganese deposits of the Olympic Peninsula, Washington: Washington Division of Mines and Mining Report of Investigations 7, 45 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Skunk Creek No. 19 (622)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Grays Harbor
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Quinault Lake	1:62,500	Shelton	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 22' 21" N	123° 52' 37" W	sec. 30, 22N, 9W (probable location)	
LOCATION: about 3.3 mi north-northwest of Burnt Hill lookout. Access to the Skunk Creek No. 19 mine is by way of the Quinault Ridge Road northeasterly about 1.75 mi to an unmarked road junction; from the junction the easterly branch extends 1 mi to the workings.			
HOST ROCK: NAME		LITHOLOGY	AGE
Crescent Formation		basalt	Eocene
Crescent Formation		red limy argillite	Eocene
ASSOCIATED IGNEOUS ROCK: DESCRIPTION			AGE
Crescent Formation volcanic rocks			Eocene Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite neotocite pyrolusite rhodonite goethite hematite	manganocalcite, quartz, limonite, goethite, hematite, spessartite garnet, chlorite, and pyrite.	
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

**PRODUCTION:** Produced 150 tons that averaged 19.5% Mn; shipped to Provo, Utah, steel mill in 1952 (Magill, 1952, p. 71). In 1953, a 45-ton test shipment was sent to the U.S. Bureau of Mines in Albany, Oregon (data from USGS MRDS, 1990).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** The deposit is lens shaped. Limestone, so prevalent in deposits of the area, is present only in small quantities at the Skunk Creek No. 1 deposit (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

**GEOLOGIC SETTING:** The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

**REFERENCES**

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Elkhorn** (616)

ALTERNATE NAMES		DISTRICT	COUNTY
Karnes			Jefferson
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mt Jupiter	1:24,000	Mt Olympus	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 44' 18" N	123° 00' 00" W	secs. 13 and 24, 26N, 3W	
LOCATION: about 3.5 mi northeast of the Mt. Jupiter lookout. The Elkhorn group of claims is on the south slope of Mount Constance. The claims are bounded by Bull Elk Creek on the west, Miner's Creek on the east, and the Dosewallips River on the south. The property is reached from Brinnon on US Highway 101 by following the road along the Dosewallips River westerly for about 13 mi. The main workings range in elev. from 1,200 to 4,200 ft.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	red limestone	Eocene	
Crescent Formation	basalt	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite	hematite	
Fe	neotocite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

PRODUCTION: Produced in the 1940s (data from USGS MRDS, 1990).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Basalt in the mine area is locally highly fractured and cut by many minor faults. Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: The area is underlain by basalt lava flows and intercalated tuff and red argillaceous limestone; much of the basalt is altered. Pillow basalt is exposed in places. The basaltic flows generally trend north and stand nearly vertical; locally, they are greatly deformed and fractured. Three red limestone beds and intercalated basalts form three or more distinct parallel ridges extending from the Dosewallips River northward to the top of the divide. Three limestone interbeds in basalt contain lenses of manganese ore, some as much as 25 ft thick. The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**Anderson (600)**

ALTERNATE NAMES		DISTRICT	COUNTY
Baring		Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Index	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 45' 17.70" N	121° 30' 19.24" W	S1/2 sec. 10 and near SE corner sec. 11, 26N, 10E	
LOCATION: about 0.75 to 1 mi from Baring			
HOST ROCK: NAME	LITHOLOGY	AGE	
Index batholith unnamed sedimentary rocks	granodiorite and tonalite limy quartzite	Oligocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Pb Zn Ag Au Cu	magnetite galena sphalerite	amphibole	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic		Oligocene	

**PRODUCTION:** One carload shipped (Hunting, 1956, p. 196).

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** Ore body is in limy quartzite and is presumably a contact metamorphic deposit. The exposed portion is a lenticular mass that is 25-30 ft high and 15 ft wide and that exposed approximately 2,000 tons (Hunting, 1956, p. 196). Shedd (1924, p. 66) reports the presence of titanium in assays from the Anderson property.

**GEOLOGIC SETTING:** Tabor and others (1982) map the area as all within granodiorite and tonalite of the Index batholith. The mineralization presumably occurs in a small roof pendant or included block.

**REFERENCES**

- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.
- Shedd, Solon, 1924, The mineral resources of Washington with statistics for 1922; with an article on coal and coke, by G. W. Evans: Washington Division of Geology Bulletin 30, 224 p.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

**Apex (595)**

ALTERNATE NAMES		DISTRICT	COUNTY
Bondholders Syndicate Apex Gold National Gold		Miller River Money Creek	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Si	1:62,500	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 41' 42.00" N	121° 30' 44.13" W	SW1/4 sec. 34, 26N, 10E	
LOCATION: near the headwaters of Money Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Snoqualmie batholith	tonalite and granodiorite	Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Pb	chalcopyrite galena sphalerite tetrahedrite stibnite(?)	pyrite, arsenopyrite, arsenolite, quartz, tourmaline, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced 300 tons worth \$80,000 prior to 1901. Also produced ore worth a total of \$300,000 from mining in 1905, 1908, 1910, 1912, 1913, 1916-1920, 1926, 1928, and 1936-1943 (Huntting, 1956, p. 126). Production from 1905 to 1940 was valued at \$220,000 (Moen, 1976, p. 166). Shipments in 1921 averaged between 21% and 26% As, 18 and 20 oz/ton Ag, 1.5 and 2.5 oz/ton Au, and 4.5 and 6% Pb (Livingston, 1971, p. 163).

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** Mineralization is in a quartz vein 2-6 ft wide which follows a continuous fissure in granodiorite. High-grade ore occurs in shoots in the vein (Huntting, 1956, p. 126). The paragenetic sequence is quartz and tourmaline followed by pyrite and quartz, then quartz and stibnite(?), sphalerite, quartz, tetrahedrite and galena, chalcopyrite, and calcite (Coats, 1932, p. 19). Arsenolite (a secondary arsenic mineral) is found in the mine workings (Coats, 1932, p. 17).

**GEOLOGIC SETTING:** The Apex mine occurs in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and locally clinopyroxene. Age of the northern phase is about 25 m.y. based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

**REFERENCES**

- Coats, R. R., 1932, The ore deposits of the Apex gold mine, Money Creek, King County, Washington: University of Washington Master of Science thesis, 48 p., 2 pl.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.



**Carmack (597)**

ALTERNATE NAMES		DISTRICT Snoqualmie	COUNTY King
PRIMARY QUADRANGLE Snoqualmie Pass	SCALE 1:62,500	1/2° x 1° QUAD Snoqualmie Pass	1° x 2° QUAD Wenatchee
LATITUDE 47 24 2.52 N	LONGITUDE 121 27 11.71 W	SECTION, TOWNSHIP, AND RANGE N½ sec. 7 and S ½ sec. 18, 22N, 11E	
LOCATION: on the South Fork Snoqualmie River near Snoqualmie Pass			
HOST ROCK: NAME Snoqualmie batholith	LITHOLOGY tonalite and granodiorite	AGE Miocene	
COMMODITIES Au Ag Pb Cu	ORE MINERALS chalcopyrite galena	NON-ORE MINERALS	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: Shipped 20 tons of ore prior to 1901 (Hunting, 1956, p. 127).

TECTONIC SETTING: The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

ORE CONTROLS: Gold, silver, lead, and copper ore is found in three veins which are 12 ft, 2.5 ft and 1 ft wide (Hunting, 1956, p. 127).

GEOLOGIC SETTING: The Carmack mine is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular, and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and, locally, clinopyroxene. The age of the northern phase is about 25 m.y. based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Frizzell and others, 1984, p. 17).

## REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.

***Cleopatra group*** (602)

ALTERNATE NAMES		DISTRICT	COUNTY
Cleopatra Aces Up		Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 38' 7.97" N	121° 27' 31.79" W	SE1/4 sec. 24, 25N, 10E and SW1/4 sec. 30, 25N, 11E	
LOCATION: The Cleopatra is in sec. 24, and the Aces Up is in sec. 30.			
HOST ROCK: NAME		LITHOLOGY	AGE
Snoqualmie batholith		tonalite and granodiorite	Miocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	argentiferous galena	pyrite, arsenopyrite, quartz, kaolinite, sericite	
Pb	chalcopyrite		
Cu	tetrahedrite		
Zn	sphalerite		
Sb	jamesonite		
Au	dyscrasite(?)		
DEPOSIT TYPE		MINERALIZATION AGE	
mineralized shear zone vein			

**PRODUCTION:** Intermittent prior to 1914. Produced in 1938, 1940, and 1941 (Hunting, 1956, p. 296). The mine was shut down in 1941 under the war powers act when it was classified a nonessential mine (Livingston, 1971, p. 139).

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** Livingston (1971, p. 140) notes two mineralized zones. The country rock along both zones was altered to kaolinite and locally sericite. Argentiferous galena and chalcopyrite are the principal ore minerals; pyrite is abundant. Hunting (1956, p. 296) reports the alteration and mineralization lie along joints in the granodiorite. The two main zones are each about 2.5 ft wide.

**GEOLOGIC SETTING:** The Cleopatra Group of deposits is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and locally clinopyroxene. The northern phase is about 25 m.y. old based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Clipper (605)**

ALTERNATE NAMES		DISTRICT	COUNTY
Snoqualmie Copper Katie Belle		Snoqualmie Burns Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Big Snow Mtn	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 31' 5.46" N	121° 20' 39.67" W	N1/2, sec. 1, 23N, 11E; NW1/4NW 1/4, sec. 3, 23N, 12E; S1/2, sec. 36, 24N, 11E; SW1/4, sec. 34, 24N, 12E.	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Snoqualmie batholith, northern phase		granodiorite and tonalite	Miocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au Mo	chalcopyrite molybdenite galena	quartz, pyrite, siderite, pyrrhotite, arsenopyrite, chlorite; potassic alteration, silicification	
DEPOSIT TYPE		MINERALIZATION AGE	
veins fractures porphyry system			

**PRODUCTION:** Test stope 25 ft by 35 ft long; ore stockpiled on dump.

**TECTONIC SETTING:** Part of a northeast-trending belt of en echelon (NW-striking) faults, fractures, and breccia pipes defining a porphyry copper system.

**ORE CONTROLS:** The Clipper group consists of seven zones, the Clipper, Pedro, Katie Belle, Hawk, Three Brothers, Red Face, and Crawford Creek. The group is in a large area of mineralization in which five narrow structures of higher grade mineralization have been indicated. The mineralization is in shear zones, along fractures, and in veins. One structure, explored by drilling and an adit, was centered on a surface exposure of mineralization that was 40 ft by 800 ft. Samples collected on the surface there had assay values ranging from 0.52% Cu to 1.2% Cu and 0.2% MoS<sub>2</sub>. One drill hole had a 69-ft intercept averaging 0.73% Cu. Copper content of some underground samples ranged from 0.31%-0.98%. Two samples on the J.T. Claim show copper and assay 0.015 oz/ton Au, and 0.040 oz/ton Au (see report in the Clipper file). Inferred reserves of 226,800 tons of 0.9 % Cu are reported in Gualtieri and others (1975) for the Clipper zone. Potential reserves from drilling in the Three Brothers zone are 1.8 million tons averaging 0.7%-0.9% Cu.

**GEOLOGIC SETTING:** The Clipper deposit is in the Snoqualmie batholith, northern phase granodiorite and tonalite with biotite and hornblende; locally it contains clinopyroxene. The rocks are light colored, medium crystalline, mostly equigranular with hypidiomorphic texture, and are coarsely jointed. The northern phase is about 25 m.y. old, on the basis of interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Frizzell and others, 1984, p. 18).

**COMMENTS:** On the Last Chance claim, a 390-ft adit bearing N40W intersects mineralization from 265 to 310 ft; the area was stoped. On the Pedro claim, a 10-ft adit was driven in a 50-ft-wide mineralized zone. On the Katie Belle and Tracy claims, a 15-ft adit and a 100-plus-ft adit were driven. Both the patented and unpatented claims are owned (1990) by United Cascade Mining Company, Inc. Companies who have explored the deposit include Anaconda Copper Co., Howe Sound Co., Climax Molybdenum Corp., Anaconda Co., Bear Creek Mining Co., Cities Services Minerals Corp., Westland Mines Ltd., Natural Resources Development Corp., Houston Oil and Minerals, and Electras Resources.

**UNPUBLISHED INFORMATION:** Melrose, J. W., 1941, Mineralization of the area to be served by the Middle Fork of Snoqualmie River road. Carithers, Ward, 1942, Memorandum to Sheldon L. Glover. These reports are in DGER files.

**REFERENCES**

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Grant, A. R., 1969, Chemical and physical controls for base metal deposition in the Cascade Range of Washington: Washington Division of Mines and Geology Bulletin 58, 107 p.
- Gualtieri, J. L.; Thurber, H. K.; Miller, M. S.; McMahan, A. B.; Federspiel, F. F., 1975, Mineral resources of additions to the Alpine Lakes study area, Chelan, King, and Kittitas Counties, Washington: U.S. Geological Survey Open-File Report 75-3, 161 p., 2 pl.

## **King**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- McIntyre, A. W., 1907, Copper deposits of Washington: American Mining Congress, 9th Annual Session, Report of Proceedings, p. 238-250.
- Purdy, C. P., Jr., 1954, Molybdenum occurrences of Washington: Washington Division of Mines and Geology Report of Investigations 18, 118 p., 6 pl.
- Western Miner, 1967, Westland Mines Ltd.: p. 128.

**Condor-Hemlock (606)**

ALTERNATE NAMES		DISTRICT	COUNTY
Condor Hemlock		Snoqualmie Burns	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Snoqualmie Pass Big Snow Mtn	1:62500 1:24,000	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 29' 48.08" N	121° 21' 36.74" W	sec. 11, 23N, 11E	
LOCATION: sec. 11, 23N, 11E			
HOST ROCK: NAME	LITHOLOGY	AGE	
Snoqualmie batholith, northern phase Snoqualmie batholith	granodiorite and tonalite monzonite dike	Miocene Miocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
hypabyssal andesite dikes and plugs both pre- and post-ore		Tertiary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Mo Au Ag	chalcopryite molybdenite covellite chalcocite	quartz, pyrite, pyrrhotite, chlorite; potassic alteration, silicification, sericite.	
DEPOSIT TYPE		MINERALIZATION AGE	
veins fractures breccia pipe porphyry system			

**PRODUCTION:** No production has been recorded.

**TECTONIC SETTING:** Part of a northeast-trending belt of en echelon (NW-striking) faults, fractures, and breccia pipes defining a porphyry copper system.

**ORE CONTROLS:** The Condor-Hemlock is in a northwest-trending mineralized zone. The Condor is located at the center of the zone, and the Hemlock is located at the southeast end of the zone. The mineralization is in shear zones, along fractures, and in veins in granodiorite and tonalite. Hydrothermal alteration grades from propylitic at the surface to quartz-sericite-chlorite to K-feldspar predominating at depth. Pyrrhotite increases and pyrite decreases with depth. Secondary enrichment of covellite and chalcocite was found in a drill hole when the West Condor fault was intercepted between 151 ft and 203 ft. This 52-ft intercept averaged 1.66 % Cu, more than 1.0 oz/ton Ag, and \$0.75 gold per ton (November 15, 1965 prices). The mineralized zone has a strike length of 1,000 feet, a thickness of 400 feet, and a depth of more than 400 feet. The Condor portion of the zone has potential reserves of 25-30 million tons averaging 0.616 % Cu and 0.032 % MoS<sub>2</sub>. Gold and silver values range from \$0.40 to \$1.25 per ton (October 1967 prices). The Hemlock zone (including part of another zone called the Porter and some adjacent zones) has been tested by 23 drill holes (total footage unknown) and three adits totalling 3,200 ft. By 1975, potential reserves of 91 million tons averaging 0.6-0.8% Cu and 0.02-0.05% MoS<sub>2</sub> in the combined Porter and Hemlock zones was indicated (Gualtieri and others, 1975).

**GEOLOGIC SETTING:** Of the numerous mineralized zones in the Middle Fork Snoqualmie system (Livingston, 1971, p. 152-153), the Condor-Hemlock contains the best demonstrated reserves. The deposit is open at depth (below 1,495 ft). The Condor-Hemlock zone is in the Snoqualmie batholith, northern phase granodiorite and tonalite that contains biotite and hornblende; locally it contains clinopyroxene. The rocks are light colored, medium crystalline, mostly equigranular with hypidiomorphic texture, and coarsely jointed. The northern phase is about 25 m.y. old, on the basis of interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Frizzell and others, 1984, p. 18).

**COMMENTS:** Both patented and unpatented claims are owned (1990) by United Cascade Mining Company, Inc. The deposit was explored by Anaconda Copper Co., Howe Sound Co., Climax Molybdenum Corp., Anaconda Co., Bear Creek Mining Co., Cities Services Minerals Corp., Westland Mines Ltd., Natural Resources Development Corp., Houston Oil and Minerals, and Electras Resources.

**UNPUBLISHED INFORMATION:** Grant, Alan R., 1965, Preliminary report Middle Fork Snoqualmie area, eastern King County, Washington. This report is in DGER files.

#### REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Grant, A. R., 1969, Chemical and physical controls for base metal deposition in the Cascade Range of Washington: Washington Division of Mines and Geology Bulletin 58, 107 p.
- Gualtieri, J. L.; Thurber, H. K.; Miller, M. S.; McMahan, A. B.; Federspiel, F. F., 1975, Mineral resources of additions to the Alpine Lakes study area, Chelan, King, and Kittitas Counties, Washington: U.S. Geological Survey Open-File Report 75-3, 161 p., 2 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- McIntyre, A. W., 1907, Copper deposits of Washington: American Mining Congress, 9th Annual Session, Report of Proceedings, p. 238-250.
- Purdy, C. P., Jr., 1954, Molybdenum occurrences of Washington: Washington Division of Mines and Geology Report of Investigations 18, 118 p., 6 pl.
- Western Miner, 1967, Westland Mines Ltd.: p. 128.

**Coney Basin (598)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 39' 7.63" N	121° 27' 41.94" W	N1/2 sec. 13, 25N, 10E and S1/2 sec. 19, 25N, 11E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Snoqualmie batholith	tonalite and granodiorite	Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopyrite	pyrite, quartz, arsenopyrite	
Ag	galena		
Cu	sphalerite		
Zn	tetrahedrite		
Pb	bournonite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			
stockwork			

PRODUCTION: Produced 40 tons in 1895; also produced in 1934, 1937-39, and 1941 (Hunting, 1956, p. 127).

TECTONIC SETTING: The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

ORE CONTROLS: The deposit consists of small persistent quartz veinlets along joint planes in granodiorite. A silicified and mineralized zone as much as 15 ft wide was also noted (Hunting, 1956, p. 127; Livingston, 1971, p. 164). Purdy (1951, p. 87), on the basis of the presence of antimony in an assay result, suggests that the probable antimony mineral is jamesonite, which is typical of other deposits in the area.

GEOLOGIC SETTING: The Coney Basin deposit is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and locally clinopyroxene. Age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Damon and Pythias (596)**

ALTERNATE NAMES		DISTRICT	COUNTY
Damon Pythias		Miller River Money Creek	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Si	1:62,500	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 41' 54.21" N	121° 31' 33.79" W	near center sec. 33, 26N, 10E	
LOCATION: at the head of Money Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Snoqualmie batholith		tonalite and granodiorite	Miocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopyrite	arsenopyrite, pyrite, quartz, tourmaline, calcite	
Ag	galena		
Pb	sphalerite		
Cu	tetrahedrite		
Zn			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Shipped 23 tons prior to 1940 (Hunting, 1956, p. 127). Ore shipped averaged 0.87 oz/ton Au, 9 oz/ton Ag, and 4% Pb (Moen, 1976, p. 167).

TECTONIC SETTING: The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

ORE CONTROLS: Similar to the Apex mine. Consists of two veins, one averaging 3 ft wide over a distance of 900 ft (Hunting, 1956, p. 127).

GEOLOGIC SETTING: The Damon and Pythias mines are in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and, locally, clinopyroxene. The age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

COMMENTS: This property is adjacent to the Apex property. Coats (1932, p. 17-19) describes the geology of the veins in both mines. Because the mines were operated by the same company at the time of Coats' study, some production from the Damon may be included in Apex production figures.

## REFERENCES

- Coats, R. R., 1932, The ore deposits of the Apex gold mine, Money Creek, King County, Washington: University of Washington Master of Science thesis, 48 p., 2 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.



**Dutch Miller (592)**

ALTERNATE NAMES		DISTRICT Snoqualmie	COUNTY King
PRIMARY QUADRANGLE Mount Daniel	SCALE 1:24,000	1/2° x 1° QUAD Skykomish River	1° x 2° QUAD Wenatchee
LATITUDE 47° 33' 32.65" N	LONGITUDE 121° 14' 19.34" W	SECTION, TOWNSHIP, AND RANGE NE¼ sec. 20, 24N, 13E	
LOCATION: at the head of the Middle Fork Snoqualmie River			
HOST ROCK: NAME Snoqualmie batholith	LITHOLOGY tonalite and granodiorite	AGE Miocene	
COMMODITIES Cu Au Ag	ORE MINERALS chalcopyrite tetrahedrite galena sphalerite	NON-ORE MINERALS arsenopyrite, pyrite, quartz, tourmaline, siderite, pink chlorite	
DEPOSIT TYPE vein breccia zone porphyry system		MINERALIZATION AGE	

PRODUCTION: Several shipments prior to 1901 (Hunting, 1956, p. 57).

TECTONIC SETTING: The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

ORE CONTROLS: Consists of three parallel, en echelon veins in granodiorite (Hunting, 1956, p. 57). Landes and others (1902, p. 86) report the ore vein has a maximum width of 18 1/2 ft. The presence of tourmaline and the proximity to an area of extensively explored porphyry copper mineralization (Clipper zone about 5 mi to the southwest) suggest this may be similar mineralization.

GEOLOGIC SETTING: The Snoqualmie batholith, northern phase, consists of granodiorite and tonalite with biotite and hornblende: it is medium crystalline and mostly equigranular and has hypidiomorphic texture. It locally contains clinopyroxene. It is light-colored, coarsely jointed rock. The age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Grant, A. R., 1969, Chemical and physical controls for base metal deposition in the Cascade Range of Washington: Washington Division of Mines and Geology Bulletin 58, 107 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II—The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Grand Central (590)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Money Creek	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 42' 26.67" N	121° 25' 43.09" W	sec. 29, 26N, 11E	
LOCATION: south of Money Creek. A more exact location is not available; however, the Grand Central may be part of a block of patented claims in the SW1/4SW1/4 sec. 29; SE1/4 sec. 30; E1/2 sec. 31; W1/2 sec. 32.			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation, Silver Pass Volcanic member		andesitic breccia	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb Au	stibnite	pyrite, calcite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced in 1908 (Hunting, 1956, p. 17).

TECTONIC SETTING: The Grand Central mine is located in a volcanic member of the Swauk Formation, which is early Eocene (52 m.y.) and pre-Cascade arc magmatism (Tabor 1982, geol. map, p. 13).

ORE CONTROLS: Stibnite is present in the upper workings of the mine. Narrow veinlets of quartz, pyrite, and calcite were observed on the lower level of the mine (Smith, 1915, p. 171, 185). These veinlets occur in a 40-ft-wide zone in andesite (Livingston, 1971, p. 157).

GEOLOGIC SETTING: The mine is in andesite of the Silver Pass Volcanic member of the Swauk Formation (Tabor and others, 1982, geol. map, p. 13). It is also at the contact between shales and andesite (Smith, 1913, p. 32-33).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Smith, W. S., 1913, The geology and mineral resources of north-eastern King County, Washington: Columbia University Master of Arts thesis, 35 p., 1 pl.
- Smith, W. S., 1915, Petrology and economic geology of the Skykomish Basin, Washington: Columbia University School of Mines Quarterly, v. 36, no. 2, p. 154-185.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Great Republic (591)**

ALTERNATE NAMES		DISTRICT	COUNTY
Happy Thought		Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 41' 54.75" N	121° 24' .10" W	N1/2SW1/4 sec. 33, 26N, 11E	
LOCATION: at the first falls on Happy Thought Creek about one-half mi from the Miller River			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation, Silver Pass Volcanic member		andesitic breccia	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb Au Ag	stibnite	pyrite, quartz, calcite, sericite, kaolinite, argillic alteration minerals	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced 1938-1941 (Hunting, 1956, p. 17).

**TECTONIC SETTING:** The Great Republic mine is in a volcanic member of the Swauk Formation, which is early Eocene (52 m.y.) and pre-Cascade magmatic arc (Tabor and others, 1982, geol. map, p. 13).

**ORE CONTROLS:** The best mineralization occurs along a gently dipping fault in andesite (Purdy, 1951, p. 75). Stibnite and pyrite are present in a quartz, calcite, and andesite gangue (Livingston, 1971, p. 157).

**GEOLOGIC SETTING:** Mineralization occurs in the Silver Pass Volcanic member of the Swauk Formation. The volcanic member consists of rhyolitic to andesitic breccia and feldspar porphyry (Tabor and others, 1982, p. 13).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Guye (601)**

ALTERNATE NAMES		DISTRICT	COUNTY
Mt. Logan Summit		Snoqualmie	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Snoqualmie Pass	62500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 26' 45.10" N	121° 25' 11.57" W	secs. 28, 32, and 33, 23N, 11E	
LOCATION: about a mile north of Snoqualmie Pass			
HOST ROCK: NAME	LITHOLOGY	AGE	
Naches Formation, Guye Sedimentary Member	sandstone, shale, and conglomerate	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Pb Zn Ag Au	magnetite galena sphalerite	pyrite, garnet	
DEPOSIT TYPE	MINERALIZATION AGE		
contact metamorphic			

**PRODUCTION:** Deposit has been extensively explored (Hunting, 1956, p. 196).

**TECTONIC SETTING:** The Snoqualmie batholith complex is part of early magmatism of the Cascade magmatic arc. Mineralization probably occurred during emplacement of a later phase of the batholith.

**ORE CONTROLS:** Where granodiorite is in contact with limestone, the limestone is recrystallized to marble and partially replaced by garnet and magnetite (Shedd and others, 1922, p. 90-92). A vein near the center of one of the claims, reported to be 6 ft wide, contains abundant galena and sphalerite (Hunting, 1956, p. 196).

**GEOLOGIC SETTING:** At the Guye mine, limestone and conglomerate of the Guye Sedimentary Member of the Naches Formation (Frizzell and others, 1984, geol. map) was intruded by granodiorite of the Snoqualmie batholith (Shedd and others, 1922, p. 90-92).

**REFERENCES**

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Shedd, Solon, 1924, The mineral resources of Washington with statistics for 1922; with an article on coal and coke, by G. W. Evans: Washington Division of Geology Bulletin 30, 224 p.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

**Quartz Creek (599)**

ALTERNATE NAMES		DISTRICT	COUNTY
Rainy Western States Copper		Taylor River Snoqualmie	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Si	1:62,500	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 34' 14.34" N	121° 33' 15.86" W	NW¼ sec. 16, SW¼ sec. 9, 24N, 10E	
LOCATION: on Quartz Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Snoqualmie batholith		tonalite and granodiorite	Miocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Mo Au Ag	chalcopyrite brannerite molybdenite scheelite	pyrite, pyrrhotite, arsenopyrite, quartz, tourmaline	
DEPOSIT TYPE		MINERALIZATION AGE	
breccia pipe stockwork porphyry copper			

**PRODUCTION:** One load of concentrates shipped to Tacoma in 1952. Also produced 1953-1955 (Hunting, 1956, p. 129). Concentrates shipped to smelter in 1942 (Livingston, 1971, p. 161). Reportedly produced 2,000 tons of ore between 1951 and 1957 and a flotation mill was constructed on the property (Report of Mineral Examination for 28 unpatented mineral claims of the Rainy property)

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** The property consists of two mineralized breccia pipes, called the east and west areas. The west pipe is the area of the old Rainy mine, and the east pipe is located about 800 ft to the east (Grant, 1969, p. 79-81). Grant also reports (p. 81) two rock types are directly related to zones with proportionately higher contents of sulfide minerals, biotite-quartz diorite replacement breccias, and a syenite-quartz monzonite-granite complex. He believes the replacement breccias are products of deuteric alteration and the syenite-quartz monzonite-granite complex rocks occur as matrix material within breccia blocks. The following description of the ore bodies is from Livingston (1971, p. 149, as obtained by personal commun. with A. R. Grant). The east pipe's surface dimension is about 300 by 600 ft, and it consists of fragments of quartz diorite that contain secondary biotite and a matrix of quartz and minor sulfide. The breccia is composed of angular fragments that range from a few inches to several feet long. The sulfides are scattered in breccia fragments and along fractures, and sulfides are generally finely divided in less mineralized parts of the pipe. Vugs range up to 2 in. and are lined with quartz. Quartz is much less abundant in more sulfide-rich parts of the pipe. The west zone is a zone of shattering and contains quartz veins along west-trending fractures. Sulfides are present in the quartz veins and as disseminations in the country rock.

**GEOLOGIC SETTING:** The Quartz Creek deposit is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and, locally, clinopyroxene. The age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

**COMMENTS:** An annual report for Inland Copper Ltd. reports the Quartz Creek property has an indicated reserve of 56,000 tons grading 5.5% Cu and 2.5 oz/ton Ag, or 905,000 tons of material averaging 1.1% Cu.

**REFERENCES**

- Grant, A. R., 1969, Chemical and physical controls for base metal deposition in the Cascade Range of Washington: Washington Division of Mines and Geology Bulletin 58, 107 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.

**Royal Reward (604)**

ALTERNATE NAMES		DISTRICT	COUNTY
			King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cumberland	1:24,000	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 19' .83" N	121° 55' 54.77" W	NW1/4SW1/4SW1/4 sec. 9, or SE1/4 SE1/4 sec. 8, 21N, 7E	
LOCATION: on a small terrace on the south bank of the Green River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Puget Group	sandstone, carbonaceous shale, coal	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
andesite dike			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg As Sb	cinnabar orpiment realgar stibnite	There are no reported alteration or gangue minerals.	
DEPOSIT TYPE		MINERALIZATION AGE	
vein disseminated			

**PRODUCTION:** The property produced about 20 flasks of mercury (Livingston, 1971, p. 154).

**TECTONIC SETTING:** The Eocene rocks were extensively folded and faulted prior to deposition of the cinnabar (Livingston, 1971, 153-154).

**ORE CONTROLS:** Two sets of faults cut the folded rocks; mineralization is present along both sets of faults. Cinnabar mineralization occurs in pods and disseminations in the fault breccia. Stibnite is common in the nearby Cardinal Reward property (Rice, 1962, private report in DGER files).

**GEOLOGIC SETTING:** The Puget Group consists of predominantly fluvial and some nearshore marine sandstone, siltstone, claystone, and coal (Frizzell and others, 1984, p. 26). Much of the mineralization occurs at the contact between sandstone and carbonaceous claystone (Rice, 1962, private report).

**UNPUBLISHED INFORMATION:** Rice, W. L., 1962, Summary of the geology and mineralization of the Cardinal Reward and Royal Reward mines, King County, Washington: Private Report for Northern Pacific Railway Company, 3 p. Livingston, V. E., 1957, Memorandum report on the 1957 examination of the Royal Reward mine, 3 p. These reports are in DGER files.

**REFERENCES**

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.

**Seattle-Cascade (603)**

ALTERNATE NAMES		DISTRICT	COUNTY
Triple S Silver Dollar Copper Plate Silver Dollar and Copper Plate		Miller River	King
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 40' 5.85" N	121° 25' 5.50" W	center of the N1/2 sec. 17, 25N, 11E	
LOCATION: west of the Miller River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Snoqualmie batholith	tonalite and granodiorite	Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Zn Cu Au	galena sphalerite chalcopyrite	pyrite, arsenopyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
mineralized shear zone vein			

**PRODUCTION:** Minor production prior to 1900; produced again in 1940 (Hunting, 1956, p. 297).

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** Mineralization occurs in a 1.5-ft-wide shear zone in granodiorite, in bands as much as 8 in. thick and which assay as much as 30 oz/ton Ag (Moen, 1976, p. 167).

**GEOLOGIC SETTING:** The Seattle-Cascade mine is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and, locally, clinopyroxene. The age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Snoqualmie (593)**

ALTERNATE NAMES		DISTRICT Miller River	COUNTY King
PRIMARY QUADRANGLE Grotto	SCALE 1:24,000	½° x 1° QUAD Skykomish River	1° x 2° QUAD Wenatchee
LATITUDE 47° 38' 40.73" N	LONGITUDE 121° 29' 17.61" W	SECTION, TOWNSHIP, AND RANGE secs. 14 and 23, 25N, 10E	
LOCATION: topographic map shows mines in the SE¼ sec. 14 and the NE¼ sec. 23. Location selected is the two portals in sec. 14			
HOST ROCK: NAME Snoqualmie batholith	LITHOLOGY tonalite and granodiorite	AGE Miocene	
COMMODITIES Cu	ORE MINERALS chalcopyrite	NON-ORE MINERALS pyrite	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: Produced in 1925 (Hunting, 1956, p. 59).

TECTONIC SETTING: The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

ORE CONTROLS: Gash veins are present in granodiorite. Some high-grade ore came from the No. 6 vein (Hunting, 1956, p. 59).

GEOLOGIC SETTING: The Snoqualmie batholith, northern phase, is granodiorite and tonalite with biotite and hornblende. It is medium crystalline, mostly equigranular, and hypidiomorphic. It locally contains clinopyroxene. It is light-colored, coarsely jointed rock. The age of the northern phase is about 25 m. y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.



**Una (567)**

ALTERNATE NAMES		DISTRICT	COUNTY
John Stevens Little Una		Miller River	King
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Grotto	1:24,000	Skykomish River	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 39' 32.44" N	121° 24' 51.54" W	SE¼ sec. 17, 25N, 11E	
LOCATION: on the West Fork Miller River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Snoqualmie batholith	tonalite and granodiorite	Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au	malachite chalcopyrite	tourmaline	
DEPOSIT TYPE		MINERALIZATION AGE	
vein breccia pipe			

**PRODUCTION:** Produced in 1908 (Hunting, 1956, p. 59).

**TECTONIC SETTING:** The Snoqualmie batholith is part of early magmatism of the Cascade magmatic arc.

**ORE CONTROLS:** A 20-ft body of tourmaline at the Una mine is described by Smith (1915, p. 184-185). The brief description suggests the tourmaline is massive crystalline, but may also be a portion of a breccia pipe.

**GEOLOGIC SETTING:** The Una mine is in rocks of the Snoqualmie batholith, northern phase, which consists of granodiorite and tonalite and is light colored, medium crystalline, and mostly equigranular and has hypidiomorphic texture. The rocks are coarsely jointed and contain biotite and hornblende and, locally, clinopyroxene. The age of the northern phase is about 25 m.y., based on interpretation of numerous discordant K-Ar ages of both hornblende and biotite (Tabor and others, 1982, p. 8).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1971, Geology and mineral resources of King County, Washington: Washington Division of Mines and Geology Bulletin 63, 200 p., 8 pl.
- Smith, W. S., 1915, Petrology and economic geology of the Skykomish Basin, Washington: Columbia University School of Mines Quarterly, v. 36, no. 2, p. 154-185.
- Tabor, R. W.; Frizzell, V. A., Jr.; Booth, D. B.; Whetten, J. T.; Waitt, R. B.; Zartman, R. E., 1982, Preliminary geologic map of the Skykomish River 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 82-747, 31 p., 1 pl.

**Boulder Creek (567)**

ALTERNATE NAMES		DISTRICT	COUNTY
Burke			Kittitas
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Kachess Lake	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 26' 18.00" N	121° 2' 31.22" W	SE¼NE¼ sec 35, 23N, 14E	
LOCATION: on the west slope of Mt. Hawkins			
HOST ROCK: NAME		LITHOLOGY	AGE
Ingalls Complex		serpentinite	Jurassic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cr	chromite	serpentine	
DEPOSIT TYPE		MINERALIZATION AGE	
podiform chromite		Jurassic	

PRODUCTION: Together with ore from the Mount Hawkins mine, one carload of ore was shipped (Hunting, 1956, p. 37).

TECTONIC SETTING: The Ingalls Complex is an ophiolite complex (Miller, 1985, p. 27).

ORE CONTROLS: Chromite is present as small lenses in serpentinitized peridotite (Hunting, 1956, p. 37). The chromite originated as a magmatic segregation in peridotite.

GEOLOGIC SETTING: The Ingalls Complex is an ophiolite complex of Middle to Late Jurassic age (Miller, 1985, p. 27-42).

## REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.

***Boulder Creek placer*** (584)

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 15' 16.15" N	120° 39' 27.21" W	sec. 1, 20N, 17E	
LOCATION: at the junction of Boulder and Williams Creeks			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel.	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Considerable production (Hunting, 1956, p. 185).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Cascade Chief (572)**

ALTERNATE NAMES		DISTRICT	COUNTY
Morrison First of August Gladstone		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenāṭchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 16' 52.98" N	120° 38' 18.14" W	SE1/4SW1/4 sec. 26, 21N, 17E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Swauk Formation	feldspathic to lithofeldspathic sandstone	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz, calcite, talcose mineral	
DEPOSIT TYPE	MINERALIZATION AGE		
mineralized shear zones vein			

**PRODUCTION:** Produced in 1911, 1938, and 1939 (Huntting, 1956, p. 131).

**TECTONIC SETTING:** Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

**ORE CONTROLS:** Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). At the Cascade Chief mine, stringers of mineralized quartz occur in three shear zones that average 4 ft wide in sandstone (Huntting, 1956, p. 131).

**GEOLOGIC SETTING:** Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

**REFERENCES**

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Clarence Jordin (573)**

ALTERNATE NAMES		DISTRICT	COUNTY
Gold King		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 15' 21.92" N	120° 38' 42.08" W	sec. 2, 20N, 17E	
LOCATION: on Snowshoe Ridge			
HOST ROCK: NAME	LITHOLOGY	AGE	
Swauk Formation	feldspathic to lithofeldspathic sandstone	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz, calcite	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Reported \$35,000 production. Produced from the Ace of Diamonds claim in 1952 (Hunting, 1956, p. 131).

TECTONIC SETTING: Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

ORE CONTROLS: Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80).

GEOLOGIC SETTING: Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Cle Elum River Iron (581)**

ALTERNATE NAMES		DISTRICT	COUNTY
Cle Elum River, north deposit Cle Elum River, south deposit Balfour Guthrie			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Kachess Lake	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 25' 53.42" N	121° 3' 16.25" W	secs. 26, 34, and 35, 23N, 14E and SE1/4 and NW1/4 sec. 1; N1/2 sec. 2; and E1/2 sec. 3, 22N, 14E	
LOCATION: on both sides of the Cle Elum River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex Swauk Formation	serpentine sandstone and shale	Jurassic Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Ni Cr	magnetite hematite millerite chrome spinel garnierite	serpentine and other weathered residuum from serpentinized peridotite	
DEPOSIT TYPE		MINERALIZATION AGE	
residual deposit			

**PRODUCTION:** There is no recorded production, but drilling (average from 52 holes) indicates a reserve of 6.25 million tons of 40.82% Fe, 0.83% Ni, and 2.4% Cr<sub>2</sub>O<sub>3</sub> (Huntting, 1956, p. 197).

**TECTONIC SETTING:** The Eocene Swauk Formation was deposited on tectonically emplaced, serpentinized Ingalls Complex.

**ORE CONTROLS:** Weathering of serpentinized peridotite prior to deposition of the Eocene Swauk Formation. The iron minerals derived from the peridotite remain in the residuum on the peridotite and in the basal beds of the Swauk Formation overlying and flanking the peridotite (Shedd and others, 1922, p. 77-79).

**GEOLOGIC SETTING:** Iron minerals are concentrated at the serpentinized peridotite-Swauk Formation contact.

**REFERENCES**

- Broughton, W. A., 1944, Economic aspects of the Blewett-Cle Elum iron ore zone, Chelan and Kittitas Counties, Washington: Washington Division of Geology Report of Investigations 12, 42 p., 7 pl.
- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

**Dolphin (570)**

ALTERNATE NAMES		DISTRICT	COUNTY
Bonanza			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mount Stuart	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 26' 41.54" N	120° 58' 40.52" W	sec. 33, 23N, 13E	
LOCATION: on the southeast slope of Hawkins Mountain. The adit shown on the topographic map in the N1/2 of the section is assumed to be the location of the mine.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	serpentinite	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au Co	chalcopyrite	pyrite, serpentine, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
silicified zone			

PRODUCTION: Produced in 1905 and again in 1907 about 5 tons of ore (Hunting, 1956, p. 60).

TECTONIC SETTING: The Ingalls Complex was accreted to the North American continent in the late Mesozoic (Miller, 1985, p. 27-42).

ORE CONTROLS: The deposit consists of a silicified zone as much as 50 ft wide in serpentinite (Hunting, 1956, p. 60). The deposit consists of copper and iron sulfides and could be a magmatic segregation deposit.

GEOLOGIC SETTING: The Ingalls Complex is an ophiolite complex of Middle to Late Jurassic age (Miller, 1985, p. 27-42).

## REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.

***Esther and Louisa*** (583)

ALTERNATE NAMES		DISTRICT	COUNTY
		Gold Creek	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Snoqualmie Pass	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 27' 38.69" N	121° 19' 16.51" W	N1/2NE1/4 sec. 27, 23N, 12E	
LOCATION: on the east side of a cirque at the head of Gold Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Naches Formation	andesite flows	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granodiorite		Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	pyrargyrite	pyrite, quartz	
Au	galena		
Pb	sphalerite		
Zn			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced about \$1,000 in 1896 (Hunting, 1956, p. 298).

TECTONIC SETTING: The Miocene igneous activity is part of the initial stages of Cascade magmatic arc development.

ORE CONTROLS: Ore is sporadically distributed but is locally of high grade in the vein (Hunting, 1956, p. 298; Moen, 1976, p. 142-143).

GEOLOGIC SETTING: The vein occurs near a wide granodiorite dike (Hunting, 1956, p. 298), which probably is part of the nearby Snoqualmie batholith (Tabor and others, 1982, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.



**Flodine (576)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 16' 52.88" N	120° 38' 21.59" W	sec. 25, 21N, 17E, and sec. 30, 21N, 18E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Swauk Formation	feldspathic to lithofeldspathic sandstone	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
fissure zone			
vein			

**PRODUCTION:** Produced several thousand dollars of gold from the oxidized zone prior to 1928 (Hunting, 1956, p. 132).

**TECTONIC SETTING:** Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

**ORE CONTROLS:** Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). Mineralization at the Flodine deposit is in a fissure zone (Hunting, 1956, p. 132).

**GEOLOGIC SETTING:** Eocene sandstone of the Swauk Formation is cut by numerous dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Francis Virdin Park (588)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Swauk Pass	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 16' 11.50" N	120° 36' 37.47" W	NW1/4 sec. 32, 21N, 18E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Swauk Formation	feldspathic to lithofeldspathic sandstone	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
silicified zone vein			

**PRODUCTION:** A small washing plant is currently (1990) operating. The operator is also recovering leaf-type gold specimens. The specimen gold is very light yellow and probably has a fairly high silver content.

**TECTONIC SETTING:** Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

**ORE CONTROLS:** Gold occurs at the center of quartz-calcite veinlets. Paragenetically, the gold is younger than the quartz and calcite.

**GEOLOGIC SETTING:** Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Golden Fleece (577)**

ALTERNATE NAMES		DISTRICT	COUNTY
Mercer T-Bone		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 18' 28.51" N	120° 39' 19.73" W	NE1/4SW1/4 sec. 13, 21N, 17E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation		feldspathic to lithofeldspathic sandstone	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
mineralized shear zone vein			

PRODUCTION: Reported production totals \$30,000 (Hunting, 1956, p. 132).

TECTONIC SETTING: Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

ORE CONTROLS: Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). At the Golden Fleece mine, a mineralized shear zone about 4 in. wide cuts beds of carbonaceous shale (Hunting, 1956, p. 132).

GEOLOGIC SETTING: Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**H-O-M-E (582)**

ALTERNATE NAMES		DISTRICT	COUNTY
Home			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Frost Mtn	1:24,000	Chelan	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 6' 13.05" N	120° 57' 25.05" W	secs. 26, 27, 28, and 34, 19N, 15E	
LOCATION: near the junction of Frosty and Taneum Creeks			
HOST ROCK: NAME	LITHOLOGY	AGE	
pre-Jurassic phyllite (Easton Schist) Manastash Formation	carbonaceous phyllite sandstone, shale, and conglomerate	pre-Jurassic Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg	cinnabar mercury (quicksilver)		
DEPOSIT TYPE	MINERALIZATION AGE		
vein shear zone			

**PRODUCTION:** This deposit has produced a reported 9.5 tons of ore (Hunting, 1956, p. 264).

**TECTONIC SETTING:** Mineralization probably took place following deformation of metamorphic rocks and deposition of the overlying Eocene rocks.

**ORE CONTROLS:** Cinnabar occurs in faults and fracture zones near the contact between carbonaceous phyllite and highly altered calcareous sedimentary rocks (Hunting, 1956, p. 264).

**GEOLOGIC SETTING:** Mineralization occurs scattered through older (pre-Jurassic) phyllite and calcareous sandstone of the Manastash Formation (Tabor and others, 1982, geol. map, p. 10).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Huckleberry (569)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Kittitas
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Kachess Lake	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 27' 41.39" N	121° 2' 21.39" W	secs. 24 and 26, 23N, 14E	
LOCATION: on the northwest flank of Huckleberry Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	diabase and gabbro	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au	chalcopyrite	quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Four carloads of ore reported shipped to Tacoma prior to 1935 (Hunting, 1956, p. 60).

TECTONIC SETTING: The Ingalls Complex was accreted to the North American continent in the late Mesozoic (Miller, 1985, p. 27-42).

ORE CONTROLS: The deposit is said to be a vein that ranges from 8 in. to 4 ft wide (Hunting, 1956, p. 60).

GEOLOGIC SETTING: The mineralization is hosted in diabase and gabbro of the Ingalls Complex.

## REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.

**Liberty (575)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 17' 49.37" N	120° 38' 25.68" W	NW1/4NE1/4 sec. 25, 21N, 17E, and sec. 19, 21N, 18E	
LOCATION: in Lyons Gulch			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation		feldspathic to lithofeldspathic sandstone	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
silicified zone vein			

PRODUCTION: Produced in 1935 and 1936 (Hunting, 1956, p. 132).

TECTONIC SETTING: Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

ORE CONTROLS: Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). At the Liberty deposit, a 4-ft-wide shear zone in carbonaceous shale contains stringers of quartz and calcite (Hunting, 1956, p. 132).

GEOLOGIC SETTING: Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Mineral Creek (571)**

ALTERNATE NAMES		DISTRICT	COUNTY
Durrwachter Liberty Lode			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Snoqualmie Pass	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 25' 29.40" N	121° 15' 15.71" W	SE 1/4 sec. 6, 22N, 13E	
LOCATION: on Mineral Creek 2 mi above Little Kachess Lake			
HOST ROCK: NAME	LITHOLOGY	AGE	
Naches Formation	sandstone and volcanic rocks	Eocene	
Easton schist	phyllite	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granodiorite dike (Snoqualmie batholith?)		Miocene	
COMMODITIES-	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite	pyrite, pyrrhotite, quartz	
Au	bornite		
Ag	molybdenite		
Mo			
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone disseminated			

**PRODUCTION:** 20 tons shipped to the Tacoma smelter prior to 1920 (Hunting, 1956, p. 60).

**TECTONIC SETTING:** Mineralization occurs in shear zones, along joint planes, and in a brecciated zone, which suggests mineralization took place following deformation of metamorphic rocks and deposition of the overlying Eocene rocks (Frizzell and others, 1984, geol. map; Hunting, 1956, p. 60).

**ORE CONTROLS:** Mineralization occurs in a narrow shear zone and in joint planes in a 20-40-ft-wide brecciated zone in a granodiorite dike (Patty, 1921, p. 277), and a breccia zone as much as 500 ft wide between rhyolite and basalt is mineralized (Hunting, 1956, p. 60).

**GEOLOGIC SETTING:** Host rocks for the deposit include a granodiorite dike probably related to the Snoqualmie batholith (Patty, 1921, p. 277-278) and rhyolitic and basaltic rocks probably of the Naches Formation (Frizzell, and others, 1984, geol. map).

**REFERENCES**

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Mount Hawkins (568)**

ALTERNATE NAMES		DISTRICT	COUNTY
Crowe Gallagher Head Skipper			Kittitas
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Mount Stuart	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 26' 30.20" N	120° 58' 56.88" W	SW¼NW¼ sec. 33, 23N, 15E	
LOCATION: in Drew Creek drainage on the southeast slope of Hawkins Mountain, near a small lake			
HOST ROCK: NAME	LITHOLOGY	AGE	
Ingalls Complex	serpentine	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cr	chromite	serpentine	
DEPOSIT TYPE		MINERALIZATION AGE	
podiform chromite		Jurassic	

**PRODUCTION:** The Mount Hawkins produced about 15 tons of ore during World War II (Huntting, 1956, p. 37-38).

**TECTONIC SETTING:** The Ingalls Complex is an ophiolite complex (Miller, 1985, p. 27). Chromite occurs in the ultramafic rocks.

**ORE CONTROLS:** Chromite is present as small lenses in serpentized peridotite (Huntting, 1956, p. 37). The chromite originated as a magmatic segregation in peridotite.

**GEOLOGIC SETTING:** The Ingalls Complex is an ophiolite complex of Middle to Late Jurassic age (Miller, 1985, p. 27-42).

**REFERENCES**

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.



**Mountain Daisy (579)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 15' 26.59" N	120° 38' 44.78" W	sec. 1, 20N, 17E and sec. 6, 20N, 18E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation		feldspathic to lithofeldspathic sandstone	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced from 1934 to 1938 (Hunting, 1956, p. 133).

TECTONIC SETTING: Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

ORE CONTROLS: Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80).

GEOLOGIC SETTING: Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

Kittitas

**Old Bigney placer (585)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 15' 10.85" N	120° 39' 59.06" W	W1/2 sec. 1, 20N, 17E	
LOCATION: near Liberty			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: More than \$200,000 prior to 1903; production reported for 1908, 1915, 1916, and 1923 (Hunting, 1956, p. 186).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

UNPUBLISHED INFORMATION:

REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Ollie Jordin (574)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Liberty	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 15' 30.95" N	120° 39' 2.36" W	sec. 2, 20N, 17E	
LOCATION: about 0.75 mi up Williams Creek from Liberty			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation		feldspathic to lithofeldspathic sandstone	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
silicified zone vein			

**PRODUCTION:** Produced a total of about \$20,000 in gold during 2 years prior to 1920 (Hunting, 1956, p. 133)

**TECTONIC SETTING:** Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

**ORE CONTROLS:** Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). At the Ollie Jordin mine, mineralization is present in a silicified zone that is 4 ft wide. Wire gold is found at the deposit (Hunting, 1956, p. 133).

**GEOLOGIC SETTING:** Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Silver Creek (580)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Kachess Lake	1:62,500	Snoqualmie Pass	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 29' 37.67" N	121° 1' 45.38" W	sec. 12, 23N, 14E	
LOCATION: in the Fish Lake area			
HOST ROCK: NAME		LITHOLOGY	AGE
Ingalls Complex		serpentinite	Jurassic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced in 1937 and 1938 (Hunting, 1956, p. 134).

TECTONIC SETTING: The Ingalls Complex was accreted to the North American continent in the late Mesozoic (Miller, 1985, p. 27-42).

ORE CONTROLS: Mineralization is irregularly distributed in a 15-20-ft wide quartz vein (Hunting, 1956, p. 133-134).

GEOLOGIC SETTING: The Ingalls Complex is an ophiolite complex of Middle to Late Jurassic age (Miller, 1985, p. 27-42).

## REFERENCES

- Frizzell, V. A., Jr.; Tabor, R. W.; Booth, D. B.; Ort, K. M.; Waitt, R. B., 1984, Preliminary geologic map of the Snoqualmie Pass 1:100,000 quadrangle, Washington: U.S. Geological Survey Open-File Report 84-693, 42 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Miller, R. B., 1985, The ophiolitic Ingalls Complex, north-central Cascade mountains, Washington: Geological Society of America Bulletin, v. 96, no. 1, p. 27-42.

**Silver Tip (589)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Frost Mtn	1:24,000	Chelan	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 6' 15.22" N	120° 57' 57.91" W	SE1/4 sec. 28, 19N, 15E	
LOCATION: near the junction of Frosty and Taneum Creeks			
HOST ROCK: NAME	LITHOLOGY	AGE	
pre-Jurassic phyllite (Easton Schist) Manastash Formation	carbonaceous phyllite sandstone, shale, and conglomerate	pre-Jurassic Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg	cinnabar mercury (quicksilver)		
DEPOSIT TYPE	MINERALIZATION AGE		
vein shear zone			

PRODUCTION: Deposit has reported production of 5 tons of ore (Hunting, 1956, p. 264).

TECTONIC SETTING: Mineralization probably occurred following deformation of metamorphic rocks and deposition of overlying Eocene rocks.

ORE CONTROLS: Cinnabar occurs in faults and fracture zones near the contact between carbonaceous phyllite and highly altered calcareous sedimentary rocks. At the Silver Tip, the shear zone is about 12 ft wide at the contact between the phyllite and sandstone; mineralization consists of small specks and thin seams of cinnabar (Hunting, 1956, p. 264).

GEOLOGIC SETTING: Mineralization occurs scattered through older (pre-Jurassic) phyllite and calcareous sandstone of the Manastash Formation (Tabor and others, 1982, geol. map, p. 10).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Swauk Creek placers (586)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Swauk Prairie	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 12' 49.17" N	120° 41' 58.58" W		
LOCATION: Hunting (1956, p. 187) reports the Swauk Creek placers are along Swauk Creek between the mouths of Baker and First Creeks. Only Baker Creek is identified on the topographic map; consequently, the Swauk placers are placed just south of the mouth of Baker Creek.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Considerable production (Huntting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Wall Street (578)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Swauk Pass	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 17' 19.28" N	120° 37' 28.49" W	sec. 30, 21N, 17E	
LOCATION: 5.5 mi up Cougar Gulch from Liberty; the last 1.5 mi is by trail.			
HOST ROCK: NAME		LITHOLOGY	AGE
Swauk Formation		feldspathic to lithofeldspathic sandstone	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
silicified fracture zone vein			

PRODUCTION: Produced \$50,000 worth of ore prior to 1935; also produced in 1938 (Hunting, 1956, p. 134).

TECTONIC SETTING: Quartz veins of the Swauk district are subparallel to north-northeast-trending basalt dikes in the area. The veins also cut some basalt dikes (Smith, 1904, p. 9).

ORE CONTROLS: Gold-quartz veins in the Swauk district are subparallel to numerous basalt dikes of the region (Smith, 1903, p. 80). At the Wall Street mine, gold mineralization is present in silicified fracture zones in sandstone (Hunting, 1956, p. 134).

GEOLOGIC SETTING: Eocene sandstone of the Swauk Formation is cut by numerous basalt dikes and gold-quartz veins. The gold-quartz veins appear to be localized along the same fracture system as the dikes.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Smith, G. O., 1903, Gold mining in central Washington. *In* Contributions to economic geology, 1902: U.S. Geological Survey Bulletin 213, p. 76-80.
- Smith, G. O., 1904, Geological atlas of the United States—Mount Stuart folio, Washington: U.S. Geological Survey Geologic Folio 106, 10 p.
- Tabor, R. W.; Waitt, R. B.; Frizzell, V. A., Jr.; Swanson, D. A.; Byerly, G. R.; Bentley, R. D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, central Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-1311, 1 sheet, scale 1:100,000, with 26 p. text.

**Williams Creek placers (587)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Swauk	Kittitas
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Swauk Prairie	1:24,000	Wenatchee	Wenatchee
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 14' 45.58" N	120° 40' 55.61" W	secs. 2, 10, and 11, 20N, 17E	
LOCATION: along Williams Creek between the junction with Swauk Creek and Liberty			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Considerable production (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.



**Barnum-McDonnell (632)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Morton	Lewis
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Glenoma	1:24,000	Centralia	Hoquiam
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 32' 46.69" N	122° 13' 59.87" W	N1/2NW1/4 sec. 7, 12N, 5E	
LOCATION: about 2.25 mi southeast of Morton			
HOST ROCK: NAME	LITHOLOGY	AGE	
Puget Group	shale, siltstone, sandstone	Eocene	
Puget Group	basic dikes and sills	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg	cinnabar		
DEPOSIT TYPE	MINERALIZATION AGE		
shear zone			
vein and veinlet			

PRODUCTION: Produced 75 flasks of mercury in 1916, 489 flasks in 1926, 1,265 flasks 1926-1929, and 441 flasks in 1931 (Hunting, 1956, p. 265).

TECTONIC SETTING: Deltaic deposits of the Eocene Puget Group were folded and faulted during the Tertiary (Gard, 1968, p. 27-29). Mineralization probably took place during or shortly after deformation.

ORE CONTROLS: Cinnabar occurs in fractures and breccia along fault zones in sandstone, shale, and siltstone of the Puget Group. Mafic rocks intruded the sediments, and all were subjected to folding and faulting prior to mineralization (Mackin, 1944, p. 8). The shear zone that localizes mineralization at the Barnum-McDonnell deposit has a maximum thickness of 10 ft. The best ore is in tuffaceous sandstone (Hunting, 1956, p. 265).

GEOLOGIC SETTING: The sedimentary rocks of the Puget Group contain coal, and fossils suggest brackish water and estuarine depositional conditions (Mackin, 1944, p. 7).

## REFERENCES

- Gard, L. M., Jr., 1968, Bedrock geology of the Lake Tapps quadrangle, Pierce County, Washington: U.S. Geological Survey Professional Paper 388-B, 33 p., 2 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mackin, J. H., 1944, Relation of geology to mineralization in the Morton cinnabar district, Lewis County, Washington: Washington Division of Mines and Mining Report of Investigations 6, 47 p., 2 pl.
- Schasse, H. W., compiler, 1987, Geologic map of the Centralia quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 87-11, 28 p., 1 pl.

**Eagle Peak (631)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Lewis
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mt Rainier West	1:24,000	Mt Rainer	Yakima
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 45' 20.96" N	121° 46' 48.47" W	near the NW corner sec. 27, 15N, 8E	
LOCATION: on the west slope of Eagle Peak in the Mt. Rainier area			
HOST ROCK: NAME	LITHOLOGY	AGE	
Stevens Ridge Formation	rhyodacite ash-flow tuff and volcaniclastic rocks	Oligocene - Miocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
undifferentiated felsic rocks related to the Tatoosh pluton		Miocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite	pyrite, arsenopyrite	
Au	bornite		
Ag	covellite		
Co	scheelite		
U	molybdenite		
W	sphalerite		
	linnaeite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein and veinlet			

**PRODUCTION:** Produced 100 tons in 1919; also produced in 1925 and 1928, but total production does not exceed 200 tons. An 18-ton shipment yielded 8.05% Cu and 0.09 oz/ton Au. Reported 1% to 5% Co in some samples (Hunting, 1956, p. 61).

**TECTONIC SETTING:** Rocks are part of the Cascade magmatic arc.

**ORE CONTROLS:** Mineralization occurs along joints or slip planes in granite. The ore zone ranges from 0.5 to 5 ft in width; the zone carries a streak of high-grade ore that is 1 to 14 in. wide. Gold values are associated with the arsenopyrite. The ore from near the portal of an old drift about 15 ft above the present drift was slightly radioactive (Hunting, 1956, p. 61).

**GEOLOGIC SETTING:** Mineralization occurs in granite (Hunting, 1956, p. 61) and would appear to be in dikes emanating from the Tatoosh pluton (Schasse, 1987, p. 29).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Schasse, H. W., compiler, 1987, Geologic map of the Mount Rainier quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 87-16, 43 p., 1 pl.

**Lytle-Lynch (634)**

ALTERNATE NAMES		DISTRICT	COUNTY
Charlotte Ann Kropolis		Morton	Lewis
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Glenoma	1:24,000	Centralia	Hoquiam
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 33' 32.18" N	122° 14' 3.39" W	NW¼ and N½NE¼ sec. 6, 12N, 5E	
LOCATION: about 0.75 mi north of the Roy and Barnum-McDonnell mines			
HOST ROCK: NAME	LITHOLOGY	AGE	
Puget Group	shale, siltstone, sandstone	Eocene	
Puget Group	basic dikes and sills and porphyry	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg	cinnabar	pyrite or marcasite	
DEPOSIT TYPE	MINERALIZATION AGE		
fault breccia vein and veinlet			

**PRODUCTION:** Produced from 1930 to 1932 (Hunting, 1956, p. 265).

**TECTONIC SETTING:** Deltaic deposits of the Eocene Puget Group were folded and faulted during the Tertiary (Gard, 1968, p. 27-29). Mineralization probably took place during or shortly after deformation.

**ORE CONTROLS:** Cinnabar is present in fractures and breccia along fault zones in sandstone, shale, and siltstone of the Puget Group. Mafic rocks intrude the sedimentary rocks, and all were subjected to folding and faulting prior to mineralization (Mackin, 1944, p. 8).

**GEOLOGIC SETTING:** The sedimentary rocks of the Puget Group contain coal and fossils that suggest brackish water and estuarine depositional conditions (Mackin, 1944, p. 7). At the Lytle-Lynch the country rocks are shale, sandstone, coal, and porphyry (Hunting, 1956, p. 265).

**REFERENCES**

- Gard, L. M., Jr., 1968, Bedrock geology of the Lake Tapps quadrangle, Pierce County, Washington: U.S. Geological Survey Professional Paper 388-B, 33 p., 2 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mackin, J. H., 1944, Relation of geology to mineralization in the Morton cinnabar district, Lewis County, Washington: Washington Division of Mines and Mining Report of Investigations 6, 47 p., 2 pl.
- Schasse, H. W., compiler, 1987, Geologic map of the Centralia quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 87-11, 28 p., 1 pl.

**Mineral Creek (635)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Lewis
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Anderson Lake	1:24,000	Centralia	Hoquiam
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 39' 59.89" N	122° 6' 1.92" W	SW¼ sec. 30, 14 N, 6E	
LOCATION: on Mineral Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed	basaltic andesite and andesite flows	Eocene - Oligocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
As	sphalerite		
Zn	galena		
Pb	realgar		
Ag			
DEPOSIT TYPE	MINERALIZATION AGE		
hydrothermal			

**PRODUCTION:** Reportedly 1,000 tons of arsenic ore produced in 1903 and some production in 1904 and 1905. A sample said to come from a 300-ft face assayed 3.8% Zn, 3.6% Pb, and 2.56 oz/ton Ag (Hunting, 1956, p. 363).

**TECTONIC SETTING:** Rocks of the Mineral Creek mine area are the earliest stages of Cascade magmatic arc volcanism (Walsh and others, 1987, correlation diagram; Swanson and others, 1989, p. 5-7).

**GEOLOGIC SETTING:** Host rocks at the Mineral Creek deposit are in a sequence of Eocene to Oligocene basaltic andesite and andesite flows (Schasse, 1987, geol. map).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Schasse, H. W., compiler, 1987, Geologic map of the Centralia quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 87-11, 28 p., 1 pl.
- Swanson, D. A.; Cameron, K. A.; Evarts, R. C.; Pringle, P. T.; Vance, J. A., 1989, Excursion 1A—Cenozoic volcanism in the Cascade Range and Columbia Plateau, southern Washington and northernmost Oregon. In Chapin, C. E.; Zidek, Jiri, editors, Field excursions to volcanic terranes in the western United States, Volume II—Cascades and intermountain west: New Mexico Bureau of Mines and Mineral Resources Memoir 47, p. 1-50.
- Walsh, T. J.; Korosec, M. A.; Phillips, W. M.; Logan, R. L.; Schasse, H. W., 1987, Geologic map of Washington—Southwest quadrant: Washington Division of Geology and Earth Resources Geologic Map GM-34, 2 sheets, scale 1:250,000, with 28 p. text.

**Roy (633)**

ALTERNATE NAMES		DISTRICT	COUNTY
Fisher Morton Gillispie		Morton	Lewis
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Glenoma	1:24,000	Centralia	Hoquiam
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
46° 32′ 52.90″ N	122° 13′ 57.88″ W	S½SW¼ and NW¼SW¼ sec. 6, 12N, 5E	
LOCATION: 2 mi southeast of Morton			
HOST ROCK: NAME	LITHOLOGY	AGE	
Puget Group Puget Group	shale, siltstone, sandstone basic dikes and sills	Eocene Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Hg	cinnabar	pyrite, calcite	
DEPOSIT TYPE	MINERALIZATION AGE		
vein and veinlet breccia zones			

**PRODUCTION:** Produced about 2,500 flasks in 1928-1929, 1,079 flasks in 1930, 1,581 tons of ore in 1931, and recovered 65 flasks of mercury from 1,000 tons of ore in 1940. Also produced in 1941 (Hunting, 1956, p. 265).

**TECTONIC SETTING:** Deltaic deposits of the Eocene Puget Group were folded and faulted during the Tertiary (Gard, 1968, p. 27-29). Mineralization probably took place during or shortly after deformation.

**ORE CONTROLS:** Cinnabar occurs in fractures and breccia along fault zones in sandstone, shale, and siltstone of the Puget Group. Mafic rocks intruded the sedimentary rocks, and all were subjected to folding and faulting prior to mineralization (Mackin, 1944, p. 8). The shear zone has a maximum probable thickness of 10 ft (Hunting, 1956, p. 265).

**GEOLOGIC SETTING:** The sedimentary rocks of the Puget Group contain coal and fossils that suggest brackish water and estuarine depositional conditions (Mackin, 1944, p. 7).

**REFERENCES**

- Gard, L. M., Jr., 1968, Bedrock geology of the Lake Tapps quadrangle, Pierce County, Washington: U.S. Geological Survey Professional Paper 388-B, 33 p., 2 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Mackin, J. H., 1944, Relation of geology to mineralization in the Morton cinnabar district, Lewis County, Washington: Washington Division of Mines and Mining Report of Investigations 6, 47 p., 2 pl.
- Schasse, H. W., compiler, 1987, Geologic map of the Centralia quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 87-11, 28 p., 1 pl.

***Barnell placer*** (298)

ALTERNATE NAMES		DISTRICT	COUNTY
			Lincoln
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Keller Ferry	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 56' 3.10" N	118° 42' 4.85" W	sec. 7, 28N, 33E	
LOCATION: in Swawilla basin, near Plum, 0.5 mi below the ferry			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced \$200-\$400 per week in 1938 (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**China Bar placer** (299)

ALTERNATE NAMES		DISTRICT	COUNTY
			Lincoln
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Olsen Canyon	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 50' 48.20" N	118° 21' 6.08" W	secs. 12 and 13, 27N, 35E	
LOCATION: on the east side of the Columbia River			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced in 1928. Said to have been worked out by Chinese miners (Huntting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

- Collier, A. J., 1907, Gold-bearing river sands of northeastern Washington. *In* Contributions to economic geology 1906; Part I—Metals and nonmetals, except fuels: U.S. Geological Survey Bulletin 315, p. 56-70.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Clark placer** (300)

ALTERNATE NAMES		DISTRICT	COUNTY
			Lincoln
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Keller Ferry	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 56' 28.61" N	118° 40' 52.21" W	SE¼NE¼ sec. 8, 28N, 33E	
LOCATION: along the Columbia River			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced \$4,657 in 1933, and \$8,243 from 19,700 yards of gravel in 1934 (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.



**Keller Ferry placer (301)**

ALTERNATE NAMES		DISTRICT	COUNTY
Angle Placer			Lincoln
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Keller Ferry	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 56' 8.70" N	118° 40' 51.57" W	E½ sec. 8, 28N, 33E	
LOCATION: opposite the mouth of the Sanpoil River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced in 1932, and 186.8 oz of Au from 11,628 yd<sup>3</sup> of gravel in 1933-1934 (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCE

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Spokane Molybdenum (297)**

ALTERNATE NAMES		DISTRICT	COUNTY
Pitney Butte Egypt		Deer Trail Cedar Canyon	Lincoln
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
McCoy Lake	1:24,000	Coulee Dam	Ritzville
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 52' 51.56" N	118° 9' 50.56" W	NE1/4SE1/4 sec. 32, 28N, 37E	
LOCATION: on the northeast side of Pitney Butte			
HOST ROCK: NAME	LITHOLOGY	AGE	
granite of Pitney Butte	monzogranite	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
andesite dike of Sanpoil Volcanics		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mo Au Ag	molybdenite chalcopyrite sphalerite pitchblende	quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced in 1941 (Hunting, 1956, p. 269).

**TECTONIC SETTING:** Cretaceous granitic rocks intrude rocks of the Kootenay arc and adjacent terranes in northeastern Washington (Rhodes and Hyndman, 1988).

**ORE CONTROLS:** Quartz that is found in muscovite-rich pegmatite veins is of two types: (1) milky white quartz and (2) gray, massive quartz in segregations. Both types of quartz contain sparse flakes of molybdenite (Becraft and Weis, 1963, p. 67-68). A cross fracture in the main vein contains lenses of black radioactive material (Hunting, 1956, p. 269).

**GEOLOGIC SETTING:** At the Spokane Molybdenum mine the Cretaceous, fine-grained monzogranite is cut by a dike of the Sanpoil Volcanics (Becraft and Weis, 1963, p. 67).

**REFERENCES**

- Becraft, G. E.; Weis, P. L., 1963, Geology and mineral deposits of the Turtle Lake quadrangle, Washington: U.S. Geological Survey Bulletin 1131, 73 p., 6 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1954, Molybdenum occurrences of Washington: Washington Division of Mines and Geology Report of Investigations 18, 118 p., 6 pl.
- Rhodes, B. P.; Hyndman, D. W., 1988, Regional metamorphism, structure, and tectonics of northeastern Washington and northern Idaho. In Ernst, W. G., editor, Metamorphism and crustal evolution of the western United States: Prentice-Hall [Englewood Cliffs, N.J.], Rubey Volume VII, p. 271-295.

**Black Hump (619)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Mason
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mt Steel	1:62,500	Mt. Olympus	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 31' 51" N	123° 19' 41" W	N1/2N1/2 sec. 33, 24N, 5W	
LOCATION: about 1 mi north of the Lincoln guard station, 2 mi by trail north of Staircase Resort at the head of Lake Cushman			
HOST ROCK: NAME	LITHOLOGY	AGE	
Needles-Gray Wolf lithic assemblage unnamed basalt	micaceous sandstone basalt	Eocene Eocene - Oligocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	bementite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene - Oligocene	

PRODUCTION: Some production prior to 1924 (Hunting, 1956, p. 260).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312). Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: The Crescent Formation on the Olympic Peninsula consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts (Snively, 1987, p. 306-308). Clastic rocks and associated basalts overlie the Crescent Formation host manganese mineralization at the Black Hump deposit (Tabor and Cady, 1978, geol. map).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.
- Tabor, R. W.; Cady W. M., 1978, Geologic map of the Olympic Peninsula, Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-994, 2 sheets, scale 1:125,000.

**Black and White (617)**

ALTERNATE NAMES		DISTRICT	COUNTY
Three Friends			Mason
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mt Steel	1:62,500	Mt. Olympus	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 34' 31" N	123° 20' 40" W	sec. 17 or NW 1/4 sec. 21, 24N, 5W	
HOST ROCK: NAME		LITHOLOGY	AGE
unnamed unit		foliated sandstone and semischist	Eocene - Oligocene
unnamed basalt		altered basalt	Eocene - Oligocene
ASSOCIATED IGNEOUS ROCK: DESCRIPTION			AGE
altered basalt			Eocene - Oligocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	copper	jasper	
Mn	cuprite		
Ag	chalcocite		
Cu	chalcotrichite		
Ni	malachite		
Mg	azurite		
	bementite		
	neotocite		
	rhodonite		
	rhodochrosite		
	manganocalcite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene - Oligocene	

**PRODUCTION:** Produced 5 tons of ore in 1915 containing 0.40 oz/ton Ag, 7.85% Cu, 3.2% Fe, and 65% insoluble (Hunting, 1956, p. 62).

**TECTONIC SETTING:** Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

**ORE CONTROLS:** Mineralization occurs along the contact between altered basalt and foliated sandstone-semischist (data from USGS MRDS, 1990; Tabor and Cady, 1978, geol. map). The deposit is in lenses that pinch and swell. Its maximum width is 8 ft, and the thickness is 50 ft. Lenticular bodies of manganese silicates occur conformably along the eastward or upper contacts of the basalt members at elevations approaching 4,400 ft. A manganiferous lens at the collar of the shaft pinches and swells along strike and down dip. No red limestone was exposed near the lens, but the limestone is found in the same stratigraphic horizon as the mineralization a short distance to the northeast. Manganese minerals are largely buff, tan, and gray bementites; also present are several thin veinlets of neotocite (data from USGS MRDS, 1990).

**GEOLOGIC SETTING:** The principal country rocks in the area are altered basalts and argillite, graywacke, limestone, and some phyllite. Near Kidney (Black and White) Lakes, they trend approximately N20E and dip steeply eastward.

**REFERENCES**

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Green, S. H., 1945, Manganese deposits of the Olympic Peninsula, Washington: Washington Division of Mines and Mining Report of Investigations 7, 45 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.

- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.
- Tabor, R. W.; Cady W. M., 1978, Geologic map of the Olympic Peninsula, Washington: U.S. Geological Survey Miscellaneous Investigations Series Map I-994, 2 sheets, scale 1:125,000.

**Triple Trip** (618)

ALTERNATE NAMES		DISTRICT	COUNTY
Brown Mule McKean			Mason
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mt. Steel	1:62,500	Mt. Olympus	Seattle
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
47° 30' 14" N	123° 19' 12" W	secs. 4 and 9 23N, 5W	
LOCATION: elev. 1,000 ft, 1 mi from Lincoln guard station along Apex trail, 0.25 mi up Copper Creek from the North Fork Skokomish River; on the north bank of Copper (Boulder) Creek; near the northwest corner of Lake Cushman.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Crescent Formation	red limestone	Eocene	
Crescent Formation	basalt	Eocene	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Crescent Formation volcanic rocks		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn Fe	bementite manganiferous magnetite		
DEPOSIT TYPE		MINERALIZATION AGE	
replacement, disseminated exhalative/diagenetic		Eocene	

PRODUCTION: A carload of ore produced during World War I ran 35% to 45% Mn and 17% to 30% SiO<sub>2</sub> (Hunting, 1956, p. 261).

TECTONIC SETTING: Host rocks of manganese deposits of the Olympic Peninsula formed at an ocean ridge or in a back arc basin at the active margin of the North American plate (Garrison, 1973; Snively, 1987, p. 306-309).

ORE CONTROLS: Lens-shaped ore body occurs in red limestone about 1 ft from its contact with basalt (data from U.S.G.S. MRDS, 1990; Hunting, 1956, p. 261). Manganese mineralization consists of a tabular body of bementite about 1.5 ft wide, 70 ft long, and 30 ft thick. Small amounts of secondary manganese oxides and jasper are associated with the bementite. Other small masses of manganese mineralization were found along Copper Creek to the southwest (data from USGS MRDS, 1990). Manganese deposits of the region are associated with spilites (Park, 1942, p. 311-312) and commonly are found with reddish pelagic limestone interbedded with the basalt. Manganese mineralization may be either disseminated in replacement bodies or volcanogenic exhalative/diagenetic bodies (Sorem and Gunn, 1967).

GEOLOGIC SETTING: The Crescent Formation, which hosts manganese mineralization on the Olympic Peninsula, consists of Paleocene? and Eocene pillow basalts compositionally similar to ocean ridge basalts. Associated pelagic limestones were deposited in deep water (Snively, 1987, p. 306-308).

## REFERENCES

- Garrison, R. E., 1973, Space-time relations of pelagic limestones and volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin, v. 84, no. 2, p. 583-594.
- Green, S. H., 1945, Manganese deposits of the Olympic Peninsula, Washington: Washington Division of Mines and Mining Report of Investigations 7, 45 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Magill, E. A., 1960, Manganese deposits of the Olympic Peninsula, Wash.: U.S. Bureau of Mines Report of Investigations 5530, 82 p.
- Park, C. F., Jr., 1942, Manganese resources of the Olympic Peninsula, Washington—A preliminary report: U.S. Geological Survey Bulletin 931-R, p. 435-457, 2 pl.
- Snively, P. D., Jr., 1987, Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin. In Scholl, D. W.; Grantz, Arthur; Vedder, J. G., Geology and resources potential of the continental margin of western North America and adjacent ocean basins—Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, p. 305-335.
- Sorem, R. K.; Gunn, D. W., 1967, Mineralogy of manganese deposits, Olympic Peninsula, Washington: Economic Geology, v. 62, no. 1, p. 22-56.

**49th Parallel (369)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 59' 45.89" N	119° 29' 30" W	NE¼ sec. 6, 40N, 27E	
LOCATION: 1,500 ft northwest of the O.K. copper mine			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation (?)	greenstone	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Silver Nail Lake pluton		Jurassic - Cretaceous (?)	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au W	chalcopyrite bornite scheelite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
porphyry vein			

PRODUCTION: Gold-copper ore in 1914 (Hunting, 1956, p. 65).

TECTONIC SETTING: The Kobau Formation was deposited along an active continental margin proximal to an island arc.

ORE CONTROLS: The 49th Parallel deposit is in the chlorite alteration zone of the Kelsey porphyry copper-molybdenum deposit (Roper, 1973).

GEOLOGIC SETTING: The 49th Parallel mine is in greenstone tentatively identified as the Permian or Triassic (?) Kobau Formation and also is adjacent to an altered quartz diorite phase of the Jurassic-Cretaceous (?) Silver Nail Lake pluton (Roper, 1973).

COMMENTS: Part of the Kelsey porphyry copper-molybdenum deposit (Roper, 1973).

## REFERENCES

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Horton, F. W., 1916, Molybdenum, its ores and their concentration, with discussion of markets, prices, and uses: U.S. Bureau of Mines Bulletin 111, 132 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Roper, M. W., 1973, Geology of the Kelsey copper-molybdenum property, Okanogan County, Washington: Montana State University Master of Science thesis, 97 p., 5 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Adelia** (375)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 52' 24.05" N	119° 34' .26" W	SE1/4 sec. 16, 39N, 26E	
LOCATION: on Palmer Mountain east of the Ivanhoe property			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	slate	Permian	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold	quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Several shipments prior to 1902 (Hunting, 1956, p. 135).

TECTONIC SETTING: Late Paleozoic sedimentary rocks were deposited along an active continental margin.

ORE CONTROLS: A vein about 5 ft wide in slate was said to carry good values in gold and silver (Hunting, 1956, p. 135).

GEOLOGIC SETTING: Vein in fine-grained metasedimentary rocks of the Permian Spectacle Formation of the Anarchist Group (Rinehart and Fox, 1972, geol. map).

## REFERENCES

- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.



**Alder** (452)

ALTERNATE NAMES		DISTRICT	COUNTY
		Twisp	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Twisp West	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 19' 19.46" N	120° 9' 28.72" W	secs. 25, 26, 35, and 36, 33N, 21E	
LOCATION: elev. 3,600 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Newby Group		dacite and rhyolite breccia	Jurassic - Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopryrite	pyrrhotite; silicification, chloritization, sericitization	
Ag	gold		
Cu	copper		
Zn	sphalerite		
	chalcocite		
	galena		
	malachite		
	azurite		
DEPOSIT TYPE		MINERALIZATION AGE	
massive sulfide vein		Jurassic - Cretaceous (?)	

**PRODUCTION:** Production includes 6,831 tons shipped in 1939 that averaged 0.55 oz/ton Au, about 0.50 oz/ton Ag, and 0.16% to 0.55% Cu (Hunting, 1956). Approximately 9,000 tons of ore were shipped in 1940 and 4,000 tons in 1942; 1,899 tons of concentrate were shipped in 1950 (Burnet, 1976). In 1951, 1,546 tons of copper concentrate yielded 268,202 lb Cu, and 2,072 oz Au. Shipments were also made in 1952-1953 (Hunting, 1956, p. 135).

**TECTONIC SETTING:** Newby Group rocks were probably deposited along or proximal to an island arc.

**ORE CONTROLS:** Sulfide deposits in the Alder mine are concordant with bedding in the host rocks and are strata-bound; hydrothermal alteration at the mine is also strata-bound. Sulfide minerals, including pyrite, sphalerite, chalcopryrite, and galena, are found in veins. These veins are generally small, have no relation to the Alder stock that intrudes volcanic rocks near the mine, and have a random orientation (Burnet, 1976; Bunning, 1990).

**GEOLOGIC SETTING:** Dacitic breccias of the Jurassic-Cretaceous Newby Group are host to volcanogenic massive sulfide mineralization at the Alder mine. The Newby Group was intruded by the Alder Creek stock, which has been dated at  $137 \pm 3.4$  m.y. (Burnet, 1976; Bunning, 1990).

**COMMENTS:** The deposit was developed by three adits that total several hundred feet and a large open pit mine. A 300-ton/day flotation mill was built at the site (Hunting, 1956).

## REFERENCES

- Barksdale, J. D., 1948, Stratigraphy in the Methow quadrangle, Washington: Northwest Science, v. 22, no. 4, p. 164-176.
- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Burnet, F. W., 1976, Felsic volcanic rocks and mineral deposits in the Buck Mountain Formation andesites, Okanogan County, Washington: University of Washington Master of Science thesis, 26 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- McFarland, C. R.; McLucas, G. B.; Rigby, J. G.; Stoffel, K. L., 1979, Directory of Washington mining operations, 1979: Washington Division of Geology and Earth Resources Information Circular 69, 100 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.

**American Flag (376)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Wauconda	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bodie Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 45' 13.28" N	118° 50' 43.10" W	NE1/4 sec. 36, 38N, 31E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Klondike Mountain Formation	trachyte	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopyrite	quartz, fluorite, sanidine	
Cu	bornite		
Zn	sphalerite		
Ag	galena		
	gold		
	pyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene	

**PRODUCTION:** Two 30-ton shipments of copper ore made in 1918-1919 (Moen, 1980, p. 69).

**TECTONIC SETTING:** East-west extension during the Eocene resulted in formation of the Toroda Creek graben and other structures in which a thick section of volcanic and sedimentary rocks was deposited and preserved (Holder and others, 1989).

**ORE CONTROLS:** The main mineralized structure is a 2-8-ft-wide silicified shear zone that strikes N45E and dips 50NW. Brecciated and sheared volcanic rock is cemented by quartz and sanidine and as much as 5% fluorite. The silicified and fluoritized breccia body exposed on the surface above the adit is as much as 250 ft in diameter. The American Flag is the southernmost property of the Zalla M-Silver Bell mineralized belt that is about 1 mi long and several hundred feet wide (Moen, 1980, p. 69).

**GEOLOGIC SETTING:** Volcanic rocks in this part of the Toroda Creek graben plot as rhyolite on a TAS diagram, but they contain sparse plagioclase and mafic phenocrysts in an aphanitic groundmass that contains abundant plagioclase microlites. Plagioclase phenocrysts are commonly strongly sericitized, and mafic minerals are generally replaced by some combination of chlorite, calcite, epidote, and sphene; shapes of altered mafic minerals suggest that they were originally pyroxene. The presence of silica stringers and veinlets in some of these rocks suggests that silica was introduced during alteration (Stoffel, 1990, p. 11).

**COMMENTS:** The main adit is 97 ft long and trends N45E along a mineralized shear zone. Several small prospect pits were present. Dump samples collected by Moen (1980, p. 69) contained a trace to 12.5 oz/ton Ag.

**REFERENCES**

- Dorisy, C. E., 1937, Index of mineral occurrences in the state of Washington (revised edition): Washington State Planning Council Research Publication 3, 47 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.

**American Flag** (446)

ALTERNATE NAMES		DISTRICT	COUNTY
Oriental and Central			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mazama	1:62,500	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 35' 35.36" N	120° 23' 9.08" W	SE1/4 sec. 30, 36N, 20E	
LOCATION: on a cliff about 1 mi northeast of Mazama			
HOST ROCK: NAME	LITHOLOGY	AGE	
Fawn Peak stock	diorite to quartz diorite	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au Zn Pb	chalcopyrite arsenopyrite sphalerite	quartz, calcite, pyrite, tourmaline, sericite	
-			
DEPOSIT TYPE	MINERALIZATION AGE		
breccia pipe			

**PRODUCTION:** Produced a few hundred tons before 1910; had a small amount of production in 1940 (Hunting, 1956, p. 135).

**TECTONIC SETTING:** Magmatic arc rocks of the area include the Fawn Peak stock, which is probably the intrusive equivalent of the volcanic member of the Midnight Peak Formation. The stock was probably the intrusive rock core of an island arc (Riedell, 1979). The unit has yielded K-Ar magmatic biotite ages of approximately 88 m.y. (Stoffel, 1990, p. 28).

**ORE CONTROLS:** The deposit is in a breccia pipe. Strongly brecciated and partially rotated fragments of quartz diorite as much as 3 cm long are in a matrix of quartz, tourmaline, calcite, pyrite, and chalcopyrite. The American Flag deposit is part of the Mazama porphyry copper-molybdenum system (Riedell, 1979).

**GEOLOGIC SETTING:** The American Flag deposit is in a breccia pipe of the Fawn Peak stock of Cretaceous age. It is an elongate, northwest-trending intrusion along the northeast side of the Methow River valley; it is situated along the axis of the Goat Peak syncline (Stoffel and McGroder, 1990).

**COMMENTS:** The American Flag deposit is part of the Mazama porphyry copper-molybdenum deposit (Riedell, 1979).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Riedell, K. B., 1979, Geology and porphyry copper mineralization of the Fawn Peak intrusive complex, Methow Valley, Washington: University of Washington Master of Science thesis, 52 p., 4 pl.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**American Rand** (407)

ALTERNATE NAMES		DISTRICT	COUNTY
Spokane			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 53' 40.39" N	119° 32' 13.61" W	W1/4 corner sec. 11, 39N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	quartzite, argillite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Whisky Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mo	molybdenite	quartz	
Au	chalcopyrite		
Ag	gold		
Cu	galena		
	pyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced in 1916-1918 and 1935-1938 (Hunting, 1956, p. 149).

TECTONIC SETTING: Late Paleozoic marine sedimentary rocks were deposited along an active continental margin.

ORE CONTROLS: A 15-in.-wide, north-trending, 20-40E-dipping, quartz vein in argillite a short distance above the contact with a pluton (Umpleby, 1911, p. 98).

GEOLOGIC SETTING: The vein is in metasedimentary rocks of the Spectacle Formation approximately 75 ft above the contact with the Whisky Mountain pluton (Umpleby, 1911, p. 98; Rinehart and Fox, 1972, geol. map).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Andy O'Neil** (473)

ALTERNATE NAMES		DISTRICT	COUNTY
Andy O.		Nespelem	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Armstrong Creek	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 9' 53.75" N	119° 1' 37.09" W	N¼ sec.27, T 31N., R 30E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
porphyritic granodiorite of Manila Creek unnamed quartzite		granite, granodiorite quartzite, limestone	(Paleocene ?) - Eocene Paleozoic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Zn Cu	galena chalcopyrite tetrahedrite bornite	pyrite, marcasite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene - (Paleocene?)	

**PRODUCTION:** A test shipment was made prior to 1940 (Hunting, 1956, p. 300).

**TECTONIC SETTING:** The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). Keller Butte suite rocks were emplaced contemporaneously with ductile deformation associated with the formation of the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** The deposit consists of fissure veins and breccia in granodiorite at and near its contact with quartzite and limestone. The vein that contains the ore-rich horizon is in a cross shear that intersects the main vein, which strikes N68W and dips 40NE. Sulfide minerals are crudely zoned in the vein; pyrite is more abundant at the walls, and galena, sphalerite, and chalcopyrite are more abundant at the center of the vein. The hanging wall has a gradational contact with the host rock, whereas the contact at the foot wall is sharp (Broch, 1979, p. 139).

**GEOLOGIC SETTING:** The deposit is in the (Paleocene?)-Eocene- porphyritic granodiorite of Manila Creek (Gulick and Korosec, 1990, geol. map).

**COMMENTS:** The main level crosscut intersects the main vein at approximately 150 ft. Two drifts were driven along the vein (Broch, 1979).

**REFERENCES**

- Broch, M. J., 1979, Igneous and metamorphic petrology, structure, and mineral deposits of the Mineral Ridge area (Moses mining district), Colville Indian Reservation, Washington: Washington State University Master of Science thesis, 204 p., 1 pl.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Antimony Bell** (441)

ALTERNATE NAMES		DISTRICT	COUNTY
Antimony Belle			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Hungry Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 9' 20.97" N	120° 8' 43.58" W	NE1/4SE1/4 sec. 25, 31N, 21E	
LOCATION: on a hill southeast of the South Fork Gold Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Newby Group		greenstone	Jurassic - Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb	stibnite stibiconite	calcite, breccia, antigorite	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

**PRODUCTION:** In 1940, 1,300 lb of stibnite ore were shipped (Hunting, 1956, p. 18).

**TECTONIC SETTING:** The Newby Group was deposited in an active margin setting, along or near an island arc.

**ORE CONTROLS:** Stibnite occurs along a shear zone in greenstone. The shear zone is 3-10 in. wide, strikes S66E, dips nearly vertical, and has been exposed at intervals over a 200-ft length (Hunting, 1956, p. 18).

**GEOLOGIC SETTING:** The mineralized shear zone is in the Jurassic-Cretaceous Newby Group (Bunning, 1990, geol. map).

**COMMENTS:** The deposit is developed by a 32-ft shaft and several adits (Hunting, 1956).

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

**Antimony Queen (442)**

ALTERNATE NAMES		DISTRICT	COUNTY
New Deal Dixie Queen Reedy Silver Seal		Gold Creek area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Hungry Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 12' 4.30" N	120° 10' 47.20" W	SW1/4NW1/4 sec. 11, 31N, 21E	
LOCATION: elev. 2,000 ft, 10 mi northwest of Methow, 4.5 mi up Copper Creek from the Methow highway			
HOST ROCK: NAME	LITHOLOGY	AGE	
Newby Group	argillite, graywacke	Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb Au Pb Zn W Ag	stibnite jamesonite arsenopyrite chalcopyrite pyrite galena sphalerite scheelite	calcite, pyrite, and pyrrhotite	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

**PRODUCTION:** A small amount of ore was extracted and may have been shipped in 1906. About 10 years later, 1,000 tons of stibnite ore and several hundred tons of antimony oxide, which had been manufactured in a small plant near the mine, were shipped. About 2/3 of a carload of ore was shipped in 1941 (Purdy, 1951, p. 109).

**TECTONIC SETTING:** The Newby Group was deposited in an active margin setting, along or near an island arc.

**ORE CONTROLS:** The host rocks of the deposit strike N30-40W and dip 50-80W. The dip of shear fractures containing fine stringer veins is at a lower angle than that of the bedding. Alteration, while extensive, is limited to a small area around the vein (Purdy, 1951).

**GEOLOGIC SETTING:** Mineralization is in veins in the graywacke and argillite of the Jurassic-Cretaceous Newby Group (Bunning, 1990).

**COMMENTS:** The mine was developed by three main adits and about 1,000 ft of workings (Hunting, 1956 p. 18).

**REFERENCES**

- Barksdale, J. D., 1948, Stratigraphy in the Methow quadrangle, Washington: Northwest Science, v. 22, no. 4, p. 164-176.  
 Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.  
 Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.  
 Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

**Apache (474)**

ALTERNATE NAMES		DISTRICT Nespelem	COUNTY Okanogan
PRIMARY QUADRANGLE Armstrong Creek	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 9' 28.47" N	LONGITUDE 119° 1' 11.66" W	SECTION, TOWNSHIP, AND RANGE center E1/2 sec. 27, 31N, 30E	
LOCATION:			
HOST ROCK: NAME porphyritic granodiorite of Manila Creek		LITHOLOGY granite, granodiorite	AGE (Paleocene?) -Eocene
COMMODITIES Ag Au Pb Mn Zn Cu	ORE MINERALS stephanite argentite silver tetrahedrite galena stephanite pyrargyrite chalcopyrite sphalerite chalcopyrite	NON-ORE MINERALS quartz, rhodocrosite	
DEPOSIT TYPE shear zone		MINERALIZATION AGE (Paleocene?) - Eocene	

**PRODUCTION:** Estimated production from 1911 to 1940 was about \$250,000 in Ag, and \$20,000 in Au. Shipments to the Tacoma smelter in 1911 contained about \$350/ton Au and Ag. Small hand-sorted, high-grade shipments contained as much as 2,444 oz/ton Ag. The bulk of the ore averaged 10-20 oz/ton Ag and was concentrated at the Great Metals mill near Nespelem Falls prior to shipment to the smelter (Moen, 1976, p. 129).

**TECTONIC SETTING:** This epizonal pluton is part of the Keller Butte suite. Intrusion of the granodiorite was contemporaneous with deformation associated with the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** The average strike of the vein is N45W, and the dip is northeast. Mineralization is a stockwork of veins that are 0.2-1.2 in. wide; the ore zone at the Apache is 1.25 ft wide. The veins pinch and swell and have sharp contacts with the host rock. Economic mineralization occurs as lenticular or tabular lenses and pods distributed along single veins. Ore shoots are complexly brecciated and contain angular fragments of broken wall rock and vein quartz. Open-space filling is exhibited by such features as euhedral quartz crystals. A wide alteration halo surrounds the veins at the surface (Broch, 1979).

**GEOLOGIC SETTING:** The veins are in a shear zone in the (Paleocene?) - Eocene porphyritic granodiorite of Manila Creek.

**COMMENTS:** The vein was accessed by several adits (Broch, 1979).

**REFERENCES**

- Broch, M. J., 1979, Igneous and metamorphic petrology, structure, and mineral deposits of the Mineral Ridge area (Moses mining district), Colville Indian Reservation, Washington: Washington State University Master of Science thesis, 204 p., 1 pl.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.



**Arlington** (420)

ALTERNATE NAMES		DISTRICT Conconully Ruby Hill area	COUNTY Okanogan
PRIMARY QUADRANGLE Ruby Hill	SCALE 1:24,000	1/2° x 1° QUAD Omak	1° x 2° QUAD Okanogan
LATITUDE 48° 28' 19.65" N	LONGITUDE 119° 44' 7.02" W	SECTION, TOWNSHIP, AND RANGE NE1/4SE1/4 sec. 6, 34N, 25E	
LOCATION: elev. 4,080 ft			
HOST ROCK: NAME Conconully pluton metamorphic complex of Conconully	LITHOLOGY granodiorite gneiss	AGE Cretaceous pre-Jurassic	
COMMODITIES Ag Cu Pb Au Zn	ORE MINERALS galena tetrahedrite pyrargyrite chalcopyrite sphalerite	NON-ORE MINERALS quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE Cretaceous?	

**PRODUCTION:** The Arlington vein was one of the first deposits to be discovered on Ruby Hill. One thousand tons of silver ore, which had a net value of \$25,000, were mined by the summer of 1893 and concentrated at a mill in Ruby City. Between 1914 and 1921, several thousand tons of ore which contained 66.6 oz/ton Ag were mined at a net profit of \$31,000. A 50-ton/day flotation mill was erected in 1937. Between 1938 and 1939, 5,700 tons of ore that had a net value of \$71,683 were concentrated in the mill. Mining and milling operations ceased early in 1940 (Moen, 1973, p. 12).

**TECTONIC SETTING:** The Conconully pluton is a directionless, post-tectonic plutonic rock that was intruded into a major structural zone (Stoffel, K. L., DGER, 1990, oral commun.).

**ORE CONTROLS:** The Arlington vein is 1-6 ft thick, strikes north, and dips 70W. The vein for the most part occurs along the contact between the Conconully pluton and gneiss. Transverse faults offset the vein as much as 10 ft, whereas faults that parallel the vein have drawn the ore minerals out into thin dark-gray bands (Moen, 1973, p. 12).

**GEOLOGIC SETTING:** The Arlington vein occurs in the Conconully pluton of Cretaceous age near the contact with gneiss of the metamorphic complex of Conconully (Moen, 1973; Rinehart, 1981; Stoffel, 1990).

**COMMENTS:** Underground workings consisted of shafts, drifts, and crosscuts that have a total length of 4,500 ft. The workings exposed the vein on four levels for a total vertical distance of 540 ft (Moen, 1973, p. 12).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Rinehart, C. D., 1981, Reconnaissance geochemical survey of gully and stream sediments, and geologic summary, in part of the Okanogan Range, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 74, 24 p., 3 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Bales (443)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Carlton-Gold Creek area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Twisp East	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 16' 41.15" N	120° 6' 25.60" W	W1/2NE1/4 sec. 17, 32N, 22E	
LOCATION: elev. 3,000 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Twisp Valley Schist	amphibolite, greenschist	pre-Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb	stibnite	quartz	
As	stibiconite		
Ni	As mineral?		
	Ni mineral?		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			
shear zone			

PRODUCTION: Mining in 1951-52 yielded an estimated 100 tons of ore (Hunting, 1956, p. 18).

TECTONIC SETTING: The metasedimentary rocks were originally deposited in an active margin environment; the interleaving of metaperidotite with supracrustal rocks suggests that significant tectonic mixing predated the complicated folding and metamorphic history of the Twisp Valley Schist (Bunning, 1990).

ORE CONTROLS: The mineralized zone consists of three elements: intensely hydrothermally altered wall rock, quartz, and lenticular masses of stibnite. The altered rock that encloses the veinlets and irregular masses of stibnite is composed of erratically distributed, alternating iron-oxide stained and bleached rock that is composed of quartz, sericite, and minor amounts of kaolin. Crisscrossing the altered rock are veinlets and microveinlets of stibiconite. The contact between the wall rock and the lenticular masses of stibnite is characterized by the presence of quartz, especially drusy quartz in the vugs. The vein or mineralized zone strikes about N47E and dips 80SE. The zone containing stibnite ranges from several inches to 5 ft in width (Purdy, 1951, p. 122-123).

GEOLOGIC SETTING: The mineralized system is within hornblende-quartz diorite of the Twisp Valley Schist (Purdy, 1951; Bunning, 1990, geol. map).

COMMENTS: Only surface work has been done at this property (Purdy, 1951).

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Barksdale, J. D., 1948, Stratigraphy in the Methow quadrangle, Washington: Northwest Science, v. 22, no. 4, p. 164-176.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.

**Bellevue (377)**

ALTERNATE NAMES		DISTRICT	COUNTY
Bellview		Wannacut Lake	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 54' 49.07" N	119° 34' 2.51" W	N1/2 sec. 4, 39N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	slate	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
metagabbro, metadiorite		Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu	chalcopyrite pyrargyrite stephanite silver gold gold-silver telluride	quartz, arsenopyrite, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** A test shipment of 1,000 lb gave returns of \$75 in Au and Ag; gold represented a little over half the total return (Umpleby, 1911).

**TECTONIC SETTING:** Late Paleozoic sediments were deposited along an active continental margin.

**ORE CONTROLS:** The N29E-trending, 45-66SE-dipping quartz vein is from 10 in. to 3 ft wide. The upper part of the vein is in slate, and limestone is present in the lower part (Hunting, 1956, p. 136; Rinehart and Fox, 1972, p. 78).

**GEOLOGIC SETTING:** The vein is in the slate, argillite, and limestone of the Spectacle Formation of the Anarchist Group of Permian age. The Spectacle Formation is intruded by gabbro and/or diorite of Permian-Triassic age.

**COMMENTS:** A small shaft was dug and 250-ft of development work was done on the property (Rinehart and Fox, 1972)

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Bi-Metallic (472)**

ALTERNATE NAMES		DISTRICT Myers Creek Buckhorn Mtn area	COUNTY Okanogan
PRIMARY QUADRANGLE Havillah	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 50' 59.02" N	LONGITUDE 119° 8' 32.60" W	SECTION, TOWNSHIP, AND RANGE NW1/4 sec. 26 SE1/4 sec. 27, 39N, 29E	
LOCATION: 3 mi northeast of Havillah			
HOST ROCK: NAME alkalic complex at Bimetallic Mountain	LITHOLOGY alaskite, aplite, monzonite	AGE Cretaceous or Tertiary	
COMMODITIES Mo Cu Ag Au W	ORE MINERALS molybdenite chalcopyrite powellite scheelite	NON-ORE MINERALS pyrite; sericitic and kaolinitic alteration	
DEPOSIT TYPE shear zone porphyry ?		MINERALIZATION AGE	

**PRODUCTION:** High-grade copper ore was shipped from one of the shafts in 1918 (Moen, 1980, p.57).

**TECTONIC SETTING:** Deposit is in an alkalic intrusive complex.

**ORE CONTROLS:** Mineralization is contained in several undulating shear zones in hydrothermally altered alaskite, aplite, and monzonite (Purdy, 1954, p. 34-44). Shear zones do not appear to continue for more than 30 ft. Sericite, kaolinite, and disseminated ore minerals are present in the shear zones. Ore minerals are localized in small ore shoots.

**GEOLOGIC SETTING:** The deposit is within the alkalic complex of Bimetallic Mountain, which is of Cretaceous or Tertiary age. The monzonite, which is brecciated and locally includes shonkinite, is intruded by hydrothermally altered, fine-grained, alaskite dikes. The rocks intrude metasedimentary rocks of the Spectacle Formation of the Permian Anarchist Group, which are locally hornfelsed (Stoffel, 1990).

**COMMENTS:** Development consists of several shafts and two adits. A total of 2,200 ft of linear trenching was done by the U.S. Bureau of Mines (USBM) in 1945 to determine the extent of mineralization; no significant mineralization was found. Assays of samples from the adit by the USBM ranged from 2.20 to 3.87% Mo and 0.03 to 0.32% Cu and showed traces of Au and Ag. Samples taken from the trench averaged 0.32% Cu; trace amounts of Au and Ag were found in some samples. Sampling suggests that the better copper values are found in samples from near the surface, whereas the higher molybdenum values are from samples collected at deeper levels in the mineralized system (Moen, 1980, p. 57).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Purdy, C. P., Jr., 1954, Molybdenum occurrences of Washington: Washington Division of Mines and Geology Report of Investigations 18, 118 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Black Bear (378)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Palmer Mtn Alice Property area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Loomis	1:24,000	Oroville	Okanogan
LATTITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 50' 16.30" N	119° 37' 37.85" W	NE1/4 sec. 36, T39N, R25E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	greenstone	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granitic rocks, felsic to intermediate		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
Ag	chalcopyrite (?)		
Cu			
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

**PRODUCTION:** Produced \$150,000 prior to 1902; produced 77 tons in 1947 (Hunting, 1956, p. 81).

**TECTONIC SETTING:** The greenstones were deposited along an active continental margin proximal to an island arc.

**ORE CONTROLS:** The 4-ft wide, N65W-trending and 75NE-dipping quartz vein is in banded greenstone (Rinehart and Fox, 1972, p. 80).

**GEOLOGIC SETTING:** Veins are in banded and schistose greenstone of the Permian-Triassic Palmer Mountain Greenstone; greenstone was intruded by felsic to intermediate rocks of probable Jurassic-Cretaceous age (Rinehart and Fox, 1972, geol. map).

**COMMENTS:** Developed by 2,500 ft of workings and a 300-ft shaft (Rinehart and Fox, 1972).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Blue Lake** (447)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Blue Goat Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 41' 14.46" N	119° 42' 20.29" W	SW1/4 sec. 21, 37N, 25E	
LOCATION: on the east side of a northward-trending ridge from Goat Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Blue Goat pluton	granodiorite	(Triassic?) - Jurassic-	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au	chalcopyrite gold	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: About 5,000 tons of ore were taken out in 1901 (Hunting, 1956, p. 63).

TECTONIC SETTING: The Blue Goat pluton is a elongate pluton that is roughly concordant with the regional foliation in the local metamorphic rocks, but contacts are locally sharp and discordant. The country rock was probably deformed and metamorphosed prior to the emplacement of the pluton (Rinehart and Fox, 1976, p. 32).

ORE CONTROLS: Mineralization consists of several small quartz stringers in granodiorite; these contain a little chalcopyrite (Hunting, 1956, p. 63).

GEOLOGIC SETTING: Veins at the Blue Lake property are in the directionless, sub-porphyritic granodiorite of the (Triassic?)-Jurassic Blue Goat pluton (Stoffel, 1990, p. 33).

COMMENTS: The property was developed by a 300-ft adit (Hunting, 1956, p. 63).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Jones, E. L., Jr., 1917, Reconnaissance of the Conconully and Ruby mining districts, Washington. *In* Contributions to economic geology (short papers and preliminary reports), 1916-Part I, metals and nonmetals except fuels: U.S. Geological Survey Bulletin 640, p. 11-36.
- Landes, Henry; Thyng, W. S.; Lyon, D A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.
- Rinehart, C. D; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Bodie (379)**

ALTERNATE NAMES		DISTRICT	COUNTY
Northern Gold		Wauconda	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bodie	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 49' 50.05" N	118° 53' 25.99" W	SW1/4 sec. 3, 38N, 31E, and sec. 34, 39N, 31E	
LOCATION: elev. 3,000 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Klondike Mountain Formation		andesite, dacite	Eocene
Klondike Mountain Formation		tuff	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Fe	gold chalcopryrite pyrite pyrrhotite magnetite	pyrite, pyrrhotite, magnetite, quartz, chalcedonic quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein, epithermal breccia		Eocene	

**PRODUCTION:** A 10-stamp amalgamation-cyanide mill processed approximately 15,000 tons of gold ore from 1902 to 1916. In 1930 a 70-ton/day mill was built to recover free gold and low-grade ore that contained about 0.65 oz/ton Au and 0.60 oz/ton Ag. In 1938, the capacity of the mill was increased to 125 tons/day, and cyanidation and flotation circuits were added. Mining and milling were carried out until 1944, with as many as 40 miners working three shifts and producing as much as \$1,500 of Au and Ag a day. From 1930 to 1944, about 50,000 tons of ore valued at about \$280,000 was mined.

**TECTONIC SETTING:** These rocks were deposited in the Toroda Creek graben, a volcano-tectonic depression that was the result of east-west extension in the Eocene (Holder and others, 1989).

**ORE CONTROLS:** The Bodie vein is 2-22 ft wide and consists of many, parallel, closely spaced quartz veins. The individual veins are several inches to 5 ft in width and strike north to N16E and dip 60-80W. Vein quartz is white to light gray, fine grained, and commonly chalcedonic and colloform. Prospects pits have exposed the vein for about 6,000 ft; underground workings indicate the vein extends in excess of 500 ft along its dip. Gold is rarely visible; sulfides and magnetite are fine grained and disseminated in the quartz. The black streaks that parallel the walls contain dustlike particles of sulfides and gold and are the richest part of the vein. The gold grade in protore diminishes with depth from about 0.15 to 0.002 oz/ton; ore from the surface to the 100 level averaged about 1 oz/ton Au. The silver to gold ratio is 1:1. In addition to the veins, gold occurs in breccia in volcanic rocks that adjoin the Bodie vein on the west. The breccia strikes north and dips 35-45W. In the upper workings of the mine the breccia forms the hanging wall of the vein. The breccia extends below the 700 level to an undetermined depth. Clasts of the breccia are highly to moderately altered volcanic rocks and are cemented by cryptocrystalline silica and gouge (Moen, 1980, p. 73).

**GEOLOGIC SETTING:** The deposit is in the Klondike Mountain Formation. Whole-rock geochemistry suggests that the rock is a rhyolite; however, thin section work by Stoffel (1990) suggests that altered mafic minerals were originally pyroxenes and that the high silica content of the rock may be due to the addition of silica in the form of stringers and veinlets.

**COMMENTS:** Most of the mining took place on the Bodie No. 1 claim. At least 10,000 ft of underground work on five main crosscut adits penetrate the breccia and intersect the vein. A glory hole mined rich parts of the Bodie vein at the surface (Moen, 1980, p. 74).

**REFERENCES**

- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.
- Holder, R. W.; Gaylord, D R.; Holder, G. A. M., 1989, Plutonism, volcanism, and sedimentation associated with core complex and graben development in the central Okanogan Highlands, Washington. IN Joseph, N. L.; and others, editors, 1989, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 187-200.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

## Okanogan

- McFarland, C. R.; McLucas, G. B.; Rigby, J. G.; Stoffel, K. L., 1979, Directory of Washington mining operations, 1979: Washington Division of Geology and Earth Resources Information Circular 69, 100 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.



**Bullfrog (380)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Palmer Mtn	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 54' 58.42" N	119° 34' 29.35" W	S 1/2 SW 1/4 sec. 33, 40N, 26E	
LOCATION: elev. 3,560 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Bullfrog Mountain Formation	schistose quartzite, sericite schist	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
felsic dike		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	black metallic sulfide	quartz	
Ag	pyrite		
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Shipped 4,600 lb of ore (Hunting, 1956, p. 137).

TECTONIC SETTING: Late Paleozoic sediments were deposited along a convergent continental margin.

ORE CONTROLS: Reportedly, a 7-ft-wide quartz vein in quartzite and sericitic schist was traceable for 3,000 ft (Hunting, 1956, p. 137).

GEOLOGIC SETTING: The vein is in sericitic schist and schistose quartzite of the Permian Bullfrog Mountain Formation (Rinehart and Fox, 1972).

COMMENTS: There is an adit and two shafts at property.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Bunker Hill** (381)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Loomis	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 50' 39.25" N	119° 38' 3.74" W	approximately sec. 25, 39N, 25E	
LOCATION: on the west slope of Palmer Mountain at elev. 3,500 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	greenstone, metagabbro	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Produced 5 tons of high-grade gold ore prior to 1890 (Bethune, 1891, p. 64).

TECTONIC SETTING: The greenstones were deposited along a convergent continental margin proximal to an island arc.

ORE CONTROLS: A 20-in.-wide (on the surface) vertical vein trends northeast and is exposed on the surface for more than 300 ft (Hunting, 1956, p. 137).

GEOLOGIC SETTING: The vein is in greenstone and metagabbro of the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972, geol. map).

COMMENTS: The property was developed by two adits (Bethune, 1991).

## REFERENCES

- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Butcher Boy (382)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Myers Creek Chesaw area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Chesaw	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 12.36" N	119° 2' 59.61" W	N1/2NW1/4 sec. 21, 40N, 30E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Kobau Formation?		greenstone, metadiorite	Permian or Triassic (?)
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	pyrrhotite	quartz, pyrite, pyrrhotite	
Ag	sphalerite		
Zn	galena		
Pb	gold		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Eleven carloads of gold ore were shipped in 1907 (Hunting, 1956, p. 137). One carload was shipped in 1908; it averaged 4 oz/ton Au. Production ended in 1909 (Moen, 1973).

TECTONIC SETTING: Greenstone was deposited along a convergent continental margin proximal to an island arc.

ORE CONTROLS: A quartz fissure vein in greenstone that varies in width from 1 in. to as much as 6 ft. The vein trends N50W and dips 74NE. Gold is rarely visible, except in high-grade ore shoots (Moen, 1980, p. 33). Gold is associated with pyrite-rich parts of the vein, which contain as much as 5 oz/ton Au (Umpleby, 1911, p. 50).

GEOLOGIC SETTING: The vein is in greenstone and metadiorite of the Permian or Triassic (?) Kobau Formation (Moen, 1980).

COMMENTS: Development work consists of a 326-ft adit, a shaft 150 ft west of and 50 ft higher in elevation than the portal that contained high-grade-gold ore that was mined in the early 1900s. The shaft does not connect with the adit (Moen, 1980, p. 33).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Caribou** (383)

ALTERNATE NAMES		DISTRICT	COUNTY
		Myers Creek Buckhorn Mtn Area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Buckhorn Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 52.93" N	118° 59' 26.54" W	SW1/4SW1/4 sec. 13, T40N, R30E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	quartzite, argillite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Buckhorn Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	pyrite	pyrrhotite, skarn, and tactite minerals	
Ag	magnetite		
Cu	chalcopyrite		
Fe	malachite		
DEPOSIT TYPE		MINERALIZATION AGE	
contact metasomatic			

**PRODUCTION:** Between 1908 and 1916, the mine was operated by British Columbia Copper Co., and eight carloads of gold-silver-copper ore were shipped to smelters in British Columbia (Moen, 1980, p. 43).

**TECTONIC SETTING:** The Spectacle Formation was deposited along an active continental margin.

**ORE CONTROLS:** Contact metamorphic deposit at the contact of metasedimentary rocks and biotite-hornblende granodiorite (Moen, 1980, p. 43).

**GEOLOGIC SETTING:** The contact metamorphic deposit is found at the contact of Permian metasedimentary rocks and the Jurassic-Cretaceous granodiorite of the Buckhorn Mountain pluton (Moen, 1980, p. 43; Stoffel, 1990, p. 29).

**COMMENTS:** This deposit appears to be a northwesterly extension of the ore bodies at the Magnetic mine. Shafts, adits, prospect pits, and trenches are on the property. A core hole drilled by Granby Consolidated Mining, Smelting & Power Co. contained intercepts of 0.2-4.3% Cu within 112 ft of the surface.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

***Cassimer Bar placer*** (487)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bridgeport	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 5' 56.99" N	119° 43' 4.83" W	sec. 17, 30N, 25E	
LOCATION: at the mouth of the Okanogan River			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Considerable production was reported for 1860 to 1890 (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Castle Creek** (466)

ALTERNATE NAMES		DISTRICT	COUNTY
		Park City	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bald Knob	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 22' 55.98" N	118° 51' 22.41" W	secs. 1, 11, and 12, 33N, 31E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed argillite and shale unit	argillite, shale	Paleozoic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granite of Moses Mountain		Paleocene - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	pyrite, marcasite, quartz	
Ag	sphalerite		
Au	chalcopyrite		
Cu	tetrahedrite		
Zn			
DEPOSIT TYPE		MINERALIZATION AGE	
veins			

**PRODUCTION:** Produced an unknown amount of ore in 1906, 1918, and 1920 (Hunting, 1956, p. 217).

**TECTONIC SETTING:** The granite of Moses Mountain is part of the Keller Butte suite of Holder and Holder (1988). Rocks of the Keller Butte suite were emplaced during regional ductile stretching associated with deformation in the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** Mineralization consists of lenticular quartz masses containing scattered patches of ore in a series of metamorphic rocks intruded by granitic dikes (Hunting, 1956, p. 217).

**GEOLOGIC SETTING:** At the Castle Creek mine, Paleozoic black shale containing hematitic, brown siltstone interbeds is intruded by dikes of the Paleocene-Eocene granite of Moses Mountain (Joseph, 1990, geol. map).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

**Central (475)**

ALTERNATE NAMES		DISTRICT	COUNTY
Trinidad			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Aeneas Lake	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 37' 59.07" N	119° 33' 23.86" W	W1/2 sec. 10, 36N, 26E	
LOCATION: elev. 2,000 ft, 8.5 mi southwest of Tonasket; by the south shore of Turtle Lake			
HOST ROCK: NAME		LITHOLOGY	AGE
unnamed hypabyssal intrusive rocks		dacite	Eocene
unnamed crystal tuff		crystal-lithic and crystal tuff	Eocene
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	chalcopyrite	pyrite	
Pb	galena		
Au	sphalerite		
Cu	gold		
Zn			
DEPOSIT TYPE		MINERALIZATION AGE	
volcanic-plutonic		Eocene	

**PRODUCTION:** In 1918, 30 tons of ore that averaged \$12/ton Ag and Au were shipped from the 32-ft level of the shaft (Moen, 1973).

**TECTONIC SETTING:** Eocene volcano-tectonic depressions formed as the result of regional east-west extension (Holder and others, 1989).

**ORE CONTROLS:** According to Moen (1973), the deposit is in brecciated and silicified black argillite that has been intruded by an north-trending aplite porphyry dike, which appears to be several hundred feet wide. Parts of the dike contain pods of galena and pyrite that are accompanied by minor chalcopyrite and sphalerite (Moen, 1973, p. 37). Rinehart and Fox (1976) indicate the deposit is in a Tertiary crystal tuff intruded by a Tertiary dike.

**GEOLOGIC SETTING:** According to Rinehart and Fox (1976) and Stoffel (1990), the deposit is within crystal-lithic and crystal tuff of Eocene age intruded by a northwest-trending dacitic dike.

**COMMENTS:** A shaft has been sunk 50 ft on the dike, and drifts were driven at 32 and 50 ft below the collar of the shaft (Moen, 1973, p. 37).

**REFERENCES**

- Holder, R. W.; Gaylord, D R.; Holder, G. A. M., 1989, Plutonism, volcanism, and sedimentation associated with core complex and graben development in the central Okanogan Highlands, Washington. IN Joseph, N. L.; and others, editors, 1989, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 187-200.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Rinehart, C. D; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Chicago** (384)

ALTERNATE NAMES		DISTRICT Squaw Creek	COUNTY Okanogan
PRIMARY QUADRANGLE Cooper Mtn	SCALE 1:24,000	1/2° x 1° QUAD Twisp	1° x 2° QUAD Concrete
LATITUDE 48° 6' 41.51" N	LONGITUDE 120° 3' 43.65" W	SECTION, TOWNSHIP, AND RANGE sec. 10, T30N, R22E	
LOCATION:			
HOST ROCK: NAME Methow gneiss	LITHOLOGY tonalitic gneiss	AGE pre-Cretaceous	
COMMODITIES Au Cu Zn Fe	ORE MINERALS gold chalcopyrite?	NON-ORE MINERALS quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: 3 tons prior to 1897 (Hunting, 1956, p. 137).

TECTONIC SETTING: Intrusion of the Methow Gneiss (orthogneiss) during the Cretaceous probably took place at moderate crustal levels and was later metamorphosed to lower amphibolite facies (Bunning, 1990).

ORE CONTROLS: An 18-in. vein is traceable for 300 ft (Hunting, 1956, p. 137).

GEOLOGIC SETTING: The vein is in the Methow Gneiss (Bunning, 1990, geol. map).

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.



**Chloride Queen (385)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Nighthawk	Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 55' 13.89" N	119° 37' 36.02" W	NE¼SE¼ sec. 36, 40N, 25E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Kobau Formation		argillite, greenstone	Permian or Triassic (?)
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced in 1936 and 1937 (Hunting, 1956, p. 137).

TECTONIC SETTING: Greenstones were deposited along a convergent continental margin proximal to an island arc.

ORE CONTROLS: A 1-4-ft wide, iron-stained quartz vein in metasedimentary and metavolcanic rocks is sparsely mineralized with pyrite and free gold (Hunting, 1956, p. 137).

GEOLOGIC SETTING: In greenstone, argillite, and quartzite (metachert) of the Permian or Triassic Kobau Formation (Rinehart and Fox, 1972, geol. map).

COMMENTS: Developed by 50-ft inclined shaft (Hunting, 1956, p. 137), although Rinehart and Fox (1972) report no workings were present at this location and suggest that the deposit may be mislocated.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Copper Glance** (448)

ALTERNATE NAMES		DISTRICT	COUNTY
		Methow Valley Mining Eightmile Creek area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Billy Goat Mtn	1:24,000	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 45' 7.74" N	120° 18' 7.99" W	NE1/4 sec. 35, 38N, 20E	
LOCATION: elev. 5,600 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
andesite of Isabella Ridge	andesite	Cretaceous?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
plagioclase porphyry dikes			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite	pyrite, hematite, quartz, calcite, barite	
Au	chalcocite		
Ag	gold		
Fe	tetrahedrite		
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone		Cretaceous?	

PRODUCTION: Two tons of hand-sorted ore were shipped in 1914 (Huntting, 1956, p. 63).

TECTONIC SETTING: The andesite of Isabella Ridge was deposited in an island arc setting during the Cretaceous.

ORE CONTROLS: A fracture zone in andesite 80-100 ft wide contains locally distributed copper and a minor amounts of gold and silver (Huntting, 1956, p. 63). Mineralization occurs in many small shears in shattered andesitic pyroclastic rocks that are cut by several plagioclase porphyry dikes.

GEOLOGIC SETTING: The deposit is in andesite of Isabella Ridge, which is of probable Cretaceous age (Stoffel and McGroder, 1990).

COMMENTS: The property was developed by a 200-ft adit, a 50-ft adit, and several open cuts (Huntting, 1956, p. 63).

## REFERENCES

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Staatz, M. H.; Weis, P. L.; Tabor, R. W.; Robertson, J. F.; Van Noy, R. M.; Pattee, E. C.; Holt, D. C., 1971, Mineral resources of the Pasayten Wilderness Area, Washington: U.S. Geological Survey Bulletin 1325, 255 p., 3 pl.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**Copper World Extension (449)**

ALTERNATE NAMES		DISTRICT	COUNTY
Iron Mask Iron Master (?)		Palmer Mtn	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 51' 25.34" N	119° 35' 34.98" W	S 1/2 sec. 20 and N 1/2 sec. 29, 39N, 26E	
LOCATION: elev. 3,940 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	andesite	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
altered greenstone, metadiabase			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite	pyrite, pyrrhotite, arsenopyrite, magnetite, marcasite, and quartz; augite extensively replaced by secondary epidote, calcite, quartz, and chlorite	
Au	azurite		
Ag	malachite		
W	gold		
Zn	sphalerite		
Fe	bornite		
	pyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
massive sulfide			

**PRODUCTION:** Produced an unknown amount of ore prior to 1911. Produced 3,486 tons of ore in 1918-1919, which averaged 3.147% Cu, 0.42 oz/ton Ag, and 0.03 oz/ton Au (Hunting, 1956, p. 64).

**TECTONIC SETTING:** Rocks of the Palmer Mountain Greenstone were probably deposited proximal to island arcs.

**ORE CONTROLS:** The mineralized zone trends N85W, dips 40SW, and contains massive sulfides in en echelon pods in schistose andesite. Hanging wall andesite is highly silicified. A fault is probably present between the mineralization and the silicified andesite; talc and serpentine developed between the mineralized zones. The andesite is altered, and thin sections show that plagioclase and augite are extensively replaced by secondary epidote, calcite, quartz, and chlorite (Patty, 1921, p. 242-243; Rinehart and Fox, 1972, p. 84).

**GEOLOGIC SETTING:** Ore is in en echelon tabular lenses in altered andesite of the Permian-Triassic Palmer Mountain Greenstone (Patty, 1921, p. 241; Rinehart and Fox, 1972, p. 84).

**COMMENTS:** A 300-ft shaft was driven; crosscuts are on the 100-, 200-, and 300-ft levels (Rinehart and Fox, 1972).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- McIntyre, A. W., 1907, Copper deposits of Washington: American Mining Congress, 9th Annual Session, Report of Proceedings, p. 238-250.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Patty, E. N.; Glover, S. L., 1921, The mineral resources of Washington, with statistics for 1919: Washington Geological Survey Bulletin 21, 155 p., 1 pl.
- Rinehart, C. D; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Crounse placer** (488)

ALTERNATE NAMES		DISTRICT	COUNTY
Strawberry Creek placer			Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Strawberry Mountain	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 23' 59.18" N	118° 52' 43.94" W	S½ sec. 35, 34N, 31E	
LOCATION: on Strawberry Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold magnetite ilmenite	stream gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Reportedly \$100 worth of gold (Hunting, 1956, p. 187).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. Terraces from 20 to 200 ft wide along the stream are underlain by a shallow layer of coarse gravel (Hunting, 1956, p. 187).

COMMENTS: The deposit has been examined by small pits. Two pans of gravel from the layer next to the bedrock yielded 1 cent in Au and 1 oz of black sand (Hunting, 1956, p. 187).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Crystal Butte** (386)

ALTERNATE NAMES		DISTRICT	COUNTY
Crystal Butte Camp Mother Lode		Myers Creek Buckhorn Mtn area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Chesaw	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 55' 15.07" N	119° 0' 16.66" W	center W1/2 sec. 35, 40N, 30E	
LOCATION: elev. 4,800 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	limestone, argillite, quartzite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
acidic to intermediate intrusive rock		Mesozoic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Pb Zn Cu	galena chalcopyrite sphalerite	pyrite, arsenopyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic disseminated			

**PRODUCTION:** A mill was built in 1908, but there is no record that the mill produced concentrate. A small-scale mining operation was carried out from 1937 to 1941; this resulted in several carloads of ore being shipped to the smelter in Trail, British Columbia, and to the Bunker Hill smelter in Idaho. A few shipments ran as high as 1 oz/ton Au, 33 oz/ton Ag, and 28% Pb. The mine has been idle since 1941 (Moen, 1980, p. 44-45).

**TECTONIC SETTING:** Late Paleozoic rocks were deposited along an active continental margin.

**ORE CONTROLS:** Quartz fissure veins are in limestone, argillite, and quartzite and in quartz- and plagioclase-bearing porphyritic rocks. The N15-80E-striking vein dips 10-35NW and is 8-12 in. wide. Gouge zones are present along the hanging and foot walls of the vein; normal faults perpendicular to the vein have resulted in minor offsets. Sulfide minerals are sparsely disseminated in the vein or are present as thin, discontinuous bands that roughly parallel the walls of the vein (Moen, 1980, p. 45).

**GEOLOGIC SETTING:** Metasedimentary rocks of the Permian Spectacle Formation were intruded by quartz-plagioclase porphyritic intrusive rock of Mesozoic age (Moen, 1980, p. 45).

**COMMENTS:** Several adits were driven into the south-facing hillside at the property. The property is on patented mining claims (Moen, 1980, p. 44).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**First Thought** (421)

ALTERNATE NAMES		DISTRICT Conconully Ruby Hill area	COUNTY Okanogan
PRIMARY QUADRANGLE Ruby Hill	SCALE 1:24,000	1/2° x 1° QUAD Omak	1° x 2° QUAD Okanogan
LATITUDE 48° 29' 35.23" N	LONGITUDE 119° 44' 19.67" W	SECTION, TOWNSHIP, AND RANGE center NE¼ sec. 31, 35N, 25E	
LOCATION:			
HOST ROCK: NAME metamorphic complex of Conconully	LITHOLOGY gneiss, schist	AGE pre-Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION Conconully pluton		AGE Cretaceous	
COMMODITIES Ag Pb Cu Zn	ORE MINERALS galena tetrahedrite chalcopyrite chalcocite sphalerite stephanite	NON-ORE MINERALS pyrite, quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	

**PRODUCTION:** The mine produced silver ore valued at \$66,000 from 1892 to 1893. Production came to an end during the silver panic of 1893. Several small shipments of silver ore were made from the mine in the 1920s (Moen, 1973, p. 16).

**TECTONIC SETTING:** The Conconully pluton is a directionless, post-tectonic plutonic rock that intruded into a major structural zone (Stoffel, K. L., DGER, 1990, oral commun.).

**ORE CONTROLS:** Mineralization occurs in discontinuous lenticular lenses of quartz that are as much as 90 ft thick and as much as 700 ft in maximum width. The lenses strike N10E and dip 55 to 60E. They are present in highly foliated micaceous gneiss and feldspathic quartzite in the metamorphic complex of Conconully. The trend of the lenses of quartz is parallel to the foliation in the gneiss. Ore minerals are concentrated in ore shoots along the walls of the lenses; some of the ore shoots are as much as 5 ft thick. The ore shoots contain 75-100 oz/ton Ag, whereas the massive quartz between the shoots contains only 6-8 oz/ton Ag (Moen, 1973, p. 16-17).

**GEOLOGIC SETTING:** The mineralization at the First Thought occurs in the gneiss and quartzite of the metamorphic complex of Conconully near the contact of the Conconully pluton of Cretaceous (Rinehart, 1981; Stoffel, 1990).

**COMMENTS:** Underground workings of the First Thought mine consist of more than 4,000 ft of drifts on three levels that have a vertical extent of 350 ft (Moen, 1973, p. 17).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Rinehart, C. D., 1981, Reconnaissance geochemical survey of gully and stream sediments, and geologic summary, in part of the Okanogan Range, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 74, 24 p., 3 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Four Metals (467)**

ALTERNATE NAMES		DISTRICT	COUNTY
Mammoth (?)			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 54.31" N	119° 39' 51.78" W	secs. 22 and 23, 40N, 25E	
LOCATION: elev. 1,160 ft, on the south slope of Little Chopaka Mountain, on the Similkameen River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	quartzite (chert)	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Similkameen composite pluton		Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	quartz, garnet, epidote	
Ag	chalcopyrite		
Cu	bornite		
Zn	sphalerite		
W	scheelite		
Mo	molybdenite		
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

**PRODUCTION:** Produced 20 or more carloads of high-grade ore and concentrates between 1918 and 1921; 600 tons were milled in 1939 (Hunting, 1956, p.219).

**TECTONIC SETTING:** The sediments and volcanic materials of the Kobau Formation were deposited proximal to an island arc along a convergent continental margin.

**ORE CONTROLS:** A vein striking N7W and dipping 32-37W is in granodiorite at its contact with a roof pendant (Patty, 1921, p. 233; Rinehart and Fox, 1972).

**GEOLOGIC SETTING:** The deposit is a vein in the Similkameen composite pluton at or near the contact with quartzite (chert), limestone, and greenstone (hornblende schist) of the Kobau Formation of Permian or Triassic (?) age.

**COMMENTS:** Two shafts and three adits are present at the property (Rinehart and Fox, 1972).

**REFERENCES**

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Fourth Of July** (476)

ALTERNATE NAMES		DISTRICT Conconully Ruby Hill area	COUNTY Okanogan
PRIMARY QUADRANGLE Ruby Hill	SCALE 1:24,000	1/2° x 1° QUAD Omak	1° x 2° QUAD Okanogan
LATITUDE 48° 28' 36.64" N	LONGITUDE 119° 43' 39.10" W	SECTION, TOWNSHIP, AND RANGE SE1/4NW1/4 sec. 5, 34N, 25E	
LOCATION: elev. 4,500 ft			
HOST ROCK: NAME Salmon Creek schist and gneiss	LITHOLOGY biotite gneiss	AGE pre-Jurassic	
COMMODITIES Ag Au Pb Cu	ORE MINERALS tetrahedrite galena argentite silver stephanite cerargyrite ruby silver galena	NON-ORE MINERALS quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	

**PRODUCTION:** The vein was discovered in 1887. In 1889 a shipment of high-grade silver ore was sent to the smelter in Helena, Montana. A 200 ft shaft was sunk on the vein, and an average of 10 tons of high-grade ore per month was shipped to smelters. Silver ore was shipped to a concentrating mill in Ruby City until mid-1893 when the mine was forced to shut down because of the silver panic. Total production to this time amounted to \$36,000. From 1958 to 1964 hand-sorted shipments were sent to the smelter in Trail, British Columbia. Except for a small shipment in 1967, those were the last shipments from the mine (Moen, 1973).

**TECTONIC SETTING:** The Salmon Creek schist and gneiss is laterally equivalent to the metamorphic complex of Conconully (Gulick and Korosec, 1990), which is thought by Rinehart and Fox (1976) to grade into and include Late Triassic rocks. Late Triassic sedimentary rocks were deposited in an active continental margin setting associated with an island arc.

**ORE CONTROLS:** The vein, which averages 6 ft in thickness, strikes N10W and dips 70-80E. The vein is in biotite gneiss and is parallel to the contact with a granodiorite gneiss that is several hundred feet to the west. The ore minerals seem to be in a 2-ft-thick band that parallels the hanging wall. Parts of the vein are sheared; shears parallel the vein (Moen, 1973, p. 13-14).

**GEOLOGIC SETTING:** The vein is in biotite gneiss of the pre-Jurassic Salmon Creek schist and gneiss parallel to and west of the contact with granodiorite gneiss (Moen, 1973).

**COMMENTS:** The underground workings at the mine consist of a double-compartment 500-ft shaft and a single-compartment 200-ft shaft. Several levels have been driven and mined (Moen, 1973).

**REFERENCES**

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Jones, E. L., Jr., 1917, Reconnaissance of the Conconully and Ruby mining districts, Washington. In Contributions to economic geology (short papers and preliminary reports), 1916-Part I, metals and nonmetals except fuels: U.S. Geological Survey Bulletin 640, p. 11-36.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Schuster, J. E., 1973, Directory of Washington mining operations, 1971-72: Washington Division of Mines and Geology Information Circular 48, 97 p.



**Friday (453)**

ALTERNATE NAMES		DISTRICT	COUNTY
Tom Hal		Squaw Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Pateros	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 4' 51.79" N	119° 59' 8.22" W	SW1/4 sec. 20, 30N, 23E	
LOCATION: about 5 mi northwest of Pateros			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopyrite	quartz	
Ag	bornite		
Cu	malachite		
	pyrite		
	arsenopyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Production includes a \$5,000 shipment made prior to 1897 and one carload shipped in 1940. Ten tons of selected ore yielded \$70/ton prior to 1902 (Huntting, 1956, p. 139).

**TECTONIC SETTING:** The Methow Gneiss is a directionless, post-tectonic pluton that was emplaced at moderate to deep levels and then metamorphosed (Gulick and Korosec, 1990).

**ORE CONTROLS:** The quartz vein in the Cretaceous Methow Gneiss is about 1 ft wide (Huntting, 1956, p. 139).

**GEOLOGIC SETTING:** The vein is in tonalitic gneiss of the Methow Gneiss (Gulick and Korosec, 1990, geol. map).

**COMMENTS:** Mineralization at the Friday mine may be similar to that at the Sullivan prospect.

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p., 1 pl.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Fuller (465)**

ALTERNATE NAMES		DISTRICT	COUNTY
Lone Pine Claims		Squaw Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Pateros	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 4' 14.81" N	119° 55' 13.59" W	near center sec. 26, 30N, 23E	
LOCATION: elev. 2,300 ft, about 1 mi northwest of Pateros			
HOST ROCK: NAME		LITHOLOGY	AGE
Alta Lake migmatite		migmatite	Jurassic - Cretaceous (?)
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Cu Ti	magnetite chalcopyrite	hornblende, actinolite, biotite, calcite, epidote	
DEPOSIT TYPE		MINERALIZATION AGE	
disseminated			

**PRODUCTION:** Reportedly, 1,000 tons of ore was shipped prior to 1943; 1,250 tons were shipped in 1943 (Hunting, 1956, p. 198).

**TECTONIC SETTING:** Migmatitic parts of the gneiss of Alta Lake; migmatization appears to have been a polyphase event that involved both anatexis and igneous injection (Gulick and Korosec, 1990).

**ORE CONTROLS:** Magnetite is intimately intermixed with large amounts of hornblende and actinolite. The ore is coarse grained and has a schistose appearance. Ore occurs in seams and thin lenses. The deposit varies from a few feet to 15 ft thick and is exposed along strike for 400-500 ft. The mineralized zone grades into the gneiss and migmatite (Shedd and others, 1922).

**GEOLOGIC SETTING:** The Fuller deposit is in the migmatitic part of the amphibolite, schist, and gneiss of the Alta Lake migmatite. The migmatite is gradational with the layered metamorphic rocks and is characterized by irregular, lenticular, swirled, and contorted granitic, tonalitic, aplitic, and pegmatitic leucosomes (Gulick and Korosec, 1990, p. 40).

**COMMENTS:** Development consists of 150 ft of adits and a quarry (Hunting, 1990).

**REFERENCES**

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

**Gold Axe (387)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Myers Creek Buckhorn Mtn area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Buckhorn Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 3.99" N	118° 58' 48.94" W	SE1/4SW1/4 sec. 24, 40N, 30E	
LOCATION: elev. 5,280 ft, about 500 ft east of the summit of Buckhorn Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	limestone, greenstone	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Buckhorn Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu Ag Fe	pyrite pyrrhotite magnetite	pyrite, pyrrhotite, magnetite, garnet, epidote	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone contact metamorphic			

**PRODUCTION:** Sporadic production of copper and gold from 1910 to 1934; total production probably did not exceed 5,000 tons (Moen, 1980, p. 47). Handy (1916, p. 8) reports that six carloads of ore containing \$10-\$20 per ton in Au and minor amounts of silver were shipped. Hunting (1956, p. 139) reports 16 or 17 carloads of ore were shipped in 1914-15.

**TECTONIC SETTING:** The Kobau Formation was deposited along an active continental margin proximal to an island arc.

**ORE CONTROLS:** The shear zone trends N34W and dips from 65SW to vertical; magnetite and pyrrhotite occur in lens-like bodies with garnet and epidote along the shear zone and range from several inches to 1 ft in width. The contact metamorphic zone is not as well developed here as at the Magnetic mine (Moen, 1980, p. 47).

**GEOLOGIC SETTING:** The Jurassic-Cretaceous Buckhorn Mountain pluton intruded and contact metamorphosed limestones and greenstones of the Permian or Triassic (?) Kobau Formation.

**COMMENTS:** Development of the property consists of an adit and many prospect pits. Drilled by Crown Resources Corp. in 1989 and by Battle Mountain Gold in 1990. The property is part of a joint venture between Crown Resources and Battle Mountain Gold and is known as the Buckhorn Mountain deposit.

**REFERENCES**

- Handy, F. M., 1916, An investigation of the mineral deposits of northern Okanogan County: State College of Washington Department of Geology Bulletin 100, 27 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Gold Crown (388)**

ALTERNATE NAMES		DISTRICT	COUNTY
Golden Crown		Palmer Mountain	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 50' 4.94" N	119° 36' 38.11" W	SE1/4 sec. 31, 39N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	meta-andesite, metagabbro	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

**PRODUCTION:** Amount not known. Assay from a dump sample was \$105/ton Au (Bethune, 1891, p. 63).

**TECTONIC SETTING:** The Palmer Mountain Greenstone was deposited along a convergent continental margin proximal to an island arc.

**ORE CONTROLS:** The 10-ft-wide vein strikes northwest and dips 50E (Bethune, 1991, p. 63).

**GEOLOGIC SETTING:** Quartz vein in the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972, geol. map).

**COMMENTS:** Ore from this mine was handled at a 5-stamp mill at the War Eagle-Black Bear mine (Bethune, 1991).

**REFERENCES**

- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Gold Key** (454)

ALTERNATE NAMES		DISTRICT	COUNTY
		Mazama	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mazama	1:62,500	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 35' 58.82" N	120° 24' 4.21" W	near northwest corner sec. 30, 36N, 20 E	
LOCATION: about 1,000 ft northwest of the upper workings of the Mazama Pride mine			
HOST ROCK: NAME	LITHOLOGY	AGE	
Midnight Peak Formation	andesite	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Fawn Peak stock		Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu	chalcopyrite	quartz, calcite, pyrite, arsenopyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
stringer zone stockwork		Cretaceous	

PRODUCTION: In 1931, 37 tons of ore were shipped (Hunting, 1956, p. 140).

TECTONIC SETTING: The Midnight Peak Formation and the Fawn Peak stock were part of a Cretaceous island arc (Riedell, 1979).

ORE CONTROLS: The ore consists of quartz stringer veins that are mineralized to some extent (Hunting, 1956, p. 140). The deposit is in the actinolite-chlorite zone of the Mazama molybdenum porphyry system. Veins peripheral to the porphyry system generally trend N10-40E and N70-85W (Riedell, 1979).

GEOLOGIC SETTING: The Gold Key mine is in andesites of the Midnight Peak Formation, which are intruded by the Cretaceous Fawn Peak stock. The stock is probably the intrusive equivalent of the volcanic rocks (Stoffel and McGroder, 1990, p. 15).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Riedell, K. B., 1979, Geology and porphyry copper mineralization of the Fawn Peak intrusive complex, Methow Valley, Washington: University of Washington Master of Science thesis, 52 p., 4 pl.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**Golden Chariot (370)**

ALTERNATE NAMES		DISTRICT Oroville	COUNTY Okanogan
PRIMARY QUADRANGLE Oroville	SCALE 1:24,000	½° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 59' 23.78" N	LONGITUDE 119° 28' 52.44" W	SECTION, TOWNSHIP, AND RANGE SE¼ sec. 6, and NE¼ sec. 7, 40N, 27E	
LOCATION:			
HOST ROCK: NAME Silver Nail Lake pluton Kobau Formation	LITHOLOGY quartz diorite argillite, quartzite	AGE Jurassic - Cretaceous (?) Permian or Triassic (?)	
COMMODITIES Cu Au Ag W Mo	ORE MINERALS chalcopyrite scheelite molybdenite	NON-ORE MINERALS pyrite, quartz	
DEPOSIT TYPE porphyry vein		MINERALIZATION AGE	

**PRODUCTION:** Produced \$4,000 worth of ore prior to 1911; nine carloads of hand-sorted ore shipped sometime later (Hunting, 1956, p. 65).

**TECTONIC SETTING:** The Kobau Formation was deposited along an active continental margin proximal to an island arc.

**ORE CONTROLS:** The 5-ft-wide ore zone strikes north and dips 35W; it consists of a series of S-shaped mineralized quartz lenses. The quartz lenses range from a fraction of an inch to a foot in thickness (Culver and Broughton, 1945, p. 32-33). The deposit contains stockwork with quartz-K-feldspar flooding and sericitic alteration which is part of the Kelsey copper-molybdenum porphyry copper system (Roper, 1973).

**GEOLOGIC SETTING:** The deposit is in argillite of the Permian or Triassic (?) Kobau Formation and quartz diorite and fine-grained diorite of the concentrically zoned Jurassic-Cretaceous (?) Silver Nail Lake pluton (Stoffel, 1990, p. 28).

**COMMENTS:** The property was opened up by an inclined shaft down the 30 degree dip of the ore zone for a reported distance of 350 ft; several open cuts were also present (Culver and Broughton, 1945, p. 32).

**REFERENCES**

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Roper, M. W., 1973, Geology of the Kelsey copper-molybdenum property, Okanogan County, Washington: Montana State University Master of Science thesis, 97 p., 5 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Golden Zone** (389)

ALTERNATE NAMES		DISTRICT	COUNTY
		Nighthawk Mt Chopaka area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 58' 33.30" N	119° 43' 52.23" W	near SE corner sec. 7, 40N, 25E	
LOCATION: at the base of Mount Chopaka, elev. 1,600 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Similkameen composite pluton	granodiorite	Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, arsenopyrite, quartz	
Ag	galena		
Pb	chalcopyrite		
Cu	malachite		
Zn	azurite		
Mo	sphalerite		
	molybdenite?		
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

PRODUCTION: Produced prior 1911 and in 1939 (Hunting, 1956, p. 140).

TECTONIC SETTING: The Similkameen batholith is a concentrically zoned plutonic suite with early mafic alkalic rocks bordering and intruded by calc-alkalic granitic rocks. The undeformed body is in the upper plate of the Okanogan Valley fault (Buddington and Burmester, 1990).

ORE CONTROLS: The vein trends N20E and dips 40NW; rocks are visibly altered as much as 1 ft from the vein (Rinehart and Fox, 1972, p. 88).

GEOLOGIC SETTING: Shear zone in Similkameen composite pluton (Rinehart and Fox, 1972, geol. map).

COMMENTS: Developed by 5,000 ft of tunnels and drifts (Rinehart and Fox, 1972, p. 89).

## REFERENCES

- Buddington, A. M.; Burmester, R. F., 1990, The Similkameen batholith: A mid-Jurassic, post-tectonic complex in the Quesnel terrane, north-central Washington and south-central British Columbia [abstract]: Geological Society of America Abstracts with Programs, v. 22, n. 3, p. 10-11.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Gray Eagle** (390)

ALTERNATE NAMES		DISTRICT	COUNTY
Eagle		Myers Creek Chesaw area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Chesaw	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 47.34" N	119° 2' 36.68" W	SW1/4 sec. 16, 40N, 30E	
LOCATION: elev. 3,040 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	greenstone, siltite	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
hornblende-biotite granitic rocks			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	gold chalcopryrite arsenopyrite azurite sphalerite unknown Mo mineral galena	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein shear zone			

**PRODUCTION:** Small-scale mining in late 1900s. Minor production in 1916, 1921-24, 1937-39, and 1950-51; value probably less than \$50,000. Most mining was by pick and shovel. A small amalgamation mill was built in 1950-51; only 8 tons of ore containing a total of 4 oz Au and 4 oz Ag were mined and milled (Moen, 1980, p. 35).

**TECTONIC SETTING:** The Kobau Formation was deposited along an active continental margin proximal to an island arc.

**ORE CONTROLS:** North-striking, west-dipping, quartz and calcite veins in fractures 0.5-6 in. wide in sheared and brecciated rock. Visible gold is present in veins and stringers. The adit is in hydrothermally altered granitic rocks, greenstone, and siltite (Moen, 1980, p. 35).

**GEOLOGIC SETTING:** The vein is in siltite and greenstone of the Kobau Formation; altered granitic rocks occur with the mineralization.

**COMMENTS:** Mining on the Gray Eagle claim appears to have been carried out on two widely separated veins. One of the veins has also been mined as part of the Reco vein in the No. 2 adit of the Reco mine.

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.



*Gray Eagle no. 2* (391)

ALTERNATE NAMES		DISTRICT Squaw Creek Chesaw area	COUNTY Okanogan
PRIMARY QUADRANGLE Cooper Mtn	SCALE 1:24,000	1/2° x 1° QUAD Twisp	1° x 2° QUAD Concrete
LATITUDE 48° 5' 2.41" N	LONGITUDE 120° 1' 28.87" W	SECTION, TOWNSHIP, AND RANGE NW1/4SE1/4 sec. 24, 30N, 22E	
LOCATION:			
HOST ROCK: NAME Methow Gneiss	LITHOLOGY tonalitic gneiss	AGE Cretaceous	
COMMODITIES Au Cu	ORE MINERALS pyrite	NON-ORE MINERALS pyrite, quartz	
DEPOSIT TYPE _ shear zone		MINERALIZATION AGE	

PRODUCTION: Amount of production is unknown (Hunting, 1956, p. 140).

TECTONIC SETTING: The Methow Gneiss is a Cretaceous tonalitic gneiss that has been metamorphosed to the amphibolite facies (Bunning, 1990).

ORE CONTROLS: A lens of quartz, which is 15 ft long and 3 ft wide at its thickest portion, occurs in a shear zone in Cretaceous tonalite gneiss (Hunting, 1956, p. 140; Bunning, 1990, geol. map).

GEOLOGIC SETTING: The vein is in tonalitic gneiss of the Methow Gneiss of Cretaceous age (Bunning, 1990, geol. map).

COMMENTS: The deposit was developed by a short adit and open cuts (Hunting, 1956).

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Hiawatha** (392)

ALTERNATE NAMES		DISTRICT	COUNTY
Josie		Wannacut Lake	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 54' 1.39" N	119° 32' 44.98" W	NE¼ sec. 10, 39N, 26E	
LOCATION: elev. 3,200 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	argillite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Whisky Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	galena	quartz	
Ag	chalcopryite		
Pb	sphalerite		
Cu	gold		
Zn	pyrite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced some ore in 1938 (Hunting, 1956, p. 141).

**TECTONIC SETTING:** Marine sedimentary rocks of the Spectacle Formation were deposited along an active continental margin.

**ORE CONTROLS:** A friable white quartz vein is 1 to 12 ft wide, averages 3-4 ft wide, and crops out for about 2,500 ft. The vein dips 10W at the surface and 45W at depth. Faults offset the vein (Rinehart and Fox, 1972, p. 88).

**GEOLOGIC SETTING:** The vein is in argillite of the Spectacle Formation near the contact with monzonite and quartz diorite of the Whisky Mountain pluton (Hunting, 1956, p. 141; Rinehart and Fox, 1972, geol. map).

**COMMENTS:** Two 80-ft adits 80 ft apart are connected by a drift near the face (Rinehart and Fox, 1972, p. 89).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Hidden Treasure (455)**

ALTERNATE NAMES		DISTRICT	COUNTY
Sunshine Triangle		Squaw Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooper Mtn	1:24,000	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 24.24" N	120° 2' 26.96 "W	SE1/4 sec. 11, 30N, 22E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu	chalcopyrite galena sphalerite malachite	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced 90 tons of ore prior to 1902, and that returned \$67 per ton. Produced again in 1939-1942 (Hunting, 1956, p. 141).

**TECTONIC SETTING:** Intrusion of the Methow Gneiss protolith took place at moderate crustal levels; the intrusive was later metamorphosed to lower amphibolite grade with other metamorphic sequences in the area (Bunning, 1990).

**ORE CONTROLS:** Mineralization is in a 2-4-ft-wide vein in the Methow Gneiss (Hunting, 1956, p. 141).

**GEOLOGIC SETTING:** The vein is in tonalitic gneiss of the Methow Gneiss (Bunning, 1990, geol. map).

**COMMENTS:** The deposit is developed by two adits (Hunting, 1956, p. 141).

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.

**Highland (393)**

ALTERNATE NAMES		DISTRICT	COUNTY
Highland Light		Squaw Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooper Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 36.38" N	120° 2' 6.23" W	E1/2 sec. 11, and W1/2 sec. 12, 30N, 22E	
LOCATION: 30N, 22E			
HOST ROCK: NAME		LITHOLOGY	AGE
Methow Gneiss		tonalitic gneiss	Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, quartz	
Cu	chalcopyrite		
W	scheelite		
Zn	sphalerite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene?	

**PRODUCTION:** Production totaled \$13,000 during 1938 to 1941 in Au and Zn (Hunting, 1956, p. 141).

**TECTONIC SETTING:** The Methow Gneiss is a tonalitic gneiss of Cretaceous age that has been metamorphosed to amphibolite-facies grade (Bunning, 1990).

**ORE CONTROLS:** Several gold-bearing veins are present at the property. The Sailor Boy vein averages 6 ft thick and strikes S70E and dips 70NE. The quartz vein is exposed over a distance of 3,000 ft and is in the Methow Gneiss (Culver and Broughton, 1945, p. 47; Bunning, 1990, geol. map).

**GEOLOGIC SETTING:** The veins cut the Cretaceous Methow Gneiss (Bunning, 1990, geol. map).

**COMMENTS:** The property is developed by a 40-ft shaft and a 15-ft shaft 3,000 ft apart on the vein. An crosscut adit lies north of the shafts and 200 ft below them (Culver and Broughton, 1945). The property is bounded by the Holden-Campbell property on the west.

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.

**Holden-Campbell** (394)

ALTERNATE NAMES		DISTRICT	COUNTY
Hunter			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooper Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 59.89" N	120° 3' 24.64" W	mainly in secs. 10 and 11, 30N, 22E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
basic dike		Eocene?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, quartz	
Cu	chalcopyrite		
W	molybdenite		
Ag	scheelite		
Mo			
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene?	

**PRODUCTION:** The Holden-Campbell has produced a considerable amount of gold ore, and the adjacent Hunter mine (now part of the Holden-Campbell property) produced a small amount in 1940 (Hunting, 1956, p. 141).

**TECTONIC SETTING:** The Methow Gneiss is a tonalitic gneiss that has been metamorphosed to amphibolite grade (Bunning, 1990).

**ORE CONTROLS:** Several veins are present at the property. The mineralized rock consists of northeast-striking, steeply dipping, quartz veins that cut the gneiss. Several dikes are parallel to basic dikes that cut the gneiss (Culver and Broughton, 1945, p. 46).

**GEOLOGIC SETTING:** The veins are in tonalitic gneiss of the Cretaceous Methow Gneiss that has been cut by dikes of probable Eocene age (Bunning, 1990, geol. map; Culver and Broughton, 1945, p. 46).

**COMMENTS:** More than 1,100 ft of underground workings and several surface cuts developed the several veins on the property (Hunting, 1956, p. 141).

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p., 1 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Homestake** (414)

ALTERNATE NAMES		DISTRICT Conconully Conconully-Salmon Creek area	COUNTY Okanogan
PRIMARY QUADRANGLE Conconully East	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 34' 19.08" N	LONGITUDE 119° 44' 42.89" W	SECTION, TOWNSHIP, AND RANGE E1/2 SW1/4 sec. 31, 36N, 25E	
LOCATION: E1/2 SW1/4 sec. 31, 36N, 25E, elev. 3,200 ft			
HOST ROCK: NAME metamorphic complex of Conconully	LITHOLOGY quartz-mica schist	AGE pre-Jurassic	
COMMODITIES Pb Ag Au Bi	ORE MINERALS galena arsenopyrite unknown Bi mineral	NON-ORE MINERALS quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	
PRODUCTION: Produced 400 tons of ore prior to 1902, of which 100 tons netted \$1,200 (Moen, 1973, p. 28).			
TECTONIC SETTING: Rocks of the metamorphic complex of Conconully are thought by Rinehart and Fox (1976) grade upward into and include, in part, Late Triassic rocks. These rocks are of amphibolite grade where they are near plutonic rocks. These rocks formed in an active margin setting associated with island arcs.			
ORE CONTROLS: The 11-ft-thick, N22W-trending quartz vein dips 30SW (Moen, 1973, p. 28).			
GEOLOGIC SETTING: Quartz-mica schists of the metamorphic complex of Conconully contain garnet and sillimanite near the contact with plutonic rocks. The unit is intruded by irregular masses of alaskite and pegmatite (Rinehart and Fox, 1976, p.16)			
COMMENTS: This property was developed by a 29-ft shaft and a 75-ft adit (Moen, 1973)			
REFERENCES			
Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.			
Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.			
Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.			
Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.			

**Horn Silver (422)**

ALTERNATE NAMES		DISTRICT	COUNTY
Arizona		Nighthawk Bullfrog Mtn area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 45.44" N	119° 34' 5.27" W	S 1/2 sec. 21, 40N, 26E	
LOCATION: elev. 3,200 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	greenstone, argillite	Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granodiorite		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu	stephanite cerargyrite prousite galena chalcopyrite sphalerite pyrite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Several carloads were shipped prior to and during 1909 (Hunting, 1956, p.304); the values were mostly in silver and the ore averaged \$62/ton in 1909 (Rinehart and Fox, 1972, p. 90).

**TECTONIC SETTING:** Rocks of the Palmer Mountain Greenstone were deposited along an active continental margin proximal to an island arc.

**ORE CONTROLS:** Three veins are present on the property. The most developed vein strikes N53E and dips 40NW, and is from 4 in. to 4 ft wide, averaging 18 in. (Rinehart and Fox, 1972, p. 90).

**GEOLOGIC SETTING:** The quartz vein is in metavolcanic rocks of the Palmer Mountain Greenstone (Rinehart and Fox, 1972, geol. map).

**COMMENTS:** Development consists of a 75-ft adit and a 100-ft shaft (Moen, 1976, p. 122).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Independence** (456)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Methow	1:24,000	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 9' 24.23" N	120° 7' 12.39" W	SW1/4 sec. 29, 31N, 22E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu Mo	chalcopyrite molybdenite	pyrite, quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

**PRODUCTION:** Produced an unknown amount of ore in 1940 and 1942 (Hunting, 1956, p. 142).

**TECTONIC SETTING:** The Methow Gneiss protolith was probably intruded at moderate crustal levels; the intrusive was later metamorphosed to the amphibolite grade together with other metamorphic rocks in the area (Bunning, 1990).

**ORE CONTROLS:** Mineralization is in a 3-ft-wide quartz vein in gneissic diorite (Hunting, 1956, p. 142).

**GEOLOGIC SETTING:** The deposit consists of a vein in the Methow Gneiss (Bunning, 1990, geol. map).

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.



**Ivanhoe (423)**

ALTERNATE NAMES		DISTRICT	COUNTY
Ivanhoe Group		Palmer Mtn	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 52' 29.63" N	119° 34' 26.40" W	SW 1/4 sec. 16, 39N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Bullfrog Mountain Formation of the Anarchist Group	argillite, siltite, phyllite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
dacitic dikes		Tertiary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Au Sb Pb	stephanite cerargyrite tetrahedrite pyrite galena gold silver	quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Hand-sorted, near-surface ores mined between 1888 and 1897 contained as much as 1,000 oz/ton Ag and several ounces of Au. Approximately 1,000 tons of ore was shipped to smelters in Washington and Montana; several shipments averaged 392 oz/ton Ag and 1.2 oz/ton Au (Moen, 1976, p. 123; Rinehart and Fox, 1972, p. 91).

**TECTONIC SETTING:** Sediments of the Anarchist Group were deposited along an active continental margin.

**ORE CONTROLS:** Seven quartz veins were mined. The Ivanhoe vein is 3.5-4.5 ft thick and strikes north-northwest and dips 60W. Silver ore was richer near the surface; veins intersected at a depth of 1,560 ft had average Ag content of less than 1 oz/ton (Umpleby, 1911).

**GEOLOGIC SETTING:** The vein is in phyllite and argillite of the Permian Bullfrog Mountain Formation of the Anarchist Group.

**COMMENTS:** The Ivanhoe vein was developed by several shallow shafts, one of which was later sunk to a depth of 500 ft. The richest ores were near the surface; these were mined by stripping the hanging wall of the vein over an area 70 by 120 ft. In about 1914, a 4,400-ft crosscut was driven from the west slope of Palmer Mountain in an attempt to intersect the vein at a depth of 1,560 ft (Umpleby, 1911).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**John Judge** (396)

ALTERNATE NAMES		DISTRICT	COUNTY
Leadville Denver City Grandview		Palmer Mtn	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 51' 35.94" N	119° 37' 11.74" W	SW1/4 sec. 19, 39N, 26E	
LOCATION: elev. 2,500 ft, about 2.7 mi north-northeast of Loomis, and on the west side of Palmer Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	meta-andesite; metagabbro	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu Pb Ag	gold chalcopyrite bornite galena pyrargyrite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced 15 tons of ore in 1937; also produced in 1938 and 1939 (Hunting, 1956, p. 143).

**TECTONIC SETTING:** The Palmer Mountain Greenstone was deposited along an active continental margin proximal to island arcs.

**ORE CONTROLS:** Vertical, 2-3-ft-wide vein trending N60E in chlorite schist (Rinehart and Fox, 1972, p. 90). The vein is traceable for 200 ft in the lower adit (Hunting, 1956, p. 143).

**GEOLOGIC SETTING:** Vein in chlorite schist of the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972).

**COMMENTS:** Development consisted of 2,500 ft of adits and shafts (Rinehart and Fox, 1972, p. 91)

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Johnson Creek (445)**

ALTERNATE NAMES		DISTRICT	COUNTY
Funkhauser Omak		Conconully Omak area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Omak	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 28' 55.10" N	119° 34' 57.35" W	NE1/4 sec. 5, 34N, 26E	
LOCATION: elev. 1,750 ft, 6 mi north of Omak by road; near the intersection of the Riverside cut-off and the Omak-Conconully highway			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed serpentinite unnamed calc-silicate rocks	serpentinite calc-silicate rocks	pre-Cretaceous pre-Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cr Ni Fe	chromite white sulfide of nickel	serpentinite, antigorite, magnesite, talc, garnet, diopside	
DEPOSIT TYPE		MINERALIZATION AGE	
disseminated (?) magmatic segregation (?)			

**PRODUCTION:** Produced 5.9 tons of ore in 1955; these assayed 42% Cr<sub>2</sub>O<sub>3</sub> and had a Cr/Fe ratio of 2.9 (Hunting, 1956, p. 38).

**TECTONIC SETTING:** An ultramafic mass in the North Cascades.

**ORE CONTROLS:** The deposit is near an ultrabasic-dolomite contact. Chromite is disseminated sparsely through the ultrabasic rock. Near the ultrabasic-dolomite contact, a tabular body of nearly solid chromite is present; it measures 7.5 ft by 6 ft by 3 ft. A stringer about 2 in. wide and 18 in. long was found about 10 ft north of the main pod (Hunting, 1956, p. 38).

**GEOLOGIC SETTING:** The deposit is in calc-silicate rock derived from altered ultrabasic rocks or from dolomitic sedimentary rocks. The chromite-bearing unit consists of calc-silicate minerals and magnesite, talc, and tremolite; serpentinitized magnesite talc schist; and garnet-diopside calc-silicate granofels, hornblende schist, and serpentinitized carbonate (Gulick and Korosec, 1990, p. 27).

**REFERENCES**

- Banta, H. E., 1956, Directory of Washington mining operations, 1956: Washington Division of Mines and Geology Information Circular 25, 87 p.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Livingston, V. E., Jr., 1957, Directory of Washington mining operations, 1957: Washington Division of Mines and Geology Information Circular 27, 94 p.
- Wilson, Hewitt; Skinner, K. G.; Hurst, T. L., 1943, Some refractory properties of Washington chromite: U.S. Bureau of Mines Report of Investigations 3694, 31 p.

**Kaaba** (415)

ALTERNATE NAMES		DISTRICT	COUNTY
Caaba Kaaba-Texas		Nighthawk	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 16.13" N	119° 39' 4.71" W	center NE1/4 sec. 23, 40N, 25E	
LOCATION: elev. 2,500 ft, about 9 mi north of Loomis, near the base of the southwest slope of Little Chopaka Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Similkameen composite pluton	granodiorite	Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
lamprophyre granodiorite		Eocene? Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb Ag Zn Cu Au W Mo	galena sphalerite chalcoppyrite scheelite molybdenite	pyrite, marcasite, quartz, calcite, gouge	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced in 1918 and 1943-1951; total is about 205,000 oz of Ag. From 1943 to 1946, the mine produced 135,973 oz Ag, 1,357,185 lb Pb, 506,050 lb Zn, and 99,410 lb Cu (Moen, 1976, p. 123).

**TECTONIC SETTING:** The Similkameen batholith is a concentrically zoned plutonic suite with early mafic alkalic rocks bordering and being intruded by calc-alkalic granitic rocks. The undeformed body is on the upper plate of the Okanogan Valley fault (Buddington and Burmester, 1990).

**ORE CONTROLS:** The vein at the Kaaba mine is a persistent, but low-grade, banded quartz vein that strikes N3W, dips 43-55W, and averages 10 ft in width. This vein appears to be bordered by two dikes; however, at the 200-ft level the dike in the footwall cuts the vein, and 50 ft below the collar, the vein is entirely in the Similkameen composite pluton (Patty, 1921, p. 230). Patty (1921) suggests that the dikes are younger than the mineralization because they are not altered and because they cut mineralized rock. The depth to the bottom of the deposit is 240 ft.

**GEOLOGIC SETTING:** The vein is in the granodiorite of the Similkameen composite pluton of Jurassic age.

**COMMENTS:** An inclined shaft is 300 ft long with 1,100 ft of drifts are on four levels; more than half the drifting is on the third level. A 150-ton/day flotation mill was constructed on property (Hunting, 1956, p. 220).

**REFERENCES**

- Buddington, A. M.; Burmester, R. F., 1990, The Similkameen batholith: A mid-Jurassic, post-tectonic complex in the Quesnel terrane, north-central Washington and south-central British Columbia [abstract]: Geological Society of America Abstracts with Programs, v. 22, n. 3, p. 10-11.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Kankakee** (450)

ALTERNATE NAMES		DISTRICT Nespelem	COUNTY Okanogan
PRIMARY QUADRANGLE Armstrong Creek	SCALE 1:24,000	1/2° x 1° QUAD Omak	1° x 2° QUAD Okanogan
LATITUDE 48° 10' 42.25" N	LONGITUDE 119° 2' 13.19" W	SECTION, TOWNSHIP, AND RANGE near NW corner, sec. 22, 31N, 30E	
LOCATION:			
HOST ROCK: NAME porphyritic granodiorite of Manila Creek unnamed quartzite	LITHOLOGY granite, granodiorite quartzite	AGE (Paleocene?) - Eocene Paleozoic	
COMMODITIES Cu Pb Ag	ORE MINERALS chalcopyrite argentite?	NON-ORE MINERALS pyrite, fluorite, quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE (Paleocene?) - Eocene	

PRODUCTION: A small test shipment of oxidized lead ore was made (Hunting, 1956, p. 66).

TECTONIC SETTING: The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). Intrusion of Keller Butte suite rocks into metasedimentary country rocks was contemporaneous with regional ductile stretching associated with the formation of the metamorphic core complexes (Holder and Holder, 1988).

ORE CONTROLS: Weak mineralization is present along fractures in granite and quartzite. One quartz vein is 2-in. wide (Hunting, 1956, p. 66).

GEOLOGIC SETTING: Quartzite of probable Paleozoic age is intruded by the (Paleocene?)-Eocene porphyritic granodiorite of Manila Creek (Gulick and Korosec, 1990).

## REFERENCES

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Kelsey (438)**

ALTERNATE NAMES		DISTRICT	COUNTY
Hart Stone Stone		Similkameen Oroville	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 59' 41.66" N	119° 28' 40.74" W	secs. 5, 6, 7, and 8, 40N, 27E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Silver Nail Pluton	diorite	Jurassic - Cretaceous (?)	
Silver Nail Pluton	quartz diorite	Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
quartzo-feldspathic dikes			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Mo Ag Au	chalcopyrite molybdenite malachite azurite chrysocolla	pyrite (dominant), magnetite, pyrrhotite, chlorite, quartz, calcite, epidote, sericite, secondary biotite (weak), K-spar, actinolite, garnet; sericitization, potassic alteration, tactite development	
DEPOSIT TYPE		MINERALIZATION AGE	
porphyry copper veins veinlets disseminated			
PRODUCTION: Has not produced.			
TECTONIC SETTING: To the north in British Columbia, distribution of major porphyry copper deposits is, in part controlled by the Quesnel trough. The Kelsey property lies on the southward projection of this structural trough and is similar to mined deposits to the north.			
ORE CONTROLS: The bulk of the mineralization is in the Silver Nail pluton, a coarsely crystalline quartz diorite. The intrusive is sheeted; contacts between the various phases are fairly flat. Sills and thrust faults are present. At the south end of the property, the pluton is brecciated. Permian Anarchist Group rocks (slate, phyllite, impure marble, greenstone, metaconglomerate, and metawacke) were incorporated as fragments in the breccia and as slivers in the pluton. The most concentrated copper-molybdenum mineralization is associated with quartz-sericite alteration and intense fracturing; country rock surrounding the mineralized zone is chloritized. Mineralization in greenstone and limy portions of the Anarchist Group consists of copper-bearing tactite containing pyrrhotite, minor garnet, and epidote. Quartz veins contain chalcopyrite-rich pods. Hunting (1956, p. 67) reports 18 assays averaging 2.62% Cu, 0.6 oz/ton Ag, and 0.04 oz/ton Au. During the 1960s, exploration programs for porphyry copper deposits outlined two zones in the pluton. The Central Zone is 4,800 ft long, has a northwest-trending axis, and averages 1,530 ft wide. It contains 66,575,200 tons of indicated ore averaging 0.286% Cu and an additional 137,560,000 tons of inferred ore of similar grade. The geological potential of the central zone to a depth of 600 ft is 280,000,000 tons. Molybdenum credits are variable and range from 0.015 to 0.02% MoS <sub>2</sub> . Hole K-3 drilled by Inland Copper Ltd. encountered 85 ft of ore averaging 0.407% Cu and 0.13% Mo. The Northwest Zone contains a total of 34,320,000 tons of indicated and inferred ore grading 0.241% Cu. The Northwest Zone extends across the international boundary into British Columbia.			
GEOLOGIC SETTING: The major rock type at the Kelsey deposit is the Jurassic-Cretaceous (?) Silver Nail quartz diorite pluton. Subordinate intrusive rock types are quartz monzonite and syenite. Late aplite occurs near the south contact. All intrude Permian Anarchist Group greenstone, phyllite, slate, and minor marble, limestone, and quartzite. Tertiary conglomerate and dacite adjoining the deposit are not altered, and therefore are post ore.			
COMMENTS: The Kelsey property, in 1974, consisted of 15 patented claims, 91 unpatented claims, and land held in fee. It was discovered in the late 1800s, and attempts were made to develop the higher grade quartz veins. From 1965 to 1974 American Smelting and Refining Co., Inland Copper Ltd., Canadian Superior Exploration Ltd., St. Joe Minerals Corp., Placer-Amex Inc., and U.S. Borax explored the property for porphyry copper deposits. They drilled a total of 36,085 ft of X-ray, rotary, percussion, and core holes. The property in nov (1990) owned by Wilbur G. Hallauer of Oroville.			

UNPUBLISHED INFORMATION: Grant, A.R., 1974, Report of evaluation, Kelsey property, Okanogan County, Washington. Grant, A.R., 1980, Summary report, Kelsey property, Okanogan County, Washington. These reports are in DGER files.

#### REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Roper, M. W., 1973, Geology of the Kelsey copper-molybdenum property, Okanogan County, Washington: Montana State University Master of Science thesis, 97 p., 5 pl.
- Umpleby, J. B., 1911, Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, pt. 2, p. 55-111.

# Okanogan

## Key (424)

ALTERNATE NAMES		DISTRICT Conconully Conconully area	COUNTY Okanogan
PRIMARY QUADRANGLE Conconully West	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 34' 39.73" N	LONGITUDE 119° 45' 8.35" W	SECTION, TOWNSHIP, AND RANGE SW1/4NW1/4 sec. 31, 36N, 25E	
LOCATION: elev. 2,800 ft			
HOST ROCK: NAME metamorphic complex of Conconully	LITHOLOGY schist, migmatite	AGE pre-Jurassic	
COMMODITIES Ag Pb Au Cu Zn Bi	ORE MINERALS galena pyrite chalcopyrite sphalerite unknown Bi mineral	NON-ORE MINERALS quartz	

DEPOSIT TYPE vein	MINERALIZATION AGE
<p>PRODUCTION: Produced a total of 1,500 tons; 12 tons were produced in 1914 and netted \$444 in Ag and Pb (Moen, 1973, p. 28).</p> <p>TECTONIC SETTING: Triassic (or pre-Jurassic) sediments were deposited along an active continental margin setting.</p> <p>ORE CONTROLS: The sparsely metallized quartz vein is 3-10 ft thick, strikes N25E, and dips 60NW.</p> <p>GEOLOGIC SETTING: The vein is in schist and migmatite of the metamorphic complex of Conconully. These rocks are thought by Rinehart and Fox (1976) to be in part Late Triassic because they are intruded by a small pluton that is correlated with the Loomis pluton, dated at 194 m.y.</p> <p>COMMENTS: Development at the property consists of a 300-ft adit, a 80-ft shaft, and a 105-ft drift (Moen, 1973, p. 28).</p>	

### REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.



**Kimberly** (416)

ALTERNATE NAMES		DISTRICT	COUNTY
		Wannacut Lake	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bullfrog Mtn	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 53' 24.32" N	119° 31' 57.40" W	SW1/4 sec. 11, 39N, 26E	
LOCATION: elev.: 1,920 ft, about 7 mi northeast of Loomis			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	argillite	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Whisky mountain pluton		Jurassic to Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	quartz	
Ag	chalcopyrite		
Au	sphalerite		
Cu			
Zn			
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Rinehart and Fox (1972, p. 93) report that the mine has produced "considerable hand-sorted ore of \$40 grade." Presumably this is ore worth \$40 per ton.

TECTONIC SETTING: Marine rocks of the Spectacle Formation were derived from sediments deposited along an active continental margin.

ORE CONTROLS: Mineralization is present in en echelon lenses of quartz as much as 50 ft in long and 1-5 ft wide, in the hanging wall over contact (possibly fault) that strikes N46W and dips 52SW between a porphyritic granodiorite and quartz monzonite pluton and argillite and slate (Umpleby, 1911, p. 97; Rinehart and Fox, 1972, geol. map).

GEOLOGIC SETTING: The deposit occurs at contact of the Permian Spectacle Formation and the Jurassic-Cretaceous Whisky Mountain pluton (Rinehart and Fox, 1972, geol. map).

COMMENTS: Development at the site included a 140-ft inclined shaft and drifts at the 60-, 80-, and 100-ft levels (Rinehart and Fox, 1972).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**King Solomon (435)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Nighthawk	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 57.93" N	119° 38' 10.73" W	NE1/4SW1/4 sec. 24, 40N, 25E	
LOCATION: elev. 3,040 ft, about 9.2 mi north of Loomis			
HOST ROCK: NAME		LITHOLOGY	AGE
Kobau Formation		metachert, slate	Permian or Triassic (?)
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	pyrite, banded vuggy quartz	
Cu	chalcopryrite		
Au			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced lead ore in 1924 (Hunting, 1956, p. 220).

TECTONIC SETTING: Marine sedimentary and volcanic materials of the Kobau Formation were deposited along an active continental margin proximal to an island arc.

ORE CONTROLS: The 7-ft-wide, banded, vuggy quartz vein trends N5E and dips 50W. The vein pinches at depth (Rinehart and Fox, 1972, p. 92).

GEOLOGIC SETTING: The vein is in thinly laminated slate and quartzite (metachert) of the Kobau Formation (Rinehart and Fox, 1972, geol. map).

COMMENTS: An inclined shaft developed property (Rinehart and Fox, 1972).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Lakeview (395)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Oroville	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 58' 50.21" N	119° 28' 59.72" W	center sec. 7, 40N, 27E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	greenstone, metachert	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Silver Nail Lake pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold		
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: Two carloads were shipped prior to 1916 (Hunting, 1956, p. 143).

TECTONIC SETTING: The Kobau Formation was deposited along an active continental margin proximal to island arcs.

ORE CONTROLS: Said to be a thick blanket-like body of ore parallel to the slope of the hill (Hunting, 1956, p. 143).

GEOLOGIC SETTING: Partly chloritized greenstone and metachert of the Triassic or Permian Kobau Formation (Fox, 1970, geol. map; Roper, 1973, alteration map).

COMMENTS: Hydrothermal alteration associated with the Lakeview mine occurs in the alteration halo of the Kelsey porphyry Cu-Mo system (Roper, 1973; alteration map)

## REFERENCES

- Fox, K., F., Jr., 1970, Geologic map of the Oroville quadrangle, Okanogan County, Washington: U. S. Geological Survey Open-File Report 70-128, 3 sheets, scale 1:62,500.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Handy, F. M., 1916, An investigation of the mineral deposits of northern Okanogan County: State College of Washington Department of Geology Bulletin 100, 27 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Roper, M. W., 1973, Geology of the Kelsey copper-molybdenum property, Okanogan County, Washington: Montana State University Master of Science thesis, 97 p., 5 pl.

**Last Chance** (477)

ALTERNATE NAMES		DISTRICT	COUNTY
Ruby Lode claim		Conconully Ruby Hill area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Ruby Hill	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 29' 30.19" N	119° 44' 23.92" W	E 1/2 sec. 31, 35N, 25E	
LOCATION: elev. 3,000 ft, at the north end of Ruby Hill			
HOST ROCK: NAME		LITHOLOGY	AGE
Conconully granodiorite metamorphic complex of Conconully		granodiorite hornblende-mica schist	Cretaceous pre-Jurassic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Cu Zn	tetrahedrite galena chalcopyrite sphalerite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Ore was blocked out but was not mined because of the silver panic of 1983. In 1920 the mine was reopened, and ore that averaged 30 oz/ton Ag, 17% Pb, and 4% Cu was shipped to a smelter at Bradley, Idaho. Small shipments of ore were made again in 1921 and 1924 (Moen, 1973, p. 18).

**TECTONIC SETTING:** The quartz monzonite was probably emplaced in a magmatic arc.

**ORE CONTROLS:** The quartz fissure vein that averages 12 ft in width strikes S50E and dips 70SW. Ore minerals are concentrated in ore shoots that are 2-4 ft thick and as much as 200 ft long. Ore minerals are sparsely disseminated between ore shoots (Moen, 1973, p. 18).

**GEOLOGIC SETTING:** The vein is in granodiorite of Cretaceous age near the contact with hornblende-biotite schist of the metamorphic complex of Conconully (Gulick and Korosec, 1990, geol. map).

**REFERENCES**

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Leuena (425)**

ALTERNATE NAMES		DISTRICT	COUNTY
Laeuna Launa		Conconully Mineral Hill area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Conconully West	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 34' 19.69" N	119° 47' 32.48"W	SW1/4 sec. 35, 36N, 24E	
LOCATION: elev. 5,000 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Conconully pluton		granodiorite, quartz monzonite	Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Au	tetrahedrite stephanite argentite chalcopyrite?	quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Several carloads were shipped prior to 1890 (Moen, 1973).

**TECTONIC SETTING:** The Conconully pluton is a directionless, post-tectonic pluton that was intruded into a major structural zone (Stoffel, K. L., DGER, 1990, oral commun.)

**ORE CONTROLS:** The 7-ft-thick quartz vein in granodiorite strikes N55E and is vertical (Moen, 1973).

**GEOLOGIC SETTING:** The vein is in the Conconully pluton, which is a leucocratic, equigranular to porphyritic, medium- to coarse-grained, biotite- and hornblende-bearing granodiorite to quartz monzonite of Cretaceous age (Stoffel, 1990).

**COMMENTS:** High-grade parts of the vein contained 200-800 oz/ton Ag.

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Lilman (478)**

ALTERNATE NAMES		DISTRICT Nespelem	COUNTY Okanogan
PRIMARY QUADRANGLE Armstrong Creek	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 9' 16.46" N	LONGITUDE 119° 1' 18.92" W	SECTION, TOWNSHIP, AND RANGE SE1/4SE1/4 sec. 27, 31N, 30E	
LOCATION:			
HOST ROCK: NAME porphyritic granodiorite of Manila Creek		LITHOLOGY granite, granodiorite	AGE (Paleocene?) - Eocene-
COMMODITIES Ag Au Cu	ORE MINERALS tetrahedrite chalcopyrite	NON-ORE MINERALS pyrite, pyrrhotite, quartz	
DEPOSIT TYPE vein veinlets		MINERALIZATION AGE (Paleocene?) - Eocene	

PRODUCTION: Produced 16 tons of ore prior to 1940 (Moen, 1976, p. 131).

TECTONIC SETTING: The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). Rocks of the Keller Butte suite were emplaced during regional ductile stretching associated with deformation in the metamorphic core complexes (Holder and Holder, 1988).

ORE CONTROLS: The deposit consists of lenses and veinlets along a 1.5-20-ft-wide shear zone in granodiorite.

GEOLOGIC SETTING: The vein is in the (Paleocene?)-Eocene porphyritic granodiorite of Manila Creek (Gulick and Korosec, 1990, geol. map).

COMMENTS: Development consists of 35- and 50-ft shafts and a 150-ft adit (Moen, 1976, p. 131).

## REFERENCES

- Broch, M. J., 1979, Igneous and metamorphic petrology, structure, and mineral deposits of the Mineral Ridge area (Moses mining district), Colville Indian Reservation, Washington: Washington State University Master of Science thesis, 204 p., 1 pl.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.

**Little Chief** (479)

ALTERNATE NAMES		DISTRICT	COUNTY
Double Header Grand Coulee Ruby Silver		Nespelem	Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Armstrong Creek	1:24,000	Omak	Okanogan
LATTITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 8' 55.66" N	119° 0' 31.70" W	NE¼NW¼ sec. 35. 31N, 30E	
LOCATION: about 2 mi southwest of Nespelem village			
HOST ROCK: NAME	LITHOLOGY	AGE	
porphyritic granodiorite of Manila Creek	granite, granodiorite	(Paleocene?) - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Au Cu	argentite stephanite pyrargyrite silver chalcopyrite	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		(Paleocene?) - Eocene	

**PRODUCTION:** The mine is the second largest producer in the Nespelem district. The deposit produced \$60,000 in silver from 1911 to 1921. A small amount of concentrate was produced from 1937 to 1954 (Moen, 1976, p. 132).

**TECTONIC SETTING:** The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). The Keller Butte suite was intruded during regional ductile stretching related to deformation in the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** Two veins produced ore. The main vein strikes N45W and dips 45-60NE. Fluid inclusions in calcite in the vein homogenized at between 235 and 245 degrees C (Broch, 1979).

**GEOLOGIC SETTING:** Two veins are present in the (Paleocene?) - Eocene porphyritic granodiorite of Manila Creek (Gulick and Korosec, 1990, geol. map).

**COMMENTS:** Development consists of a 560-ft adit with 640 ft of drifts and crosscuts as well as a 200-ft shaft with stopes (Moen, 1976, p. 131).

**REFERENCES**

- Broch, M. J., 1979, Igneous and metamorphic petrology, structure, and mineral deposits of the Mineral Ridge area (Moses mining district), Colville Indian Reservation, Washington: Washington State University Master of Science thesis, 204 p., 1 pl.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Washington Division of Mines and Geology, 1971, Directory of Washington mining operations, 1969-70: Washington Division of Mines and Geology Information Circular 46, 88 p.

**Lone Star** (468)

ALTERNATE NAMES		DISTRICT	COUNTY
Star		Conconully Conconully area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Conconully West	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 34' 29.92" N	119° 45' 20.28" W	E 1/2 sec. 36, 36N, 24E	
LOCATION: elev. 2,480 ft, about 1 mi north of Conconully and on the west side of Salmon Creek			
HOST ROCK: NAME		LITHOLOGY	AGE
Conconully pluton		granodiorite and quartz monzonite	Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	quartz, banded quartz, pyrite	
Ag	tetrahedrite		
Au	stephanite		
Cu	chalcopyrite		
Zn	sphalerite		
W			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** By 1890, ore that contained as much as 200 oz/ton Ag and 30% Pb was mined from several levels. By 1897, about \$40,000 had been spent developing the mine, but production amounted to only several thousand dollars. Trial shipments of ore were made to smelters in 1913, 1943, and 1969 (Moen, 1973, p. 25).

**TECTONIC SETTING:** The quartz monzonite was probably emplaced in a magmatic arc.

**ORE CONTROLS:** The main vein has an average thickness of 3 ft, strikes north, and dips 45-50W. On the south end of the vein, where shearing and faulting in the granodiorite is prominent, the vein consists of as much as 12 ft of banded quartz. The south end of the vein has been offset as much as 50 ft by faults. Ore minerals are generally sparsely scattered through the vein. However, in some parts of the vein, the ore minerals are concentrated in bands as much as several feet wide, which parallel the walls of the vein (Moen, 1973, p. 26).

**GEOLOGIC SETTING:** The vein is in the medium- to coarse-grained granodiorite and quartz monzonite of the Conconully pluton. The pluton is hydrothermally altered on its eastern edge near the town of Conconully (Stoffel, 1990 p. 24).

**COMMENTS:** The Lone Star mine has more than 2,000 ft of underground workings in the form of shafts, drifts, and crosscuts (Moen, 1973, p. 26).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- McFarland, C. R.; McLucas, G. B.; Rigby, J. G.; Stoffel, K. L., 1979, Directory of Washington mining operations, 1979: Washington Division of Geology and Earth Resources Information Circular 69, 100 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.



**Lucky Knock (444)**

ALTERNATE NAMES		DISTRICT	COUNTY
Lawrence			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Ellisforde	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 46' 16.16" N	119° 29' 7.85" W	SE1/4SW1/4 sec. 19, 38N, 27E	
LOCATION: west of Whitestone Mountain			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	limestone, phyllite	Permian	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Sb	stibnite sphalerite	calcite, quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
veinlets			

**PRODUCTION:** About 40 tons assaying 62% Sb were shipped in about 1907. A plant to transform stibnite to antimony oxide was built at the site between 1915 and 1917. Antimony oxide and about 500 tons of high-grade stibnite ore were shipped. One carload containing 42 tons of hand-sorted ore grading 30.47% Sb was shipped in 1941. In 1948, seven shipments totaling 47 tons of hand-sorted material that averaged 55.9% Sb were made (Purdy, 1951, p. 94-96).

**TECTONIC SETTING:** Sediments of the Spectacle Formation were deposited along an active continental margin.

**ORE CONTROLS:** Stibnite crystals are found in veinlets in fractured and generally silicified limestone beds and in faults in the limestone beds overlying phyllitic beds. Stibnite sometimes is present along bedding planes in limestone, and in some places, the stibnite replaces the limestone (Purdy, 1951, p. 99-101).

**GEOLOGIC SETTING:** The deposit is in a sequence limestone and phyllite of the Permian Spectacle Formation of the Anarchist Group (Stoffel, 1990, geol. map).

**COMMENTS:** Development consists of 1,000 ft of drifts, crosscuts, and winzes (Hunting, 1956, p. 19).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N.; Glover, S. L., 1921, The mineral resources of Washington, with statistics for 1919: Washington Geological Survey Bulletin 21, 155 p., 1 pl.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Shedd, Solon, 1924, The mineral resources of Washington with statistics for 1922; with an article on coal and coke, by G. W. Evans: Washington Division of Geology Bulletin 30, 224 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.
- White, D. E., 1962, Antimony in the United States (exclusive of Alaska and Hawaii): U.S. Geological Survey Mineral Investigations Resource Map MR-20, 1 sheet, scale 1:3,168,000, with 6 p. text.

**Magnetic** (412)

ALTERNATE NAMES		DISTRICT	COUNTY
Buckhorn Natural Aztec Neutral		Myers Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Buckhorn Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 28.60" N	118° 58' 43.18" W	NE1/4NW1/4 sec. 24, 40N, 30E	
LOCATION: elev. 5,000 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed limestone	limestone	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Buckhorn Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Fe Cu Ag Au	magnetite pyrrhotite pyrite chalcopyrite scheelite gold	garnet, epidote, pyrrhotite, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic replacement			

**PRODUCTION:** Small shipments of copper ore were made from the Neutral claim to a smelter in British Columbia. A total of 80,000 tons of magnetite were shipped from the Neutral claim between 1918 and 1950 (Moen, 1980, p. 47).

**TECTONIC SETTING:** Late Paleozoic sediments were deposited along an active continental margin.

**ORE CONTROLS:** Contact-metamorphic replacement of limestone in contact with the Cretaceous or Jurassic Buckhorn Mountain pluton. The magnetite zone is proximal to the pluton; tactite includes grossularite garnet, epidote, zoisite, quartz, and calcite. Tactite pods in places contain disseminated grains, pods, and pockets of intermixed magnetite, pyrrhotite, pyrite and chalcopyrite. Magnetite is irregularly distributed in the tactite zone, but near the Magnetic mine, it is roughly parallel to the northeast-trending contact, which appears to dip about 65SW (Moen, 1980, p. 47-49). Shear zones contain moderate amounts of sulfides and an estimated several million tons of iron ore (Hunting, 1956, p. 199).

**GEOLOGIC SETTING:** Magnetite bodies are located in a northwest-trending contact-metamorphic zone in limestones of Permian age adjacent to biotite-hornblende granodiorite of the Jurassic-Cretaceous Buckhorn Mountain pluton (Moen, 1980, p. 47; Stoffel, 1990).

**COMMENTS:** Development work consisted of three open pits, two adits, and numerous open cuts and prospects pits. The main pit is 280 ft long and 65-130 ft wide and has a maximum vertical face of 50 ft (Moen, 1980, p. 49). The Magnetite mine is part of the Buckhorn Mountain project drilled by Crown Resources Corp. in 1988-89. Other than iron the ore bodies contain traces to 0.02% Cu, traces to 0.15 oz/ton Au, and 0.05-0.40 oz/ton Ag (Moen, 1980, p. 49). Broughton (1943, p.17) estimated reserves of 4,826,000 tons of iron ore averaging 47.3% Fe, 1.57% S, 0.03% P, and 0.03% Ti.

**REFERENCES**

- Broughton, W. A., 1943, The Buckhorn iron deposits of Okanogan County, Washington—Results of a magnetic survey: Washington Division of Geology Report of Investigations 8, 21 p., 1 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Glover, S. L., 1942, Washington iron ores, a summary report: Washington Division of Mines and Mining Report of Investigations 2, 23 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.

## Okanogan

- Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels, and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.
- Whitwell, G. E.; Patty, E. N., 1921, The magnesite deposits of Washington, their occurrence and technology: Washington Geological Survey Bulletin 25, 194 p.
- Zapffe, Carl, 1949, A review—Iron bearing deposits in Washington, Oregon, and Idaho: Raw Materials Survey Resource Report 5, 89 p.
- Zoldok, S. W.; Cole, J. W.; Dougherty, E. Y., 1947, Iron deposits of Buckhorn Mountain, Meyers Creek mining district, Okanogan County, Washington: U.S. Bureau of Mines Report of Investigations 4051, 22 p.

**Mammoth (426)**

ALTERNATE NAMES		DISTRICT	COUNTY
Mammoth Claim		Conconully Conconully area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Conconully East	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 35' 14.63" N	119° 44' 54.49" W	sec. 30, 36N, 25E	
LOCATION: northeast of the Monitor property			
HOST ROCK: NAME	LITHOLOGY	AGE	
metamorphic complex of Conconully	schist, gneiss	pre-Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
alaskite and pegmatite			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Au		quartz	
DEPOSIT TYPE	MINERALIZATION AGE		
vein			

PRODUCTION: A shipment of ore to the smelter at Helena, Montana, in 1889 netted \$250 per ton (Hunting, 1956, p. 306).

TECTONIC SETTING: Late Triassic rocks formed along an active continental margin setting and are associated with island arcs.

ORE CONTROLS: Two veins are on the property; the upper one is 5 ft thick and the lower one is 6 ft thick (Hunting, 1956, p. 306).

GEOLOGIC SETTING: The quartz veins are in metasedimentary rocks of the metamorphic complex of Conconully (Stoffel, 1990)

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Mary Ann Creek placer (489)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Chesaw	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 24.90" N	119° 3' 6.44" W	sec. 30, 40N, 30E	
LOCATION: on Mary Ann Creek, 1 mi south of Chesaw			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	stream gravels	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: The deposit yielded \$40,000 in the 1880s (Hunting, 1956, p. 188).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action. Gold occurs from grass roots downward to bedrock. A 7-in.-wide clay seam, 4 ft above bedrock, acts as false bedrock, and better gold values are found above the clay (Hunting, 1956, p. 188).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.

**Mazama (439)**

ALTERNATE NAMES		DISTRICT	COUNTY
Lesley		Mazama	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mazama	1:62,500	Robinson Mtn	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 36' 52.91" N	120° 22' 55.20" W	SW1/4 sec. 17, SE1/4, SE1/4 sec. 18, E1/2 sec. 19, and W1/2 sec. 20, 36N, 20E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Fawn Peak stock	diorite, quartz diorite, plagioclase porphyry	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
andesite and andesite porphyry breccia fragments			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu	chalcopyrite	pyrite	
DEPOSIT TYPE	MINERALIZATION AGE		
porphyry system breccia pipe			

**PRODUCTION:** There is no recorded production.

**TECTONIC SETTING:** The Fawn Peak stock was probably the intrusive rock core of an island arc (Riedell, 1979).

**ORE CONTROLS:** Mineralization is confined to fractures and a breccia body. This appears to be primarily a porphyry copper deposit that has no other metal values. Reserves are given as nearly 150 million tons averaging 0.36% Cu. The overall waste to ore ratio is 2.5:1. Quartz veins 0.5 mi to the south at the Mazama Pride mine produced 37 tons of ore in 1931, which contained copper and 0.7 oz/ton Au.

**GEOLOGIC SETTING:** The Fawn Peak stock is an elongate, northwest-trending intrusion along the northeast side of the Methow River valley; it is situated along the axis of the Goat Peak syncline. The stock consists of finely to coarsely crystalline diorite, porphyritic quartz diorite, plagioclase porphyry, and intrusion breccia. The porphyritic rocks occur along the border of the stock. The stock intrudes the Winthrop Sandstone and the Midnight Peak Formation. The sandstone contains epidote (hornfels) alteration. Swarms of porphyritic sills and dikes extend as much as 1 km from the stock. The intrusive breccia is a pipelike body and consists of subrounded clasts as much as 10 in. long of diorite, quartz diorite, andesite, and andesite porphyry in a matrix of plagioclase porphyry. Three K-Ar magmatic biotite ages for the stock averaged 88 m.y. Altered rocks in the stock yield K-Ar secondary biotite ages of  $85.1 \pm 3.5$  m.y. and  $69.7 \pm 2.9$  m.y.

**COMMENTS:** This property consists of unpatented mining claims and is being explored (1990) by Vanderbilt Gold Corporation. Considerable drilling was conducted by Bear Creek, Noranda, Inspiration Development Co., and Quintana. Bear Creek and Inspiration spent \$700,000 on drilling.

**UNPUBLISHED INFORMATION:** Vanderbilt Gold Corporation, 1989, Mazama project description. This report is in DGER files.

**REFERENCES**

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**Methow** (397)

ALTERNATE NAMES		DISTRICT	COUNTY
London New London		Squaw Creek Hunter Mtn area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooper Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 25.19" N	120° 1' 6.13" W	E1/2, secs. 12 and 13, 30N, 22E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
basic dikes		Eocene?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au W Cu Ag Mo	scheelite gold chalcopyrite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
disseminated vein			

PRODUCTION: \$40,000 in gold was produced in 1940-41 (Hunting, 1956, p. 144).

TECTONIC SETTING: The Methow Gneiss is a Cretaceous intrusive body that has been metamorphosed to amphibolite grade (Bunning, 1990).

ORE CONTROLS: The deposit is made up of several veins. The New London vein is less than 1 ft wide and contains quartz, pyrite, gold, chalcopyrite, and scheelite. It lies along the footwall of a narrow basic dike that strikes N60E and dips 55NW. Scheelite crystals are disseminated in the quartz veins and are as much as 0.5 in. across. The veins are cut by several reverse faults. Other veins on the property are similarly mineralized and either parallel basic dikes or are contained within them (Culver and Broughton, 1945, p. 48-49).

GEOLOGIC SETTING: These veins are in basic dikes of probably Eocene age and which cut the Cretaceous Methow Gneiss (Bunning, 1990; Culver and Broughton, 1945).

COMMENTS: The quartz veins that make up the deposit were developed by several shafts and adits. More than 2,200 ft of underground workings were present; more than half of these were west of the Methow River (Hunting, 1956, p. 144).

## REFERENCES

- Barksdale, J. D., 1975, Geology of the Methow Valley, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 68, 72 p., 1 pl.
- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Mid Range (457)**

ALTERNATE NAMES		DISTRICT	COUNTY Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Gilbert	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 29' 7.33" N	120° 35' 49.04" W	sec. 34, 35N, 18E	
LOCATION: at the head of North Creek, elev. 6,000-7,760 ft, about 5 mi by trail north of Gilbert, at the terminus of the Twisp River road			
HOST ROCK: NAME	LITHOLOGY	AGE	
Golden Horn batholith	granite	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Zn	chalcopyrite sphalerite	pyrite, pyrrhotite, quartz, arsenopyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Eocene	

PRODUCTION: Ten tons of ore in 1939 and 22 tons in 1940 were shipped to the Tacoma smelter (Hunting, 1956, p. 144).

TECTONIC SETTING: The Golden Horn batholith was intruded into the Hozameen fault, the major fault bounding the southwestern side of the Methow basin. Radiometric dating suggests that magmatic crystallization occurred at approximately 47 m.y. K-Ar biotite, fission-track allanite, and Rb-Sr isochron ages that fall between 38 and 42 m.y. probably represent the age of postcrystallization hydrothermal alteration (Stoffel and McGroder, 1990).

ORE CONTROLS: The deposit consists of two mineralized quartz veins in granite (Hunting, 1956, p. 144).

GEOLOGIC SETTING: The veins are in granite of the Golden Horn batholith of Eocene age (Stoffel and McGroder, 1990).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.



**Minnie** (398)

ALTERNATE NAMES		DISTRICT	COUNTY
		Squaw Creek Leecher Creek area	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Twisp East	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48°15' 51.32" N	120° 3' 16.53" W	NW1/4NW1/4 sec. 23, 32N, 22E	
LOCATION: elev. 2,400 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Leecher Metamorphics		schist, amphibolite, marble	pre-Cretaceous
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	pyrite, marcasite, quartz	
Ag	sphalerite		
W	chalcopyrite		
Cu			
Zn			
S			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Shipped one carload of ore in 1941. A carload of ore shipped in November 1945 netted \$667.56 and gave assays of 0.46 oz/ton Au and 7.75 oz/ton Ag. A mining and gold heap-leach operation was worked by Cordilleran Exploration, Inc., in 1984-85 (Hunting, 1956, p.144; Joseph, 1986).

**TECTONIC SETTING:** Much of the rock of the Leecher Metamorphics has been recrystallized after mylonitization, probably due to the intrusion of the Okanogan complex (Bunning, 1990, p. 40).

**ORE CONTROLS:** The vein at the Minnie mine strikes N15E, dips 63SE, and parallels the orientation of the fabric in the Leecher Metamorphics. The porous white quartz vein averages 3 ft in width (Culver and Broughton, 1945, p. 52-53; Bunning, 1990, p. 28).

**GEOLOGIC SETTING:** The vein is parallel to the fabric in the biotite schist and marble in the Leecher Metamorphics. The deposit is reported to be within 1,000 ft of the Summit-Frazer gneiss of Cretaceous age (Culver and Broughton, 1945, p. 52-3; Bunning, 1990).

**COMMENTS:** The property was mined in 1984 and 1985 by Cordilleran Exploration, Inc. A small heap-leach system was operated by the company; dore was poured on site (Joseph, 1986).

## REFERENCES

- Barksdale, J. D., 1975, Geology of the Methow Valley, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 68, 72 p., 1 pl.
- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., 1986, Washington's mineral industry, 1985: Washington Geologic Newsletter, v. 14, no. 1-2, p. 1-17.
- Lemmon, D. M.; Tweto, O. L., 1962, Tungsten in the United States exclusive of Alaska and Hawaii: U.S. Geological Survey Mineral Investigations Resource Map MR-25, 1 sheet, scale 1:3,168,000, with 25 p. text.
- Washington Division of Mines and Geology, 1971, Directory of Washington mining operations, 1969-70: Washington Division of Mines and Geology Information Circular 46, 88 p.

**Mohawk (436)**

ALTERNATE NAMES		DISTRICT	COUNTY
Silver Peak Sunny Peak Chief Sunshine		Salmon River Mining	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Conconully West	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 34' 11.70" N	119° 47' 5.06" W	SW1/4SE1/4 sec. 31,T 36N, R 24E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Conconully pluton	granodiorite, quartz monzonite	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Cu Zn	galena sphalerite tetrahedrite	quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** The mine was worked as early as 1890, but mining ceased during the silver panic of 1893. Between 1951 and 1954, ore that contained as much as 60 oz/ton Ag was produced. In 1961 and 1967, small shipments of silver ore were made to the Cominco smelter in Trail, British Columbia (Moen, 1973).

**TECTONIC SETTING:** The Conconully pluton is a directionless post-tectonic body that was intruded into a major structural zone (Stoffel, K. L., DGER, oral commun., 1990).

**ORE CONTROLS:** The 1.5-3-ft-wide vein has a general strike of N30W and dips 30-40SW. Ore minerals are concentrated in bands as much as 1 ft wide that contain as much as 60 oz/ton Ag, 13% Pb, and 3% Cu (Moen, 1973, p. 24).

**GEOLOGIC SETTING:** The vein is in the biotite- and hornblende-bearing Conconully pluton of Cretaceous age (Stoffel, 1990).

**COMMENTS:** Two adits were worked at the property (Moen, 1973, p.24).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Jones, E. L., Jr., 1917, Reconnaissance of the Conconully and Ruby mining districts, Washington. *In* Contributions to economic geology (short papers and preliminary reports), 1916-Part I, metals and nonmetals except fuels: U.S. Geological Survey Bulletin 640, p. 11-36.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Montana** (451)

ALTERNATE NAMES		DISTRICT	COUNTY
		Mazama	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Mazama	1:62,500	Robinson Mountain	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 36' 59.65" N	120° 20' 59.85" W	SE1/4SW1/4 sec. 16, 36N, 20E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Fawn Peak stock	diorite	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au	chalcopyrite	pyrite, quartz, calcite, arsenopyrite, pyrrhotite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Cretaceous	

PRODUCTION: Produced an unknown amount in 1915 (Hunting, 1956, p. 68).

TECTONIC SETTING: The Fawn Peak stock was probably the intrusive rock core of an island arc (Riedell, 1979).

ORE CONTROLS: The deposit consists of veins in the actinolite alteration zone of the Mazama porphyry molybdenum system (Riedell, 1979).

GEOLOGIC SETTING: The Montana deposit is in the Cretaceous Fawn Peak stock. It is an elongate, northwest-trending intrusion along the northeast side of the Methow River valley; it is situated along the axis of the Goat Peak syncline (Stoffel and McGroder, 1990).

## REFERENCES

- Barksdale, J. D., 1975, Geology of the Methow Valley, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 68, 72 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Riedell, K. B., 1979, Geology and porphyry copper mineralization of the Fawn Peak intrusive complex, Methow Valley, Washington: University of Washington Master of Science thesis, 52 p., 4 pl.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**Mountain Beaver (458)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Billy Goat Mtn	1:24,000	Robinson Mountain	Concrète
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 47' 10.98" N	120° 19' 10.57" W	sec. 15, 38N, 20E	
LOCATION: (unsurveyed, approximate location) on Isabella Ridge adjacent to the Billy Goat prospect			
HOST ROCK: NAME	LITHOLOGY	AGE	
andesite of Isabella Ridge	andesite	Cretaceous?	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
plagioclase porphyry sill or dike			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	chalcopryrite	pyrite	
Cu	pyrite		
Ag	bornite		
Bi	chalcocite		
	gold		
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			
stockwork			
disseminated			

**PRODUCTION:** Small shipments were made in 1922, 1931, 1934, and 1935 (Hunting, 1956, p. 144). One shipment of 3,639 lb made in 1934 assayed 4.1% Cu, 2.08 oz/ton Au, and 1.9 oz/ton Ag (Staatz and others, 1971). Another shipment of crude ore yielded 1.81% Cu, 1.53 oz/ton Au, 1.13 oz/ton Ag, and a minor amount of bismuth (Hunting, 1956, p. 144). The Mountain Beaver is the only claim in the Billy Goat property to have produced (Staatz and others, 1971).

**TECTONIC SETTING:** The volcanics were deposited as part of an island arc.

**ORE CONTROLS:** Andesite tuff and breccia generally strike N30W and dip 70W to vertical. A plagioclase porphyry dike or sill is discontinuously exposed along the west edge of the Billy Goat property. Pyrite and copper minerals occur locally in 0.25-2-in.-wide quartz veinlets, which are located near steeply dipping, northwest-trending faults and fractures. The veinlets are disseminated through the fractured and layered andesite and the plagioclase porphyry (Staatz and other, 1971, p.111-112).

**GEOLOGIC SETTING:** Mineralization is present in andesite tuff and breccia of probable Cretaceous age and in a plagioclase porphyry dike or sill that intrudes the volcanic rocks (Staatz, 1990).

**COMMENTS:** The property was reportedly developed by two adits (Staatz and others, 1971).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Staatz, M. H.; Tabor, R. W.; Weis, P. L.; Robertson, J. F.; Van Noy, R. M.; Pattee, E. C., 1972, Geology and mineral resources of the northern part of the North Cascades National Park, Washington: U.S. Geological Survey Bulletin 1359, 132 p., 2 pl.
- Stoffel, K. L.; McGroder, M. F., compilers, 1990, Geologic map of the Robinson Mtn. 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-5, 39 p., 1 pl.

**Mountain Boy** (469)

ALTERNATE NAMES		DISTRICT	COUNTY
		Park City	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bald Knob	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 22' 34.87" N	118° 52' 4.34" W	NE 1/4 sec. 11, 33N, 31E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary rocks	argillite, limestone	Paleozoic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granite of Moses Mountain		Paleocene - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	pyrite, quartz	
Ag	sphalerite		
Sb	chalcopyrite		
Cu	tetrahedrite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Total production at the mine is reported to be four or five carloads with an average value of \$60/ton, presumably in Ag and Pb (Pardee, 1918, p.95).

TECTONIC SETTING: The granite of Moses Mountain is part of the Keller Butte suite of Holder and Holder (1988). Rocks of the Keller Butte suite were emplaced during regional ductile stretching associated with deformation in the metamorphic core complexes (Holder and Holder, 1988).

ORE CONTROLS: Rocks near the Mountain Boy mine trend northeast and dip 30-60NW. Quartz lenses at the mine vary from 0.5 to 18 in. in width and from 2 in. to 30 ft in length. The lenses contain irregularly scattered patches of sulfide minerals. The sulfide minerals show some banding and are brecciated. Both the quartz lenses and the metasedimentary rocks are cut by calcite-filled stringers (Pardee, 1918, p. 95).

GEOLOGIC SETTING: Veins are in metasedimentary rocks of probable Paleozoic age that are intruded by the nearby Paleocene-Eocene granite of Moses Mountain (Joseph, 1990).

COMMENTS: Three adits with short drifts, inclines, and other workings are located near each other at the Mountain Boy (Pardee, 1918)

## REFERENCES

- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

**Mountain Sheep (428)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Nighthawk	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 30.80" N	119° 42' 9.27" W	NE1/4NW1/4 sec. 28, 40N, 25E	
LOCATION: 0.75 mi north of the Ruby mine			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	greenstone	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Loomis pluton		Triassic - Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Cu Ag	pyrite malachite	quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: A few cars of ore were shipped prior to 1911 (Hunting, 1956, p. 307).

TECTONIC SETTING: The Kobau Formation was deposited along an active continental margin proximal to an island arc.

ORE CONTROLS: The deposit is on same fault as and is similar to the Ruby deposit. The vein trends N65W, dips 35SW, and can be traced for 3,000 ft along the outcrop. The vein is altered as much as 4 ft from the fault (Hunting, 1956, p. 307; Rinehart and Fox, 1972, p. 96).

GEOLOGIC SETTING: The Kobau Formation is intruded by granodiorite and tonalite of the Triassic-Jurassic Loomis pluton. Mineralization is present in a west-northwest trending-fault near the contact (Rinehart and Fox, 1972, geol. map).

COMMENTS: Three adits totaling 2,000 ft were driven (Rinehart and Fox, 1972).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

***Murray placer*** (490)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Trefry Canyon	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 29.93" N	119° 16' 23.84" W	sec. 11, 30N, 28E	
LOCATION: Kartar area			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	stream gravels	
DEPOSIT TYPE	MINERALIZATION AGE		
placer	Quaternary		

PRODUCTION: The placer has produced intermittently (Hunting, 1956, p. 188).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.

**Nevada (429)**

ALTERNATE NAMES		DISTRICT	COUNTY
War Eagle Peacock			Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Conconully East	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 30' 20.91" N	119° 44' 23.72" W	SW¼NE¼ sec. 30, 35N, 25E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
metamorphic complex of Conconully	schist, gneiss	pre-Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Conconully pluton		Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Cu Sb	galena tetrahedrite chalcopyrite sphalerite stromeyerite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		Cretaceous?	

**PRODUCTION:** In 1901 a carload of ore was shipped to a smelter; \$10/ton was received. In 1923-24 several small shipments of lead and copper ore were made; however, returns from the smelter did not exceed mining costs. In 1957 ore was concentrated at a flotation mill in Omak and shipped to the smelter (Moen, 1973, p. 20).

**TECTONIC SETTING:** Late Triassic rocks of the area formed in an active margin setting. The Conconully pluton is a directionless, post-tectonic pluton that was intruded into a major structural zone (Stoffel, K. L., DGER, 1990, oral commun.)

**ORE CONTROLS:** The 3-5-ft-thick vein has a general strike of N15W and dips 60E. The Nevada vein is the most persistent of several metallized quartz veins on Peacock Mountain. It is in hornblende-biotite schist and quartz diorite gneiss and parallels the foliation in the metamorphic rocks. Ore minerals are disseminated in the veins or concentrated in bands as much as 1 ft thick that parallel the walls of the vein. The silver is present mainly in galena. Tetrahedrite occurs near the surface. At 100 ft beneath the surface chalcopyrite is the dominant copper mineral (Moen, 1976, p. 20).

**GEOLOGIC SETTING:** The deposit is in hornblende- and biotite-bearing schist and quartz diorite gneiss near the contact with the Conconully pluton of Cretaceous age, according to Moen (1973). The deposit is shown as being within the metamorphic complex of Conconully by Stoffel (1990) and is shown by Rinehart to be (1981) at the contact of the Conconully pluton of Cretaceous age and metasedimentary and metavolcanic rocks of Permian and Triassic age that are elsewhere (Rinehart and Fox, 1976) shown as the metamorphic complex of Conconully.

**COMMENTS:** Underground workings at the Nevada mine consisted of four shafts, the deepest which is 220 ft, several hundred feet of drifts, and a 1,000-ft adit (Moen, 1973).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Rinehart, C. D., 1981, Reconnaissance geochemical survey of gully and stream sediments, and geologic summary, in part of the Okanogan Range, Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 74, 24 p., 3 pl.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.



**Nighthawk** (417)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 45.16" N	119° 38' 19.38" W	sec. 13, 40N, 25E, and sec. 18, 40N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Similkameen composite pluton	granodiorite	Jurassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granite			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb Ag	galena	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
breccia			

**PRODUCTION:** The mine has produced, but the amount is unknown (Hunting, 1956, p. 221).

**TECTONIC SETTING:** The Similkameen batholith is a concentrically zoned plutonic body with early mafic alkalic rocks bordering and intruded by calc-alkalic granitic rocks. The complex is undeformed and on the upper plate of the Okanogan valley fault (Buddington and Burmester, 1990).

**ORE CONTROLS:** Irregular mineralized bodies of friable quartz, which are commonly mixed with gouge, occur along the margins of a brecciated zone in the granodiorite. The principal ore has been mined from near the hanging wall; the contact with the granodiorite on the hanging wall is sharp, but it is gradational on the footwall. The brecciated zone is as much as 100 ft wide at one place (Umpleby, 1911, p. 90).

**GEOLOGIC SETTING:** In the Jurassic Similkameen composite pluton (Rinehart and Fox, 1972, geol. map).

**COMMENTS:** The deposit was developed by a 1,700 ft adit (Rinehart and Fox, 1972).

## REFERENCES

- Buddington, A. M.; Burmester, R. F., 1990, The Similkameen batholith: A mid-Jurassic, post-tectonic complex in the Quesnel terrane, north-central Washington and south-central British Columbia [abstract]: Geological Society of America Abstracts with Programs, v. 22, n. 3, p. 10-11.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**O.K. (371)**

ALTERNATE NAMES		DISTRICT	COUNTY
OK Copper OK O. K.			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48°59 35.24 N	119° 28 54.16 W	SE1/4 sec. 6, 40N, 27E	
LOCATION: 4 mi north of Oroville			
HOST ROCK: NAME	LITHOLOGY	AGE	
Silver Nail Lake pluton	quartz diorite	Jurassic - Cretaceous (?)	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Ag Au	chalcopyrite bornite tetrahedrite molybdenite scheelite	pyrite, quartz, serpentine	
DEPOSIT TYPE		MINERALIZATION AGE	
vein shear zone porphyry			

**PRODUCTION:** Three carloads of ore were shipped prior to 1911. Nineteen carloads of ore were shipped between 1917 and 1921; this ore averaged 7% Cu, 6 oz/ton Ag, and \$2/ton Au. Approximately 1,000-2,000 tons of 1-2% Cu ore remains on the dump and in the workings (Patty, 1921, p. 244).

**TECTONIC SETTING:** The Silver Nail Lake pluton was probably emplaced in a magmatic arc.

**ORE CONTROLS:** The deposit is in the silicified-sericitic zone of the Kelsey porphyry copper deposit (Roper, 1973). Mineralization is in a shear zone striking N10W and dipping 30-50 southwest. The quartz diorite is crumpled along the shear zone and consists of light-green quartz-mica and serpentine schist over a width of 6-10 ft. Pinch and swells have formed along which mineralization has developed; mineralization also cements broken wall rock and replaces schist along planes of schistosity (Patty, 1921, p. 245).

**GEOLOGIC SETTING:** Mineralization is in a shear zone in quartz diorite of the Jurassic-Cretaceous (?) Silver Nail Lake pluton (Fox, 1970).

**COMMENTS:** Part of the Kelsey porphyry copper-molybdenum deposit (Roper, 1973).

**REFERENCES**

- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.
- Fox, K., F., Jr., 1970, Geologic map of the Oroville quadrangle, Okanogan County, Washington: U. S. Geological Survey Open-File Report 70-128, 3 sheets, scale 1:62,500.
- Horton, F. W., 1916, Molybdenum, its ores and their concentration, with discussion of markets, prices, and uses: U.S. Bureau of Mines Bulletin 111, 132 p.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.
- Roper, M. W., 1973, Geology of the Kelsey copper-molybdenum property, Okanogan County, Washington: Montana State University Master of Science thesis, 97 p., 5 pl.

**Okanogan Free Gold (399)**

ALTERNATE NAMES		DISTRICT	COUNTY
Owasco Allison			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 57' 25.84" N	119° 29' 4.27" W	SE1/4NW1/4 sec. 19, 40N, 27E	
LOCATION: on N side of Similkameen River, elev. 1,200 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Spectacle Formation of the Anarchist Group		limestone, quartzite, schist	Permian
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Zn Pb	sphalerite galena gold	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Ore was produced in 1914, 1918, 1936, and 1938-1939 (Hunting, 1956, p. 145).

TECTONIC SETTING: Marine sedimentary rocks of the Spectacle Formation formed along an active continental margin.

ORE CONTROLS: A quartz vein as much as 12 ft wide cuts country rock of limestone, quartzite, and schist. Ore minerals are disseminated in the quartz (Hunting, 1956, p. 145).

GEOLOGIC SETTING: The deposit is in limestone, quartzite, and schist of the Permian Spectacle Formation (Fox, 1970, geol. map).

COMMENTS: Three adits and a glory hole were developed at the property (Hunting, 1956, p. 145).

## REFERENCES

- Fox, K., F., Jr., 1970, Geologic map of the Oroville quadrangle, Okanogan County, Washington: U. S. Geological Survey Open-file report 70-128, 3 sheets, scale 1:62,500.
- Handy, F. M., 1916, An investigation of the mineral deposits of northern Okanogan County: State College of Washington Department of Geology Bulletin 100, 27 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Palmer Summit** (400)

ALTERNATE NAMES		DISTRICT	COUNTY
Grand Summit		Palmer Mountain	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 51' 34.53" N	119° 33' 46.44" W	SE1/4SE1/4 sec. 21, 39N, 26E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Palmer Mountain Greenstone		gabbro	Permian - Triassic
ASSOCIATED IGNEOUS ROCK: DESCRIPTION			AGE
gabbro diorite			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu Pb	chalcopyrite galena	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Ore worth \$1,000 was produced prior to 1897; also produced in 1937 and 1939 (Hunting, 1956, p. 146).

TECTONIC SETTING: The Palmer Mountain Greenstone was deposited along a convergent continental margin.

ORE CONTROLS: The deposit consists of narrow and sparsely mineralized quartz veins in gabbro (Hunting, 1956, p. 146).

GEOLOGIC SETTING: A vein in metagabbro and greenstone of the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972, geol. map).

COMMENTS: Two adits and a shaft accessed the deposit (Hunting, 1956, p. 146).

## REFERENCES

- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

**Panama** (480)

ALTERNATE NAMES		DISTRICT Nespelem	COUNTY Okanogan
PRIMARY QUADRANGLE Armstrong Creek	SCALE 1:24,000	1/2° x 1° QUAD Omak	1° x 2° QUAD Okanogan
LATITUDE 48° 9' 16.91" N	LONGITUDE 119° 1' 40.78" W	SECTION, TOWNSHIP, AND RANGE center S1/2 sec. 27, 31N, 30E	
LOCATION:			
HOST ROCK: NAME porphyritic granodiorite of Manila Creek	LITHOLOGY granite, granodiorite	AGE (Paleocene?) - Eocene	
COMMODITIES Ag Au Sb	ORE MINERALS stephanite argentite silver chalcopyrite	NON-ORE MINERALS pyrite, quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE (Paleocene?) - Eocene	

**PRODUCTION:** A 12-ton shipment of hand-sorted ore was made to the Tacoma smelter; it returned 56.8 oz/ton Ag and 9.32 oz/ton Au (Colville Confederated Tribes, 1984). Shipments totaling 26 tons were made in 1919, 1921, and 1936 (Moen, 1976).

**TECTONIC SETTING:** The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). Rocks of the Keller Butte suite were emplaced during regional ductile stretching that resulted in the deformation associated with the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** Quartz lenses and veins are present in a 4-ft-wide shear zone in the granodiorite (Moen, 1976). The vein may be crosscut by an Eocene hypabyssal dacite dike (Colville Confederated Tribes, 1984).

**GEOLOGIC SETTING:** The veins are in the (Paleocene?)-Eocene porphyritic granodiorite of Manila Creek (Gulick and Korosec, 1990, geol. map).

## REFERENCES

- Colville Confederated Tribes Geology Department, 1984, Revised geology and mineral potential of the Colville Indian Reservation, Washington, 1984: Colville Confederated Tribes [Nespelem, Wash.], 2 v.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.

**Paymaster (459)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Squaw Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Cooper Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 3.71" N	120° 3' 14.15" W	SW1/4NW1/4 sec. 14, 30N, 22E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow Gneiss	tonalitic gneiss	Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold scheelite	quartz, iron oxide	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** An unknown amount of gold ore was produced from the oxidized zone (Hunting, 1956, p. 146).

**TECTONIC SETTING:** The Methow Gneiss protolith was intruded at moderate crustal levels and then metamorphosed to amphibolite grade (Bunning, 1990).

**ORE CONTROLS:** Mineralization is in a 3-ft-wide vein composed of sheared granitic rock and quartz. Iron-oxide bands from 1 to 3 in. wide on each wall of the vein carry free gold (Hunting, 1956, p. 146).

**GEOLOGIC SETTING:** The deposit is in the Methow Gneiss of probable Cretaceous age (Bunning, 1990).

**REFERENCES**

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

**Peacock** (481)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Conconully East	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 30' 25.98" N	119° 44' 27.62" W	NE¼ sec. 30, 35N, 25E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Conconully pluton metamorphic complex of Conconully	granite, quartz monzonite schist	Cretaceous pre-Jurassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Pb Zn Sb	galena tetrahedrite chalcopyrite sphalerite	pyrite, pyrrhotite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Extensive exploration and development was done; however, only several small shipments made (Moen, 1973, p. 20).

TECTONIC SETTING: The Conconully pluton is a directionless, post-tectonic plutonic body that was intruded into a major structural zone (Stoffel, K. L., DGER, oral commun., 1990).

ORE CONTROLS: The vein strikes N15W, dips 60E, and is at the contact between granite and quartz monzonite and schist (Moen, 1973; Stoffel, 1990).

GEOLOGIC SETTING: The vein is near the contact of the Conconully pluton and the metamorphic complex of Conconully (Stoffel, 1990, geol. map).

COMMENTS: The Peacock mine is also considered part of the Nevada mine.

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Pinnacle (401)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Loomis	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 51' 30.11" N	119° 37' 45.14" W	SE1/4 sec. 24, 39N, 25E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Palmer Mountain Greenstone	gabbro	Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Cu Pb Zn Ag	gold chalcopyrite sphalerite	pyrite, quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Produced \$200,000 worth of gold ore prior to 1910 (Hunting, 1956, p. 146).

**TECTONIC SETTING:** The Palmer Mountain Greenstone was deposited along a convergent continental margin.

**ORE CONTROLS:** The deposit consists of a vertical, N60E-trending, 4-10-ft-wide quartz vein; disseminated chalcopyrite and sphalerite are also present in altered gabbro and in sheared gabbro (Hunting, 1956, p. 146).

**GEOLOGIC SETTING:** The Pinnacle deposit is in metagabbro of the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972, geol. map).

**COMMENTS:** Development consists of 2,000 ft. of workings in three adits (Rinehart and Fox, 1972, p. 99).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Stebbins, R. H., 1951, Directory of Washington mining operations, 1951: Washington Division of Mines and Geology Information Circular 19, 75 p.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.



**Pittsburg (482)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Nespelem	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Armstrong Creek	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48°10 6.43 N	119° 2 36.36 W	SE1/4 sec. 21, 31N, 30E	
LOCATION: about 0.5 mi west of the Gould and Curry property			
HOST ROCK: NAME	LITHOLOGY	AGE	
porphyritic granodiorite of Manila Creek	granite, granodiorite	(Paleocene?) - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb	galena chalcopyrite malachite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein		(Paleocene?) - Eocene	

PRODUCTION: Produced an unknown amount of ore in 1914 (Huntting, 1956, p. 308).

TECTONIC SETTING: The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988).

ORE CONTROLS: Ore is in fractures in highly silicified and altered granite. A sizable silver-lead ore body was exposed (Huntting, 1956, p. 308).

GEOLOGIC SETTING: Veins and stockwork(?) in the porphyritic granodiorite of Manila Creek of (Paleocene?)-Eocene age (Gulick and Korosec, 1990, geol. map).

COMMENTS: The property is developed by open pits and a steeply inclined shaft (Huntting, 1956, p. 308).

## REFERENCES

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Pogue Flat (471)**

ALTERNATE NAMES		DISTRICT	COUNTY
Three Buttes			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Omak	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 26' 21.59" N	119° 33' 29.45" W	near SW corner, sec. 15, 34N, 26E	
LOCATION: elev. 1,400 ft.			
HOST ROCK: NAME	LITHOLOGY	AGE	
Pogue Mountain quartz monzonite	quartz monzonite, granite	Cretaceous?	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mn	pyrolusite rhodochrosite	quartz, calcite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein disseminated veinlets			

**PRODUCTION:** In 1916, the mine produced twenty-five 30-ton carloads of ore said to contain 20% Mn (Hunting, 1956, p. 261).

**TECTONIC SETTING:** The quartz monzonite was probably emplaced in a magmatic arc.

**ORE CONTROLS:** Two quartz veins are present in the decomposed granite; one is 2 ft wide and the other is 1 ft wide. Manganese minerals occur in stringers in the quartz and as disseminations in the granite (Hunting, 1956, p. 261).

**GEOLOGIC SETTING:** The deposit occurs in hydrothermally altered quartz monzonite of the Pogue Mountain quartz monzonite (Gulick and Korosec, 1990 p. 24).

**REFERENCES**

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1922, Deposits of manganese ore in Montana, Utah, Oregon, and Washington: U.S. Geological Survey Bulletin 725-C, p. 141-243.
- Patty, E. N.; Glover, S. L., 1921, The mineral resources of Washington, with statistics for 1919: Washington Geological Survey Bulletin 21, 155 p., 1 pl.
- Shedd, Solon, 1924, The mineral resources of Washington with statistics for 1922; with an article on coal and coke, by G. W. Evans: Washington Division of Geology Bulletin 30, 224 p.

**Poland China (402)**

ALTERNATE NAMES		DISTRICT	COUNTY
Molson Overtop		Myers Creek	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Chesaw	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 58' 26.59" N	119° 7' 21.04" W	SE1/4SE1/4 sec. 11, 40N, 29E	
LOCATION: elev. 3,850 ft			
HOST ROCK: NAME		LITHOLOGY	AGE
Spectacle Formation of the Anarchist Group		argillite	Permian
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Pb	pyrite galena marcasite gold	marcasite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** The Poland China was discovered in 1896. A 25-ton/day mill was erected in 1907 which produced a small shipment in 1914. Overtop Mining Co shipped 168 tons of concentrate containing 72 oz of Au and 70 oz of Ag in 1933-1934. From 1937 to 1939, 11 carloads of siliceous ore assaying 0.37 oz/ton Au were shipped to the smelter in Trail, British Columbia. Mining ceased in 1939 (Moen, 1980, p.35-37).

**TECTONIC SETTING:** Marine sedimentary rocks of the Spectacle Formation formed along an active continental margin.

**ORE CONTROLS:** The deposit contains a black to white, northwest-striking, east-dipping, 2-15-ft-wide quartz vein in graphitic argillite. The vein walls are generally well-defined, but in several places the vein gradually grades into the argillite; the vein also contains fragments of argillite. Locally, the vein is offset by north- and northeast-trending, high-angle faults. Ore minerals are sparsely disseminated in quartz (Moen, 1980, P. 37).

**GEOLOGIC SETTING:** The vein is in graphitic argillite of the Permian Spectacle Formation of the Anarchist Group (Moen, 1980).

**COMMENTS:** High-grade near-surface ore contained 0.15-30 oz/ton Au. Ore that was mined averaged only 0.24 oz/ton Au and 0.40 oz/ton Ag. Some wall rock contained from 0.018 to 0.06 oz/ton Au. Development consisted of more than 1,500 ft of adits, shafts, and crosscuts. This property is on patented mining claims (Moen, 1980, p. 38).

**REFERENCES**

- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1969, Mines and mineral deposits of Whatcom County, Washington: Washington Division of Mines and Geology Bulletin 57, 134 p., 14 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Poorman (485)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Park City	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bald Knob	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 22' 38.14" N	118° 52' 2.86" W	SE1/4NE1/4 sec. 11, 33N, 31E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
unnamed metasedimentary rocks		argillite	Late Paleozoic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Zn	galena	pyrite, quartz, calcite	
Pb	sphalerite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced an unknown amount of ore in 1937 (Hunting, 1956, p. 365).

TECTONIC SETTING: Late Paleozoic sediments were deposited along an active continental margin.

ORE CONTROLS: Breccia at the Poorman mine is cemented by calcite and is sparsely mineralized with pyrite and sphalerite. A 6-18-in.-wide vein trending N15E and consisting of banded quartz and some sulfide minerals was encountered in the adit (Pardee, 1918, p. 97).

GEOLOGIC SETTING: The deposit is in unnamed Late Paleozoic argillite.

## REFERENCES

- Gage, H. L., compiler, 1941, Some foreign and domestic zinc-lead mines that could supply zinc concentrates to a Pacific Northwest electrolytic zinc industry; Domestic western mines IV—The zinc-lead mines of Washington: U.S. Bonneville Power Administration Market Development Section, 235 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Prize** (418)

ALTERNATE NAMES		DISTRICT	COUNTY
Lakeview			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 55' 56.79" N	119° 38' 12.64" W	SE1/4 sec. 25 and NW1/4 sec. 36, 40N, 25E	
LOCATION: on the east side of the valley, elev. 3,200 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Kobau Formation	greenstone, phyllite	Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
metagabbro; metadiorite		Permian or Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	galena	pyrite, limonite, quartz	
Ag	chalcopyrite		
Au	malachite		
Cu	azurite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: Produced in 1906 and 1913 (Hunting, 1956, p. 222).

TECTONIC SETTING: The Kobau Formation was deposited proximal to an island arc along a convergent continental margin.

ORE CONTROLS: An east-striking vein, 15-in.-wide quartz vein in greenstone and schist; quartz is in large part intermixed with country rock, both enclosing fragments of it and extending into it as irregular stringers (Umpleby, 1911, p. 93).

GEOLOGIC SETTING: The vein is in greenstone and phyllite of the Triassic or Permian Kobau Formation, which is cut by metagabbro and metadiorite that is probably coeval with the greenstones (Rinehart and Fox, 1972).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.
- Washington Bureau of Statistics, Agriculture, and Immigration, 1903, Mines and mining: Washington Bureau of Statistics, Agriculture, and Immigration Biennial Report for 1903, p. 124-138.

**Rainbow (403)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 52' 6.00" N	119° 32' 32.30" W	NE¼ sec. 22, 39N, 26E	
LOCATION: between Palmer Mountain and Wannacut Lake, elev. 2,600 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	limestone, quartzite, schist	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
metagabbro, metadiorite		Permian - Triassic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu Pb	gold arsenopyrite chalcopyrite galena malachite	limonite, quartz, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

PRODUCTION: The mine is known to have produced (Hunting, 1956, p. 147).

TECTONIC SETTING: Marine sedimentary rocks of the Spectacle Formation formed along an active continental margin.

ORE CONTROLS: The vein consists of quartz lenses enclosed in limestone, quartzite, and schist. The lenses pinch and swell over short distances (Hunting, 1956, p. 147).

GEOLOGIC SETTING: The veins are in the Permian Spectacle Formation (Rinehart and Fox, 1972, geol. map).

COMMENTS: The property was developed by three adits and several crosscuts (Rinehart and Fox, 1972).

## REFERENCES

- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Handy, F. M., 1916, An investigation of the mineral deposits of northern Okanogan County: State College of Washington Department of Geology Bulletin 100, 27 p.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Ramore (470)**

ALTERNATE NAMES		DISTRICT	COUNTY
		Park City	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Central Peak	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 21' 44.90" N	118° 52' 28.89" W	NE1/4SW1/4 sec. 14, 33N, 31E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
unnamed metasedimentary rocks	black argillite, phyllite, limestone	Paleozoic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
granite of Moses Mountain		Paleocene - Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Pb	sphalerite	pyrite, pyrrhotite, quartz, fluorite, calcite	
Ag	galena		
Au	chalcopyrite		
Cu	tetrahedrite		
Zn			
DEPOSIT TYPE		MINERALIZATION AGE	
vein			
massive sulfide? (in core)			

PRODUCTION: No large shipment of ore reported (Hunting, 1956, p. 222).

TECTONIC SETTING: Metamorphosed sediments that were originally deposited along a Paleozoic active continental margin are intruded by the granite of Moses Mountain. The granite of Moses Mountain is part of the Keller Butte suite of Holder and Holder (1988). Rocks of the Keller Butte suite were emplaced during regional ductile stretching associated with deformation in the metamorphic core complexes (Holder and Holder, 1988).

ORE CONTROLS: Veins are at or near the contact of the metasedimentary rocks and the granite of Moses Mountain. The main vein strikes northeast and dips 45NW to vertical (Pardee, 1918).

GEOLOGIC SETTING: Black argillite and graywacke of probable Paleozoic age is intruded by the Paleocene-Eocene granite of Moses Mountain (Joseph, 1990, geol. map).

## REFERENCES

- Bancroft, Howland, 1914, The ore deposits of northeastern Washington: U.S. Geological Survey Bulletin 550, 215 p.
- Colville Confederated Tribes Geology Department, 1984, Revised geology and mineral potential of the Colville Indian Reservation, Washington, 1984: Colville Confederated Tribes [Nespelem, Wash.], 2 v.
- Gage, H. L., compiler, 1941, Some foreign and domestic zinc-lead mines that could supply zinc concentrates to a Pacific Northwest electrolytic zinc industry; Domestic western mines IV—The zinc-lead mines of Washington: U.S. Bonneville Power Administration Market Development Section, 235 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.

**Reco (404)**

ALTERNATE NAMES		DISTRICT Myers Creek	COUNTY Okanogan
PRIMARY QUADRANGLE Chesaw	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 57' 51.96" N	LONGITUDE 119° 2' 35.78" W	SECTION, TOWNSHIP, AND RANGE center, S1/2 sec. 16, 40N, 30E	
LOCATION: 0.5 mi north of Chesaw			
HOST ROCK: NAME Kobau Formation	LITHOLOGY greenstone, argillite, siltite	AGE Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION intermediate to acidic granitic rocks		AGE Mesozoic	
COMMODITIES Au Ag Cu	ORE MINERALS arsenopyrite gold chalcopyrite pyrite bornite	NON-ORE MINERALS marcasite, quartz, calcite	
DEPOSIT TYPE vein		MINERALIZATION AGE	

**PRODUCTION:** Most of work at mine took place between 1897 and 1900. From 1916 to 1953, the mine produced 1,015 tons of ore valued at \$23,000 from gold and \$365 from silver. The mine has been idle since 1953 (Moen, 1980, p. 40).

**TECTONIC SETTING:** Sedimentary and volcanic materials of the Kobau Formation were deposited proximal to an island arc along a convergent continental margin.

**ORE CONTROLS:** The 1-2-ft-wide vein strikes north and dips 50-70W. The vein consists of quartz and minor calcite and is sheared and fractured due to recurrent movement. Ore minerals are sparse and are present as disseminated grains, small blebs, and thin, dark bands in the quartz. Native gold was present in high-grade ore mined near the surface but absent in deeper parts of the vein. About 30 ft from the surface, the vein terminated at a north-dipping fault. Most of the vein is on the Gray Eagle claim. Gold content ranges from 0.04 to 16.8 oz/ton. Ore averaged 0.87 oz/ton Au, and most of it was hand sorted (Moen, 1980, p. 40).

**GEOLOGIC SETTING:** A north-striking vein in the Kobau Formation, which was intruded by Mesozoic intermediate and acidic composition dikes (Stoffel, 1990).

**COMMENTS:** Development work consists of several prospect pits, trenches, short adits, and two main adits. The longest adit (No. 2) was driven for 148 ft onto the Gray Eagle claim. The property is on patented mining claims (Moen, 1980, p. 40).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.



**Red Shirt (460)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Thrapp Mtn	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 20' 54.91" N	119° 59' 56.07" W	NE1/4 sec. 19, 33N, 23E	
LOCATION: on the lower part of the west slope of Pole Pick Hill, elev. 3,800 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Red Shirt gabbro	gabbro, diorite	Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu	pyrite chalcopyrite arsenopyrite	quartz	
DEPOSIT TYPE -		MINERALIZATION AGE	
vein shear zone			

**PRODUCTION:** The mine is credited with more than \$100,000 worth of production, and it produced intermittently for 50 years. The latest work was in 1936-1938 (Hunting, 1956, p. 147).

**TECTONIC SETTING:** The Red Shirt gabbro is part of the Frazer Creek complex. Rocks were strongly sheared along the Red Shirt thrust and have undergone retrograde metamorphism and alteration to epidote and chlorite (Gulick and Korosec, 1990).

**ORE CONTROLS:** The ore is in a quartz vein that is 1-5 ft wide (Hunting, 1956, p. 147).

**GEOLOGIC SETTING:** The vein and associated mineralization are in a shear zone in the Red Shirt gabbro of Jurassic-Cretaceous age (Gulick and Korosec, 1990).

**COMMENTS:** The property was developed by several drifts and crosscuts (Hunting, 1956, p. 147).

**REFERENCES**

- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hodges, L. K., 1897, Mining in the Pacific Northwest: Seattle Post Intelligencer, 116 p. [Facsimile reprinted 1967 in two volumes, Mining in eastern and central Washington, and, Mining in western Washington: Shorey Book Store, Seattle, Washington.]
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Review (405)**

ALTERNATE NAMES		DISTRICT Myers Ck	COUNTY Okanogan
PRIMARY QUADRANGLE Chesaw	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 57' 58.75" N	LONGITUDE 119° 2' 35.99" W	SECTION, TOWNSHIP, AND RANGE NE1/4SW1/4 sec. 16, 40N, 30E	
LOCATION: elev. 3,080 ft			
HOST ROCK: NAME Kobau Formation	LITHOLOGY greenstone	AGE Permian or Triassic (?)	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION Buckhorn Mountain pluton		AGE Jurassic - Cretaceous	
COMMODITIES Au Cu	ORE MINERALS gold chalcopryrite pyrite	NON-ORE MINERALS quartz, calcite	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: According to Landes and others (1902, p. 27), a smelter test, consisting of 50 tons of ore, averaged 0.86 oz/ton Au.

TECTONIC SETTING: The Kobau Formation was deposited proximal to island arcs along a convergent continental margin.

ORE CONTROLS: The vein consists of quartz and calcite with minor amounts of chalcopryrite, pyrite, and gold (Moen, 1980, p. 40).

GEOLOGIC SETTING: The vein is in greenstones of the Kobau Formation of Permian or Triassic age.

COMMENTS: Two adits, 360 and 813 ft in length, respectively, were driven into the hillside to intersect the vein at depth (Moen, 1980, p. 40).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II-The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.

## Roosevelt (413)

ALTERNATE NAMES		DISTRICT	COUNTY
Grant Maclean Teddy Roosevelt			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Buckhorn Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 37.54" N	118° 58' 28.93" W	secs. 24 and 25, 40N, 30E	
LOCATION: elev. 4,475 ft			
HOST ROCK: NAME	LITHOLOGY	AGE	
Spectacle Formation of the Anarchist Group	hornfels, calc-silicate	Permian	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Buckhorn Mountain pluton		Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Cu Au Ag Fe W	magnetite chalcopyrite pyrite scheelite	epidote, garnet	
DEPOSIT TYPE		MINERALIZATION AGE	
contact metamorphic			

**PRODUCTION:** Twelve carloads of hand-sorted ore were shipped to smelters in 1911; ore ran 7-9% Cu, 0.10-0.20 oz/ton Au, and 3.75-7.5 oz/ton Ag (Hunting, 1956, p. 199). In 1919-20 a total of 2,000 tons of iron ore was shipped (Moen, 1980, p. 54).

**TECTONIC SETTING:** The Spectacle Formation was deposited along an active continental margin.

**ORE CONTROLS:** Ore minerals occur at the contact between Permian metasedimentary rocks and the biotite-hornblende of the Jurassic-Cretaceous Buckhorn Mountain pluton. The contact metamorphic zone contains hornfels and calc-silicate rocks and minor quartzite, marble, and phyllite. The skarn and the ore bodies occur as irregular replacements that apparently favored beds of limestone and calcareous shale. Ore minerals are present as disseminated grains, as well as in pods and lenses. Small, sparsely disseminated grains of scheelite occur in garnet-epidote-rich areas of the contact zone.

**GEOLOGIC SETTING:** Contact metamorphic deposit at the contact of Permian metasedimentary rocks and the Jurassic-Cretaceous, biotite-hornblende Buckhorn Mountain pluton (Moen, 1980, p. 54; Stoffel, 1990).

**COMMENTS:** The main development at the Roosevelt mine consisted of an 855-ft lower adit with about 750 ft of drifts and crosscuts. About 200 ft higher in elevation are several open cuts and caved adits. The upper adits appear to be the earlier workings. The lower adit was driven in an attempt to intersect the downward extension of the ore bodies. However, these bodies apparently did not extend to the depth of the lower adit, although isolated occurrences of magnetite were found (Moen, 1980, p. 54). Broughton (1943, p. 20-21) estimated only about 2,000 tons of magnetite remains in the floor of the large stope of the lower adit. He also reported an area of moderately high dip needle readings S60E, 250 ft from the stope.

## REFERENCES

- Broughton, W. A., 1943, The Buckhorn iron deposits of Okanogan County, Washington—Results of a magnetic survey: Washington Division of Geology Report of Investigations 8, 21 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Ruby (430)**

ALTERNATE NAMES		DISTRICT	COUNTY
Pyrargyrite		Nighthawk	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Nighthawk	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 8.75" N	119° 41' 39.40" W	center E 1/2 sec. 28, 40N, 25E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Anderson Creek pluton	granodiorite	Jurassic - Cretaceous (?)	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	chalcopyrite	quartz, pyrite, arsenopyrite	
Au	galena		
Cu	sphalerite		
Pb	proustite		
Zn	pyrargerite		
	argentite		
	gold		
	malachite		
	azurite		

DEPOSIT TYPE	MINERALIZATION AGE
vein	
<p>PRODUCTION: The mine produced approximately 100,000 oz Ag from 1915 to 1922. High-grade ore contained as much as several hundred oz/ton Ag, but mill-run ore averaged only 10 oz/ton Ag. A mill (capacity 75-ton/day) was built in 1920. Prior to 1920, crude ore was shipped to smelters.</p> <p>TECTONIC SETTING: The quartz monzonite was probably emplaced in a magmatic arc.</p> <p>ORE CONTROLS: The quartz fissure vein is in a shear zone in granodiorite; the zone strikes N45W and dips 42SW. Ore minerals occur as sparsely scattered grains in the vein (Moen, 1976, p. 121).</p> <p>GEOLOGIC SETTING: The vein is in a shear zone in granodiorite of Jurassic-Cretaceous (?) age (Rinehart and Fox, 1972).</p> <p>COMMENTS: The mine was developed by at least 5,000 ft of underground workings and several stopes. The main adit intersects the vein at 950 ft from the portal and 550 ft beneath the outcrop. A 75-ton/day mill was built at the site in 1920 (Moen, 1976, p. 121).</p>	

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1976, Silver occurrences of Washington: Washington Division of Geology and Earth Resources Bulletin 69, 188 p.
- Purdy, C. P., Jr., 1951, Antimony occurrences of Washington: Washington Division of Mines and Geology Bulletin 39, 186 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Security (372)**

ALTERNATE NAMES		DISTRICT Palmer Mountain	COUNTY Okanogan
PRIMARY QUADRANGLE Loomis	SCALE 1:24,000	1/2° x 1° QUAD Oroville	1° x 2° QUAD Okanogan
LATITUDE 48° 50' 27.63" N	LONGITUDE 119° 38' 9.34" W	SECTION, TOWNSHIP, AND RANGE 1/2 N sec. 36, 39N, 25E	
LOCATION:			
HOST ROCK: NAME Palmer Mountain Greenstone	LITHOLOGY greenstone	AGE Permian - Triassic	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION granitic rocks, felsic to intermediate		AGE Jurassic - Cretaceous	
COMMODITIES Cu Pb Zn Au	ORE MINERALS chalcopyrite galena sphalerite gold	NON-ORE MINERALS pyrite, quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: Has produced (Hunting, 1956, p. 70).

TECTONIC SETTING: The Palmer Mountain Greenstone was deposited at a convergent continental margin.

ORE CONTROLS: The shear zone in argillite carries a vein 70 ft long and as much as 3 ft wide; it is sparsely mineralized. A similar vein occurs in the upper adit (Hunting, 1956, p. 70).

GEOLOGIC SETTING: The vein is in banded and schistose greenstone of the Permian-Triassic Palmer Mountain Greenstone that was intruded to the west by felsic to intermediate rocks of probable Jurassic-Cretaceous age (Rinehart and Fox, 1972, geol. map).

COMMENTS: Two adits were developed on the property (Rinehart and Fox, 1972).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. E., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.

**Sheridan** (431)

ALTERNATE NAMES		DISTRICT	COUNTY
Phil Sheridan		Sheridan	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bodie Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 46' 42.73" N	118° 51' 17.17" W	SW1/4NW1/4 sec. 24, 38N, 31E	
LOCATION: in a small gulch 1 mi north of the East Fork Toroda Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Klondike Mountain Formation	andesite, dacite	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Au	silver mineral	quartz, calcite, pyrite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein, epithermal breccia		Eocene	

**PRODUCTION:** Between 1906 and 1919, shipments of high-grade gold-silver ore were made to smelters. In 1918 a 50-ton/day flotation mill was built at the mine, and concentrates averaging as much as 300 oz/ton Ag were shipped until 1920. For this period, the mine production was estimated at \$60,000 to \$100,000 (Moen, 1980, p. 72).

**TECTONIC SETTING:** The Toroda Creek graben is a volcano-tectonic depression, which resulted from east-west extension during the Eocene (Holder and others, 1989).

**ORE CONTROLS:** Silver minerals occur in a series of closely spaced, silicified and pyritized shear zones in altered and silicified andesites and dacites. Brecciation of the rocks accompanied the shearing, and clasts of the breccia have been considerably altered and silicified by hydrothermal solutions. The main vein is 1-7 ft wide, strikes N20E, and dips 50NW. During mining operations it was necessary to closely assay the vein material in order to establish borders of the ore shoots (Moen, 1980, p. 72).

**GEOLOGIC SETTING:** The ore is in altered and silicified volcanic rocks of the Eocene Klondike Mountain Formation. The geology here is similar to that described for the American Flag mine).

**COMMENTS:** A 50-ton/day flotation mill was built in 1918. Most adits have heading of N35E. The vein was mined from several levels over a vertical distance of about 500 ft and along strike for about 400 ft. Several raises connect the different levels of the mine (Moen, 1980, p. 72).

**REFERENCES**

- Holder, R. W.; Gaylord, D. R.; Holder, G. A. M., 1989, Plutonism, volcanism, and sedimentation associated with core complex and graben development in the central Okanogan Highlands, Washington. *In* Joseph, N. L.; and others, editors, 1989, *Geologic guidebook for Washington and adjacent areas*: Washington Division of Geology and Earth Resources Information Circular 86, p. 187-200.
- Hunting, M. T., 1956, *Inventory of Washington minerals—Part II, Metallic minerals*: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, *Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington*: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Patty, E. N., 1921, *The metal mines of Washington*: Washington Geological Survey Bulletin 23, 366 p.

**Shotwell placer** (491)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bridgeport	1:24,000	Omak	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 2' 3.42" N	119° 40' 58.45" W	NW1/4 sec. 10, 29N, 25E	
LOCATION: on a terrace about 100 ft above the Columbia River			
HOST ROCK: NAME		LITHOLOGY	AGE
Quaternary alluvium		sand and gravel	Quaternary
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	stream gravels	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Produced an unknown of gold in 1954. A small amount of gold was produced as byproduct of aggregate production for the Chief Joseph Dam (Hunting, 1956, p. 188).

TECTONIC SETTING: Heavy minerals were deposited in river gravels.

ORE CONTROLS: Heavy-mineral concentration by stream action.

## REFERENCES

Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Silver Bell** (406)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Bodie Mountain	1:24,000	Republic	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 45' 34.50" N	118° 50' 17.97" W	sec. 25, 38N, 31E	
LOCATION: 12 mi northwest of Republic			
HOST ROCK: NAME	LITHOLOGY	AGE	
Klondike Mountain Formation	trachyte	Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag	chalcopyrite galena tetrahedrite pyrite	purple fluorite, sanidine, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
vein, epithermal		Eocene	

**PRODUCTION:** High-grade gold ore was reported shipped from a surface pit prior to 1907. In 1940 a 29-ton shipment gave net smelter returns of \$244.03 (Hunting, 1956, p. 148). There was a small amount of production in the 1980s.

**TECTONIC SETTING:** East-west extension during the Eocene resulted in formation of the Toroda Creek graben and other structures in which a thick section of volcanic and sedimentary materials was deposited and preserved (Holder and others, 1989).

**ORE CONTROLS:** The deposit is contained in brecciated and silicified trachyte. The breccia has a general northeast strike and dips about 45W. The breccia is about 50-75 ft wide and is cemented with quartz, sanidine, and as much as 4% fluorite. The deposit is near the center of the Zalla M-American Flag mineralized belt, which is almost a mile long and several hundred feet wide (Moen, 1980, p. 69).

**GEOLOGIC SETTING:** Geologic setting is similar to that at the American Flag mine where volcanic rocks in this part of the Toroda Creek graben plot as rhyolite on a TAS diagram, but contain sparse plagioclase and mafic phenocrysts in an aphanitic groundmass that contains abundant plagioclase microlites. Plagioclase phenocrysts are commonly strongly sericitized, and mafic minerals are generally replaced by some combination of chlorite, calcite, epidote, and sphene; shapes of altered mafic minerals suggest that they were originally pyroxene. The presence of silica stringers and veinlets in some of these rocks suggests that silica was introduced during alteration (Stoffel, 1990, p. 11).

**COMMENTS:** Underground workings consist of two adits, a 50-ft shaft, and several shallow shafts. Moen (1980, p. 70) reported that drilling done by the owner of the property in 1978 delineated 124,000 tons of rock having an average value of 5.46 oz/ton Ag and 0.014 oz/ton Au. Moen (1980, p. 69) also reported assays on the siliceous breccia that range from 0-30 oz/ton Ag; as much as 120 ft of breccia assayed 3.0 oz/ton Ag and, a 13-ft section assayed as much as 26.55 oz/ton Ag.

## REFERENCES

- Holder, R. W.; Gaylord, D. R.; Holder, G. A. M., 1989, Plutonism, volcanism, and sedimentation associated with core complex and graben development in the central Okanogan Highlands, Washington. *In* Joseph, N. L.; and others, editors, 1989, Geologic guidebook for Washington and adjacent areas: Washington Division of Geology and Earth Resources Information Circular 86, p. 187-200.
- Hunting, M. T., 1949, Directory of Washington mining operations, 1949: Washington Division of Mines and Geology Information Circular 17, 62 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1980, Myers Creek and Wauconda mining districts of northeastern Okanogan County, Washington: Washington Division of Geology and Earth Resources Bulletin 73, 96 p., 6 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Republic 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-10, 62 p., 1 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.



**Silver Bluff** (432)

ALTERNATE NAMES		DISTRICT	COUNTY
		Galena	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Riverside	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 34' 48.71" N	119° 36' 30.09" W	NE 1/4 sec. 31, 36N, 26E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Cave Mountain Formation		limestone, quartzite	Triassic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag	argentite	quartz	
Cu	chalcocite		
Au	stromeyerite		
	azurite		
	malachite		

DEPOSIT TYPE	MINERALIZATION AGE
vein	
PRODUCTION: Credited with \$80,000 worth of production by the end of 1923 (Hunting, 1956, p. 310).	
TECTONIC SETTING: The Cave Mountain Formation was deposited along an active continental margin proximal to an island arc.	
ORE CONTROLS: A 2-ft-wide vein was exposed at the surface (Hunting, 1956, p. 310).	
GEOLOGIC SETTING: The vein cuts limestone and quartzite of the Triassic Cave Mountain Formation (Rinehart and Fox, 1976, geol. map).	
COMMENTS: An inclined shaft was driven 65 ft at the property (Bethune, 1892).	

## REFERENCES

- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.

**Silver Cliff** (483)

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Belvedere	1:24,000	Nespelem	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 6' 13.90" N	118° 57' .55" W	sec. 17, 30N, 31E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
porphyritic granodiorite of Manila Creek unnamed metasedimentary rocks	granite, granodiorite argillite, quartzite, marble	Eocene - (Paleocene?) Paleozoic	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb	galena chalcopyrite	pyrite, quartz, sericite, rhodochrosite	
DEPOSIT TYPE		MINERALIZATION AGE	
vein disseminated		(Paleocene ?) - Eocene	

**PRODUCTION:** One carload of ore was said to have been shipped (Hunting, 1956, p. 310).

**TECTONIC SETTING:** The porphyritic granodiorite of Manila Creek is part of the Keller Butte suite of Holder and Holder (1988). The Keller Butte suite is contemporaneous with the formation of the metamorphic core complexes (Holder and Holder, 1988).

**ORE CONTROLS:** Mineralization is present as disseminated grains of chalcopyrite, pyrite, and galena in metasedimentary rocks and in veins that cut the metasedimentary rocks and granodiorite. Several veins at the property range in thickness from several inches to 3 ft, strike N70W to N70E, and dip to the south (Pardee, 1918, p. 75-76). Hydrothermal alteration extends for several feet on either side of the veins (Colville Confederated Tribes, 1984, p. 53). Altered granodiorite near one open cut contains as much as 30 ft of rock that is sericitized and partially replaced by quartz and contains disseminated grains of pyrite, galena, and chalcopyrite (Pardee, 1918, p. 76).

**GEOLOGIC SETTING:** Roof pendants consisting of Paleozoic argillite, quartzite, and marble overlie the (Paleocene?) - Eocene porphyritic granodiorite of Manila Creek (Joseph, 1990).

## REFERENCES

- Colville Confederated Tribes Geology Department, 1984, Revised geology and mineral potential of the Colville Indian Reservation, Washington, 1984: Colville Confederated Tribes [Nespelem, Wash.], 2 v.
- Holder, R. W.; Holder, G. A. M., 1988, The Colville batholith—Tertiary plutonism in northeast Washington associated with graben and core complex (gneiss dome) formation: Geological Society of America Bulletin, v. 100, no. 12, p. 1971-1980.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Joseph, N. L., compiler, 1990, Geologic map of the Nespelem 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-16, 47 p., 1 pl.
- Pardee, J. T., 1918, Geology and mineral deposits of the Colville Indian Reservation, Washington: U.S. Geological Survey Bulletin 677, 186 p., 1 pl.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

**Silver Ledge (462)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Hungry Mtn	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 12' 15.88" N	120° 10' 11.53" W	near center sec. 11, 31N, 21E	
LOCATION: 0.25 mi downstream from the Antimony Queen mine, elev. 300 ft. above Gold Creek			
HOST ROCK: NAME	LITHOLOGY	AGE	
Newby Group	graywacke	Jurassic - Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag  -	silver sulfide cerargyrite bromyrite pyrite arsenopyrite	quartz	

DEPOSIT TYPE	MINERALIZATION AGE
vein	
PRODUCTION: Produced 2 tons of unsorted ore prior to 1921 (Hunting, 1956, p. 149).	
TECTONIC SETTING: The Newby Group was deposited along an active continental margin in or near an island arc.	
ORE CONTROLS: The deposit consists of quartz veins in graywacke and shale. The vein at the collar of the shaft is in a 5-ft-wide shear zone and consists of two parts: one is 1.5 ft wide and is at the footwall, and the other is 4 in. wide and is on the hanging wall (Hunting, 1956, p. 148).	
GEOLOGIC SETTING: The vein is in metasedimentary rocks of the Jurassic-Cretaceous Newby Group (Bunning, 1990, geol. map).	
COMMENTS: The property is developed by a 150-ft inclined shaft and a 1,500-ft crosscut (Hunting, 1956).	

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Patty, E. N., 1921, The metal mines of Washington: Washington Geological Survey Bulletin 23, 366 p.

**Silver Mountain (433)**

ALTERNATE NAMES		DISTRICT	COUNTY
Silver Star			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Aeneas Lake	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 44' 46.12" N	119° 33' 28.98" W	W1/2 sec. 34, 38N, 26E, and NW1/4 sec. 3, 37N, 26E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Anarchist Group		metasedimentary rocks	Permian
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Ag Pb Zn Au	galena sphalerite	pyrite, quartz	
DEPOSIT TYPE		MINERALIZATION AGE	
veins stockwork			

PRODUCTION: Produced in 1943 (Huntting, 1956, p. 310).

TECTONIC SETTING: Marine sediments of the Anarchist Group were deposited along an active continental margin.

ORE CONTROLS: Quartz veins and silicified zones 1-20 ft wide are present in metasedimentary rocks; the veins are sparsely mineralized (Huntting, 1956, 310).

GEOLOGIC SETTING: Mineralized veins are in the clastic metasedimentary rocks of the Permian Anarchist Group (Rinehart and Fox, 1976, geol. map).

COMMENTS: A 400-ton/day mill was under construction in 1952. A sample for a mill test showed 28.79 oz/ton Ag, 0.351 oz/ton Au, 1.11% Pb, and 0.59% Zn (Huntting, 1956, p. 310).

## REFERENCES

- Huntting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Purdy, C. P., Jr., 1952, Directory of Washington mining operations, 1952: Washington Division of Mines and Geology Information Circular 20, 75 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.

**Similkameen placers (437)**

ALTERNATE NAMES		DISTRICT	COUNTY
			Okanogan
PRIMARY QUADRANGLE	SCALE	½° x 1° QUAD	1° x 2° QUAD
Oroville	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 56' 7.95" N	119° 26' 27.50" W		
LOCATION: along the Similkameen River between Oroville and Nighthawk			
HOST ROCK: NAME	LITHOLOGY	AGE	
Quaternary alluvium	sand and gravel	Quaternary	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	sand and gravel	
DEPOSIT TYPE		MINERALIZATION AGE	
placer		Quaternary	

PRODUCTION: Reportedly \$500,000 in the few years following 1859. Intermittent to 1955 (Hunting, 1956, p. 188).

TECTONIC SETTING: Deposition of heavy minerals in river gravels.

ORE CONTROLS: Heavy mineral concentration by stream action.

GEOLOGIC SETTING: Deposits are in river bars and lower terraces of the Similkameen River (Hunting, 1956, p. 188).

## REFERENCES

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Bethune, G. A., 1891, Mines and minerals of Washington—Annual report of G. A. Bethune, first state geologist: Washington State Printer, 122 p.
- Bethune, G. A., 1892, Mines and minerals of Washington—Second annual report of G. A. Bethune, state geologist: Washington State Printer, 183 p.
- Handy, F. M., 1916, An investigation of the mineral deposits of northern Okanogan County: State College of Washington Department of Geology Bulletin 100, 27 p.
- Umpleby, J. B., 1911, Part I. Geology and ore deposits of the Myers Creek mining district; Part II—Geology and ore deposits of the Oroville-Nighthawk mining district: Washington Geological Survey Bulletin 5, 111 p.

**Spokane** (463)

ALTERNATE NAMES		DISTRICT	COUNTY
Gold Crown			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Twisp West	1:24,000	Twisp	Concrete
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 22' 25.96" N	120° 9' 16.26" W	sec. 12, 33N, 21E	
LOCATION: on Twisp River road, 2 mi west of Twisp			
HOST ROCK: NAME	LITHOLOGY	AGE	
Newby Group	dacite, andesite	Jurassic - Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
Alder stock		Cretaceous	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	sphalerite	quartz, calcite	
Ag	arsenopyrite		
Pb	chalcopyrite		
Zn	pyrite		
Cu	galena		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			
massive sulfide?			

**PRODUCTION:** Small amounts of ore were shipped in 1939 and 1954. Six tons shipped in 1941 contained 0.40 oz/ton Au, 20.25 oz/ton Ag, 4.5% Zn, and 3.8% Pb (Hunting, 1956, p. 149; Burnet, 1976, table 2).

**TECTONIC SETTING:** The Newby Group was deposited at or near volcanic island arcs.

**ORE CONTROLS:** An irregular quartz-calcite vein in dacite ranges from a few inches to 3 ft thick. Ore minerals are in bunches in the vein (Hunting, 1956, p. 149). The host rock is silicified dacite breccia. Some mineralization is also in a fault zone at the contact with the Alder stock and in quartz-sericite phyllite that is on strike with the Alder mine (Burnet, 1976).

**GEOLOGIC SETTING:** Mineralization is in silicified dacite breccia of the Jurassic-Cretaceous Newby Group; in places the Newby Group is contact metamorphosed by the Alder Creek stock, which was dated at  $137 \pm 4$  m.y. (Burnet, 1976; Bunning, 1990).

## REFERENCES

- Banta, H. E., 1956, Directory of Washington mining operations, 1956: Washington Division of Mines and Geology Information Circular 25, 87 p.
- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Burnet, F. W., 1976, Felsic volcanic rocks and mineral deposits in the Buck Mountain Formation andesites, Okanogan County, Washington: University of Washington Master of Science thesis, 26 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Landes, Henry; Thyng, W. S.; Lyon, D. A.; Roberts, Milnor, 1902, Annual report for 1901, in six parts; Part II—The metalliferous resources of Washington, except iron: Washington Geological Survey, 123 p.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

**St. Anthony** (461)

ALTERNATE NAMES		DISTRICT Squaw Creek	COUNTY Okanogan
PRIMARY QUADRANGLE Cooper Mtn	SCALE 1:24,000	1/2° x 1° QUAD Twisp	1° x 2° QUAD Concrete
LATITUDE 48° 7' 23.99" N	LONGITUDE 120° 7' 13.19" W	SECTION, TOWNSHIP, AND RANGE SW1/4SW1/4 sec. 5, 30N, 22E	
LOCATION:			
HOST ROCK: NAME Methow Gneiss	LITHOLOGY tonalitic gneiss	AGE Cretaceous	
COMMODITIES Au Ag Cu	ORE MINERALS gold	NON-ORE MINERALS quartz	
DEPOSIT TYPE vein		MINERALIZATION AGE	

PRODUCTION: The mine produced an unknown amount of ore in 1934 (Hunting, 1956, p. 148).

TECTONIC SETTING: The Methow Gneiss protolith was intruded at moderate crustal levels, then metamorphosed along with other rocks in the area to amphibolite grade (Bunning, 1990).

ORE CONTROLS: The vein is in gneiss and is 5 ft wide (Hunting, 1956, p. 148).

GEOLOGIC SETTING: The St. Anthony mine is in the Methow Gneiss of probable Cretaceous age (Bunning, 1990).

COMMENTS: Ore from the surface to a depth of 35-ft is said to run \$50/ton in Au (Hunting, 1956, p. 148).

## REFERENCES

- Bunning, B. B., compiler, 1990, Geologic map of the east half of the Twisp 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-9, 51 p, 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Washington Division of Mines and Mining, 1941, Directory of Washington metallic mining properties: Washington Division of Mines and Mining Information Circular 7, 74 p.

***Starr*** (440)

ALTERNATE NAMES		DISTRICT	COUNTY
Silver Tip Andy Starr		Conconully Galena	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Aeneas Lake	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 42' 59.34" N	119° 35' 25.13" W	SE1/4 sec. 8, 37N, 26E	
LOCATION:			
HOST ROCK: NAME	LITHOLOGY	AGE	
Aeneas Creek pluton, central phase	quartz monzonite and granodiorite	Cretaceous	
Aeneas Creek pluton, border phase	quartz diorite, diorite, gabbro	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
pegmatite and aplite dikes and sills			
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Mo W	molybdenite molybdite scheelite chalcopyrite	quartz, limonite, hematite, sericite; kaolinization, silicification	
DEPOSIT TYPE		MINERALIZATION AGE	
disseminated veins fractures breccia porphyry system			

**PRODUCTION:** Three thousand tons of hand-sorted ore, which averaged 1.0% MoS<sub>2</sub>, was shipped in 1939 (Hunting, 1956, p. 271). Material from dumps of the No. 1 and No. 2 adits was shipped after 1944.

**TECTONIC SETTING:** In British Columbia to the north, the Quesnel trough partly controls the distribution of major porphyry copper-molybdenum deposits. The Starr molybdenum property lies on the projection of this structural trough.

**ORE CONTROLS:** The deposit is an elliptical body striking northwest and is as much as 130 ft wide and 300-440 ft long. Mineralization is disseminated in silicified granodiorite, in fractures, associated with breccia, and in quartz veins. The deposit is oxidized to a depth of 30 ft with half of the molybdenum values molybdenite. Sampling in the No. 1 adit, which is entirely within the mineralized zone, gave a weighted average of 0.51% MoS<sub>2</sub>. Samples from a winze from the No. 2 adit returned assays of 0.81-1.91% MoS<sub>2</sub>. Arsenic assays by the USGS for ten samples ranged from 0.01 to 0.04%. On the basis of trench and underground sampling, indicated and inferred reserves to a depth of 300 ft total 800,000 tons averaging 0.30% MoS<sub>2</sub>. The deposit is open at depth. There is a potential for 2 million tons averaging 0.26% MoS<sub>2</sub> (Wilbur Hallauer, personal commun., Aug. 15, 1967).

**GEOLOGIC SETTING:** The northwest-trending Aeneas Creek pluton is 1 mi wide and 3 mi long. It consists of a central phase and a border phase that is in contact with Permian Anarchist Group metasedimentary rocks. The deposit is in the central phase of the pluton. The pluton consists of medium crystalline biotite-hornblende quartz monzonite and granodiorite. The pluton has yielded K-Ar biotite ages of 92.7 ± 6.6 m.y. and 98.3 ± 3.6 m.y. A concordant K-Ar hornblende age was 92.3 ± 4.3 m.y.

**COMMENTS:** The property consists of unpatented mining claims and is owned (1990) by Wilbur G. Hallauer of Oroville. In 1928 the property was explored by Molybdenum Corporation of America and in 1935 and 1936 by Titanium Alloy Manufacturing Co. Carl Lundstrom mined the property in 1939. In total there are 2,700 feet of underground workings. In 1959 six short diamond drill holes were completed. From 1966 to 1967 Cambri Mining and Development Co. Ltd. leased the property and drilled six 200-ft holes.

**UNPUBLISHED INFORMATION:** Fawley, A.P., 1966, The Starr molybdenum mine, Washington. This report is in DGER files.

**REFERENCES**

- Creasey, S. C., 1945, Geology of the Starr molybdenum mine, Okanogan County, Washington: U.S. Geological Survey Open-File Report, 11 p., 5 pl.
- Culver, H. E.; Broughton, W. A., 1945, Tungsten resources of Washington: Washington Division of Geology Bulletin 34, 89 p., 23 pl.



- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Moen, W. S., 1973, Conconully mining district of Okanogan County, Washington: Washington Division of Mines and Geology Information Circular 49, 42 p.
- Purdy, C. P., Jr., 1954, Molybdenum occurrences of Washington: Washington Division of Mines and Geology Report of Investigations 18, 118 p., 6 pl.
- Rinehart, C. D.; Fox, K. F., Jr., 1976, Bedrock geology of the Conconully Quadrangle, Okanogan County, Washington: U.S. Geological Survey Bulletin 1402, 58 p., 1 pl.
- Stoffel, K. L., compiler, 1990, Geologic map of the Oroville 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-11, 58 p., 1 pl.

**Sullivan** (464)

ALTERNATE NAMES		DISTRICT	COUNTY
Pateros			Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Pateros	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 4' 50.48"N	119° 59' 15.05" W	SW1/2 sec. 20, 30N, 23E	
LOCATION: 5 mi northwest of Pateros, on the east side of the Methow River			
HOST ROCK: NAME	LITHOLOGY	AGE	
Methow gneiss	tonalitic gneiss	Cretaceous	
ASSOCIATED IGNEOUS ROCK: DESCRIPTION		AGE	
dacite dikes		Eocene	
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au Ag Cu	pyrite chalcopyrite	quartz, carbonate minerals	
DEPOSIT TYPE		MINERALIZATION AGE	
shear zone			

**PRODUCTION:** Reportedly, 60 carloads of ore were produced prior to 1897; the total was valued at \$72,000. Seven tons of ore that were shipped in 1940 assayed 0.63 oz/ton Au, 0.80 oz/ton Ag, and 0.17% Cu. Ore was also shipped in 1941 (Hunting, 1956, p. 149).

**TECTONIC SETTING:** The Methow Gneiss is a post-tectonic body that was originally intruded at moderate to deep levels and then metamorphosed to amphibolite-facies grade rock (Gulick and Korosec, 1990, p. 38).

**ORE CONTROLS:** The deposit consists of shear zones as much as 6 ft wide but averaging less than 3 ft. The shear zones are in granite along the margins of "andesite" dikes. The shear zones also contain small, scattered pods of shipping-grade ore (Hunting, 1956, p. 149). The andesite dikes noted in Hunting (1956) are probably dacite dikes (Gulick and Korosec, 1990, geol. map).

**GEOLOGIC SETTING:** The Methow Gneiss of Cretaceous age is sheared and intruded by hypabyssal dikes of probable Eocene age near the deposit (Gulick and Korosec, 1990, geol. map).

**COMMENTS:** The deposit may also be part of the Friday mine. The Sullivan mine was developed by 1,200 ft of workings on three levels (Hunting, 1956).

## REFERENCES

- Gulick, C. W.; Korosec, M. A., compilers, 1990, Geologic map of the Omak 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-12, 52 p., 1 pl.
- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.

**Summit** (419)

ALTERNATE NAMES		DISTRICT	COUNTY
Alice		Palmer Mtn.	Okanogan
PRIMARY QUADRANGLE	SCALE	1/2° x 1° QUAD	1° x 2° QUAD
Enterprise	1:24,000	Oroville	Okanogan
LATITUDE	LONGITUDE	SECTION, TOWNSHIP, AND RANGE	
48° 50' 48.73" N	119° 36' 27.75" W	SE 1/4 sec. 30, 39N, 26E	
LOCATION:			
HOST ROCK: NAME		LITHOLOGY	AGE
Palmer Mountain Greenstone		greenstone metagabbro	Permian - Triassic
COMMODITIES	ORE MINERALS	NON-ORE MINERALS	
Au	gold	quartz	
Ag	galena		
Pb	chalcopyrite		
Cu	tetrahedrite		
	malachite		
	azurite		
DEPOSIT TYPE		MINERALIZATION AGE	
vein			

**PRODUCTION:** Concentrates recovered from 85 tons of ore in 1937 were worth \$1,538; total production in 1937 was about \$1,900 (Hunting, 1956, p. 149).

**TECTONIC SETTING:** The Palmer Mountain Greenstone was deposited along a convergent continental margin proximal to an island arc.

**ORE CONTROLS:** The deposit consists of a 1.5-6-ft-wide, nearly vertical vein trending N55W in aphanitic greenstone and medium-grained metagabbro. Chlorite alteration is present near the vein (Rinehart and Fox, 1972, p. 78).

**GEOLOGIC SETTING:** The vein is in greenstone and metagabbro of the Permian-Triassic Palmer Mountain Greenstone (Rinehart and Fox, 1972).

**COMMENTS:** The deposit was developed by three shafts totaling 420 ft and five adits totaling 1,200 ft (Hunting, 1956, p. 149).

**REFERENCES**

- Hunting, M. T., 1956, Inventory of Washington minerals—Part II, Metallic minerals: Washington Division of Mines and Geology Bulletin 37, v. 1, 428 p.; v. 2, 67 p.
- Rinehart, C. D.; Fox, K. F., Jr., 1972, Geology and mineral deposits of the Loomis quadrangle, Okanogan County, Washington: Washington Division of Mines and Geology Bulletin 64, 124 p., 3 pl.