

ENVIRONMENTAL ASSESSMENT OF PROPOSED GEODUCK HARVEST  
ALONG THE NORTHERN SHORELINE OF HARTSTENE ISLAND  
AT THE DOUGALL NORTH GEODUCK TRACT (#15490)

---

**Commercial geoduck harvest is jointly managed by the Washington Departments of Fish and Wildlife (WDFW) and Natural Resources (DNR) and is coordinated with treaty tribes through annual harvest management plans. Harvest is conducted by divers from subtidal beds between the -18 foot and -70 foot water depth contours (corrected to mean lower low water, hereafter MLLW). Harvest is rotated throughout Puget Sound in six geoduck management regions. The fishery, its management, and its environmental impacts are presented in the Puget Sound Commercial Geoduck Fishery Management Plan and Final Supplemental Environmental Impact Statement (WDFW & DNR, May 2001). The proposed harvest along the northern shoreline of Hartstene Island is described below.**

---

Proposed Harvest Dates: 2022 - 2023

Tract name: Dougall North tract (Tract #15490)

Description: (Figure 1, Tract vicinity map)

The Dougall North geoduck tract is a subtidal area with a proposed harvest area of approximately 56 acres (Table 1) along the northern point of Hartstene Island the South Puget Sound Geoduck Management Region (Figure 1). The tract is adjacent to and shares a common boundary with the Dougall Point tract to the east and is bounded by a DOH unclassified area to the west. The commercial tract area lies between the -18 and -70 foot (MLLW) water depth contours.

The tract harvest area is bounded starting from a Control Point (CP) on the -18 foot (MLLW) water depth contour in the easterly portion of the tract at 47°18.057' N. latitude, 122°50.561' W. longitude (CP 1); westerly along the -18 foot (MLLW) water depth contour to a point at 47°18.077' N. latitude, 122°51.130' W. longitude (CP 2); then northerly to a point on the -70 foot (MLLW) water depth contour at 47°18.128' N. latitude, 122°51.130' W. longitude (CP 3); then easterly along the -70 foot (MLLW) water depth contour to a point at 47°17.874' N. latitude, 122°50.310' W. longitude (CP 4); then northwesterly to the point of origin (Figure 2).

This estimate of the tract boundary is made using Geographic Information System (GIS) data layers that were generated from NOAA soundings. All contours are corrected to mean lower low water (MLLW). The shoreline data is from DNR, digitized at 1:24,000 scale in 1999. The -70 foot (MLLW) water depth contour is used for the deep water boundary, and the shallow water boundary is defined by the -18 foot contour (MLLW). The latitude and longitude positions are reported in decimal minutes to the closest thousandth of a minute. Corner latitude and longitude positions are generated using GIS, and have not been field verified to determine consistency with area estimates, landmark alignments, or water depth contours. The delineation of the tract boundary will be field

verified by DNR prior to any geoduck harvest. Any variance to the stated boundary will be coordinated between WDFW and DNR prior to geoduck harvest.

#### Substrate:

Geoducks are found in a wide variety of sediments ranging from soft mud to gravel. The most common sediments where geoducks are harvested are sand with varying amounts of mud and/or gravel. The specific sediment type of a bed is primarily determined by water current velocity. Coarse sediments are generally found in areas of fast currents and finer (muddier) sediments in areas of weak currents. The major impact of harvest will be the creation of small holes where the geoducks are removed. The holes fill in within a few days to several weeks and have no long-term effects. The substrate holes refill in areas with strong water currents much faster than in areas with weak water currents. Water currents tend to be relatively mild in the vicinity of the Dougall North tract. Currents reach an estimated average maximum flood velocity of 0.6 knots and an average ebb velocity of 0.7 knots (Tides and Currents software; station #1911; Pickering Passage, north end).

Substrates types vary greatly across this tract (subsurface substrates from dig samples found in Table 2) with sand being the predominant surface substrate type on 36 out of 46 transects. Mud was the dominant substrate on 10 transects. Other substrate types observed were pea gravel, shell, and cobble (Table 3, Figure 3).

#### Water Quality:

Water conditions in this area are affected by the convergence of currents from Pickering Passage and Case Inlet. The following data on water quality has been provided by the Washington Department of Ecology (DOE) for the Puget Sound Main Basin: Case Inlet; For 2010 and 2011 (most recent complete data years available) at water depths between -18 to -70 feet, the mean reported dissolved oxygen concentration is 8.9 mg/l with a range from 6.3 to 13.2 mg/l. The mean salinity at this station was 28.3 ppt with a range from 27.2 to 29.4 psu. The mean water temperature at this station was 10.7°C with a range from 6.9 to 16.0 °C.

This area is classified as “Approved” by the Washington Department of Health (DOH) for commercial shellfish harvest. DNR will verify the health status of the Dougall North tract prior to any state sanctioned geoduck harvest.

Biota:

Geoduck:

The Dougall North geoduck tract is approximately 56 acres and currently contains an estimated 558,171 pounds of geoducks (Table 1). The geoduck biomass estimate at this tract is based on a 2016 Squaxin Tribe survey resulting in a pre-fishing biomass of 1,066,594 lbs. A total of 508,423 pounds of geoduck have been landed on this tract to date. On all 6 dig stations (n=65 geoducks), geoducks were considered commercial quality (Table 2). Geoduck dig station difficulty ratings ranged from “very easy” to “difficult” to dig. Factors contributing to digging difficulty on stations #12-1, and #11-3 were lack of abundance, and the presence of gravel and shell in the substrate.

The geoduck density on this tract is moderate, currently averaging 0.11 geoducks/sq.ft. compared to a Puget Sound average density of about 0.16 geoducks/sq.ft. The geoducks at the Dougall North tract are average with an average weight of 2.04 pounds compared to the Puget Sound average geoduck weight of 2.1 pounds. The lowest average whole weight is 1.63 pounds per geoduck at station #8-1 and the highest average whole weight is 2.57 pounds per geoduck at station #12-1 (Table 4). Squaxin Tribe transect line start and end positions are listed in Table 5.

Geoducks are managed for long term sustainable harvest. No more than 2.7% of the fishable stocks are harvested (total fishing mortality) each year in each management region throughout Puget Sound. The fishable portion of the total Puget Sound population includes geoducks that are found in water deeper than -18 feet and shallower than -70 feet (corrected to mean lower low water (MLLW)). Other geoducks which are not harvestable are found inshore and offshore of the harvest areas. Observations in south Puget Sound show that major geoduck populations continue to depths of 360 feet. Additional geoducks exist in polluted areas and are also unavailable for harvest, but continue to spawn and contribute to the total population.

The low rate of harvest is due to geoduck's low rate of natural recruitment. WDFW has studied the regeneration rate of geoducks on certain tracts throughout Puget Sound. The estimated average time to regenerate a tract to its original density, after removal of 65 percent of the geoducks, is 55 years. The recovery time for the Dougall North tract is unknown. The research to empirically analyze tract recovery rates is continuing.

## Fish:

Geoduck beds are generally devoid of rocky outcroppings and other relief features that attract and support many fish species, such as rockfish and lingcod. The bottoms are relatively flat and composed of soft sediments which provide few attachments for macroalgae, which also is associated with rockfish and lingcod. The fish observed during the 2017 WDFW Supplemental survey of the Dougall North tract were various species of flatfish (rock sole sand dabs, and starry flounders), buffalo sculpin and unspecified sculpins, and unspecified fish (Table 6).

WDFW marine fish managers were asked of their concerns of any possible impacts that geoduck fishing would have on groundfish and baitfish.. Greg Bargmann of WDFW stated that geoduck fishing would have no long-term detrimental impacts and may have some short-term benefits to flatfish populations by increasing the availability of food. Dan Penttila of the WDFW Fish Management Program recommended that eelgrass beds within the harvest tract should be preserved for any spawning herring. During the 2017 Squaxin Tribe eelgrass survey, no eelgrass was observed along this tract below a depth of -16 feet (MLLW). The Dougall North nearshore tract boundary will be along the -18 foot (MLLW) water depth contour to provide a vertical buffer between eelgrass beds and geoduck harvest.

There are no Pacific herring spawning grounds documented along the northern shoreline of Hartstene Island in the vicinity of the Dougall North tract (Figure 4). However, a herring pre-spawner holding area has been identified off the northeastern shoreline of Hartstene Island. With a horizontal separation from known herring fish spawning sites, a nearshore geoduck harvest restriction of -18 feet or deeper, and lack of eelgrass beds within the tract, geoduck harvest on the Dougall North tract should have no detrimental impacts on herring spawning.

There is no sand lance spawning documented along the northern shoreline Hartstene Island, with the nearest spawning documented near Jarrell Cove (Figure 4). Sand lance populations are widespread within Puget Sound, the Strait of Juan de Fuca and the coastal estuaries of Washington. They are most commonly noted in areas such as the eastern Strait and Admiralty Inlet. However, WDFW plankton surveys and ongoing exploratory spawning habitat surveys suggest that there are very few if any bays and inlets in the Puget Sound basin that will not be found to support sand lance spawning activity. Sand lance spawning occurs at tidal elevations ranging from +5 feet to about the mean higher high water line. After deposition, sand lance eggs may be scattered over a wider range of the intertidal zone by wave action. The incubation period is about four weeks. Sand lances are an important part of the trophic link between zooplanktons and larger predators

in the local marine food webs. Like all forage fish, sand lances are a significant component in the diet of many economically important resources in Washington. On average, 35 percent of juvenile salmon diets are comprised of sand lance. Sand lances are particularly important to juvenile Chinook salmon, where 60 percent of their diet is comprised of sand lance. Other economically important species, such as Pacific cod (*Gadus macrocephalus*), Pacific hake (*Merluccius productus*) and dogfish (*Squalus acanthias*) feed heavily on juvenile and adult sand lance. There is substantial vertical separation between sand lance spawning (+5 feet to mean higher high water) and geoduck harvest activity (-18 ft. to -70 ft., MLLW). Geoduck harvest on the Dougall North tract should have no detrimental impacts on sand lance spawning.

There are areas of surf smelt spawning habitat that have been identified near the Dougall North tract (Figure 4). Surf smelt deposit adhesive, semi-transparent eggs on beaches that have a specific mixture of coarse sand and pea gravel. Inside Puget Sound, surf smelt spawning is thought to be associated with freshwater seepage, where the water keeps the spawning gravel moist. Eggs are deposited near the water's edge in water a few inches deep, around the time of the high-water slack. There is substantial vertical separation between surf smelt spawning (slack high tide) and geoduck harvest activity (-18 to -70 feet, MLLW). Geoduck harvest on the Dougall North tract should have no detrimental impacts on surf smelt spawning.

NOAA Fisheries Service announced on April 27, 2010, that it was listing canary and yelloweye rockfish as “threatened” and bocaccio as “endangered” under ESA (federal Endangered Species Act). The listings became effective on July 27, 2010. Historic high levels of fishing and water quality are cited as reasons that these rockfish populations are in peril and have been slow to recover. On January 23, 2017; canary rockfish were delisted based on newly obtained samples and genetic analysis (Federal Register 82 FR 7711). Geoduck fishery managers are tracking this process and will take actions necessary to reduce the risk of “take” of any listed rockfish species that could potentially result from geoduck harvest activity.

Two salmon populations, Puget Sound Chinook salmon and Hood Canal summer run chum salmon, were listed by the National Marine Fisheries Service on March 16, 1999, as threatened species under the federal Endangered Species Act. Critical habitat for summer run chum salmon populations includes all marine, estuarine, and river reaches accessible to the listed chum salmon between Dungeness Bay and Hood Canal and within Hood Canal. The timing for summer run chum spawning is early September to mid-October. Out-migration of juveniles has been observed in Hood Canal during February and March, though out-migration may be as late as mid-April. The Dougall North tract is outside of the critical habitat range for Hood Canal summer run chum salmon.

Critical habitat for Puget Sound Chinook salmon includes all marine, estuarine and river reaches accessible to listed Chinook salmon in Puget Sound. WDFW recognizes 27 distinct stocks of Chinook salmon: 8 spring-run, 4 summer-run, and 15 summer/fall and fall-run stocks. The existence of an additional five spring-run stocks is in dispute. The majority of Puget Sound Chinook salmon emigrate to the ocean as subyearlings.

The (horizontal) geographic separation of this tract from known spawning tributaries and vertical separation of geoduck harvest (deeper and seaward of the -18 ft. MLLW contour) from juvenile salmon rearing areas and migration corridors (upper few meters of the water column) reduces or eliminates potential impacts to salmon populations. Charles Simenstad of the University of Washington School of Fisheries stated that the exclusionary principle of not allowing leasing/harvesting in water shallower than -18 ft. MLLW, within two vertical feet of the lower eelgrass margin, and within any regions of documented herring or forage fish spawning should, under most conditions, remove the influences of harvest induced sediment plumes on migrating salmon. Geoduck harvest should have no impact on salmon populations.

On May 7, 2007, NOAA Fisheries Service announced listing of Puget Sound steelhead as “threatened” under ESA. This listing includes more than 50 stocks of summer- and winter-run steelhead. Steelhead share many of the same waters as Puget Sound Chinook salmon, which are already protected by ESA, and will benefit from shared conservation strategies. There are no identified streams or rivers in the vicinity of the Dougall North tract that support steelhead stocks. The horizontal separation between tributaries that support steelhead runs and the Dougall North tract will assure that geoduck harvest will likely have no impact on steelhead populations.

Green sturgeon have undergone ESA review in recent years, due to depressed populations. NOAA Fisheries Service produced an updated status review on February 22, 2005, and reaffirmed that the northern green sturgeon Distinct Population Segment (DPS) warranted listing as a Species of Concern, however proposed that the Southern DPS should be listed as Threatened under the ESA. NMFS published a final rule on April 7, 2006, listing the Southern DPS as threatened [pdf] (71 FR 17757), which took effect June 6, 2006. The green sturgeon critical habitat proposed for designation includes the outer coast of Washington within 110 meters (m) depth (including Willapa Bay and Grays Harbor) to Cape Flattery and the Strait of Juan de Fuca to its United States boundary. Puget Sound proper has been excluded from this critical habitat designation. The Dougall North geoduck tract is outside of the critical habitat range of green sturgeon and geoduck harvest at this location will have no adverse effects on ESA recovery efforts for green sturgeon populations.

Invertebrates:

Many different kinds of invertebrates were observed which are frequently found on geoduck beds during the 2017 WDFW Supplemental survey, including anemones, bivalves, cnidarians, crab, cucumbers, gastropods, nudibranchs, sea stars, shrimp and annelid worms (Table 6). Geoduck harvest has not been shown to have long-term adverse effects on these invertebrates. Geoduck harvest can depress some benthic invertebrates, however most of these animals recover within one year.

There is on-going interest from recreational and commercial crab fishers about interactions between geoduck harvest activity and Dungeness crab populations. Dungeness crab were observed on only one transect during the Dougall North 2017 WDFW Supplementary survey. Dr. Dave Armstrong at the University of Washington has determined that Dungeness crab utilize Puget Sound bottoms from the +1 foot level out to the -330 foot level. The California Department of Fish and Wildlife suggest that coastal Dungeness crab can be found in waters as deep as 750 feet. Jensen (2014) and WDFW information (personal comm. WDFW Biologist Don Velasquez, 7/23/15) confirm a similar vertical distribution in Puget Sound, though the highest densities are found between the 0 to 360 foot water depth contours.

To determine the potential impacts to Dungeness crab, the percentage of substrate disturbed during fishing was calculated and compared to the entire crab habitat within the tract and shoreward of the tract to the +1 foot level and seaward out to mid-channel (Figure 5, Potential crab habitat map). The entire crab habitat along this tract is approximately 671 acres. There were about 523,365 harvestable geoducks on this tract, from the 2016 survey estimate. With a minimum harvest level of 65 percent, the total number harvested would be 340,187 geoducks. Approximately 1.18 square feet of substrate is disturbed for every geoduck harvested, so  $340,187 \times 1.18 = 401,421$  square feet of substrate. This equals about 9.22 acres. This is about 1.4 percent of the total available crab habitat in the vicinity of this tract.

WDFW and DNR have studied the effects of geoduck harvest on the population of Dungeness crab at Thorndyke Bay in Hood Canal. The results of 4.6 years of study have shown no adverse effects on crab populations due to geoduck fishing. Based on historic low abundance of Dungeness crab in the area of the Dougall North tract, the low amount of disturbance, and the lack of effects observed at the Thorndyke Bay study, we conclude that any effects on Dungeness crab populations will be very minor, if they occur at all.

Aquatic Plants:

Large attached aquatic plants are not generally found in geoduck beds in large quantities. Light restriction often limits plant growth to areas shallower than where most geoduck harvest occurs. Aquatic plants observed during the 2017 WDFW Supplemental survey of the Dougall North tract include (Table 7) laminarian algae; *Ulva* (sea lettuce); and small foliose red algae.

John Boettner and Tim Flint, from the WDFW Habitat Division, have stated that as long as geoduck fishing was restricted to seaward of the eelgrass beds, they have no concerns about the fishing. This was confirmed by WDFW Habitat Division who stated that the existing conditions in the fishery SEIS are sufficient to protect fish and wildlife habitat and natural resources. The shallow boundary of geoduck harvest is set at least two vertical feet seaward of the deepest eelgrass to protect all eelgrass from harvest activities. An eelgrass survey was done by the Squaxin Tribe in 2017 and no eelgrass was documented below a depth of -16 feet (MLLW). The shoreward boundary of this tract will be no shallower than the -18 foot water depth contour (MLLW), which will provide a vertical buffer of at least two vertical feet between any eelgrass beds in the vicinity of the tract and geoduck harvest activity.

#### Marine Mammals:

Several species of marine mammals, including seals, sea lions, and river otters may be observed in the vicinity of this geoduck tract. Killer whales (*Orcinus orca*) were frequently observed in the vicinity of this tract in 2015 (B. Sizemore personal observation and reported at <http://www.thurstontalk.com/2015/02/06/orcas-visiting-olympia/>). The Southern Resident stock of killer whales resides mainly in the San Juan Islands throughout spring and summer, but incursions south into Puget Sound occur more frequently during winter months (Brent Norberg, NOAA, pers. comm. 5/15/06). The Southern Resident stock of killer whales was listed as “endangered” under the federal Endangered Species Act (ESA) by the National Marine Fisheries Service on November 15, 2005. This is in addition to the designation of this stock in May 2003 as “depleted” under the Marine Mammal Protection Act. More information and a draft conservation plan for this stock can be found at the NOAA website (<https://www.fisheries.noaa.gov/action/listing-southern-resident-killer-whale-under-esa>).

Hand pick shellfish fisheries, like geoduck harvesting, are considered Category III under the Marine Mammal Authorization Program for Commercial Fisheries. This means that there is a “rare or remote” likelihood of marine mammal “take,” (Brent Norberg, NOAA, pers. comm. 5/15/06). Precautions should be taken by commercial divers, when marine mammals are in the area, to be aware of marine mammal movements and behavior to eliminate the remote risk of entanglement with diver hoses and lines.



#### Birds:

A variety of marine birds are common in Puget Sound and in the general vicinity of this tract. The most significant of these are guillemots, murrelets, grebes, loons, scoters, dabbling ducks, black brant, mergansers, buffleheads, cormorants, gulls, and terns. Blue heron, bald eagles, and osprey are regularly observed. Geoduck harvest does not appear to have any significant effect on these birds or their use of the waters where harvest occurs. A study by DNR and the WDFW was conducted at northern Hood Canal to learn the effects of geoduck fishing on bald eagles (Watson et al., 1995). A significant conclusion of this study is that geoduck clam harvest is unlikely to have any adverse impacts on bald eagle productivity.

#### Other uses:

##### Adjacent Upland Use:

The upland property along the Dougall Point tract has an Environmental Designation of Rural. To minimize possible disturbance to adjacent residents, harvest vessels are not allowed within 200 yards of the ordinary high tide line (OHT) or shallower than -18 feet (MLLW) whichever is farther seaward. Harvest is only allowed during daylight hours, and no harvest is allowed on Saturdays, Sundays, or state holidays.

The only visual effect of harvest is the presence of the harvest vessels on the tract. These 35-40 foot boats are anchored during harvest and all harvest is conducted out of sight by divers. Noise from the boats, compressors and pumps may not exceed 50 dBA measured 200 yards from the noise source, 5 dBA below the state noise standard.

##### Fishing:

This area is not a prime sportfishing area, however, some recreational salmon fishing could occur seasonally in proximity to the geoduck bed. The WDFW Sport Fishing Rules pamphlet describes additional seasons, size limits, daily limits, specific closed areas, and additional rules for salmon and other marine fish species. A few small-scale commercial fisheries may take place in the area. The fishing, which does occur, should not create any problems for the geoduck harvesting effort in the area.

Geoduck fishing on this tract is managed in coordination with the southern Puget Sound treaty tribes through annual state/tribal harvest management plans. The non-Indian geoduck fishery should not be in conflict with any concurrent tribal fisheries.

Navigation:


Pickering Passage and Case Inlet experience a moderate amount of recreational vessel traffic, with seasonal fluctuations. The Dougall North tract is not within a major traffic lane and areas close to shore are used primarily by small shoal draft boats. Geoduck harvesting at this site should not result in any significant navigational conflicts. The Department of Natural Resources will notify the local boating community prior to harvests.

Summary:

Commercial geoduck harvest is proposed for the Dougall North geoduck tract located along the northern shoreline of Hartstene Island. The pre-fishing survey of this tract was done by the Squaxin Tribe in 2016, and the supplemental survey was done by WDFW in 2017. The tract biomass estimate is based on the 2016 survey and recent geoduck landings. The anticipated environmental impacts of this harvest are within the range of conditions discussed in the Final Supplemental Environmental Impact Statement (2001) for the commercial geoduck clam fishery. To reduce possible impacts to baitfish and eelgrass, harvest will be deeper and seaward of the -18 foot (MLLW) water depth contour. No significant impacts are expected from this harvest.


File: 220822\_DougallNorth\_#15490\_EA.doc

Figure 1. Vicinity Map,  
Dougall North Commercial Geoduck Tract #15490



1:80,000  
1 inch = 1.26 miles

Data Sources:  
Projection for data is GCS\_Washington Geographic System 1984,  
Units: Meters. Coastline layer is from DNR, 1: 24,000 scale, created  
09-20-99. Contours are from NOAA soundings.




Map Date: August 25, 2017  
Map Author: O. Working  
File: Data\Ocean\Geoduck

# Figure 2. Control Points Map, Dougall North Commercial Geoduck Tract #15490

Control Point	Latitude	Longitude
1	47° 18.057 122° 50.561'	
2	47° 18.077 122° 51.130'	
3	47° 18.128 122° 51.130'	
4	47° 17.874 122° 50.310'	

Coordinates are in WGS84;  
Degrees, Decimal Minutes



**Legend**


- Control Points
- 18 ft. (MLLW)
- 70 ft. (MLLW)
- 200 Yard line
- Tract Within 200 Yard Line= 13 Acres
- Dougall North Geoduck Tract= 56 Acres

1:8,000  
1 inch = 0.13 miles

Data Sources:  
Projection for data is GCS\_Washington Geographic System 1984,  
Units: Meters. Coastline layer is from DNR, 1: 24,000 scale, created  
09-20-99. Contours are from NOAA soundings.

Map Date: January 19, 2022  
Map Author: O. Working  
File: Data/Ocean/Geoduck

# Figure 3. Transect and Dig Station Map, Dougall North Commercial Geoduck Tract #15490



**Legend**

- 2016 Squaxin Tribe Transect Points
- 2016 Squaxin Tribe Transect Lines
- ▲ 2016 Squaxin Tribe Dig Stations
- 2017 WDFW Mapped Contours**
- 18 ft. (MLLW)
- 70 ft. (MLLW)
- Dougall North Geoduck Tract= 56 Acres

1:10,000  
1 inch = 0.16 miles

Data Sources:  
Projection for data is GCS\_Washington Geographic System 1984,  
Units: Meters. Coastline layer is from DNR, 1: 24,000 scale, created  
09-20-99. Contours are from NOAA soundings.





0 0.05 0.1 0.2 0.3 Miles

Washington  
Department of  
**FISH and  
WILDLIFE**

Map Date: August 25, 2017  
Map Author: O. Working  
File: Data\Ocean\Geoduck




# Figure 4. Fish Spawning Areas Near the Dougall North Commercial Geoduck Tract #15490



# Figure 5. Dungeness Crab Habitat Map, Dougall North Commercial Geoduck Tract #15490

Note: Potential Dungeness crab habitat may extend to the -330 ft. (MLLW) depth contour. To make an estimate of potential habitat, the boundaries were drawn starting at shoreline, following the geoduck tract side boundaries to mid channel, as the water depth does not reach -330 ft. (MLLW) in this area.



1:40,000  
1 inch = 0.63 miles

Data Sources:  
Projection for data is GCS\_Washington Geographic System 1984,  
Units: Meters. Coastline layer is from DNR, 1: 24,000 scale, created  
09-20-99. Contours are from NOAA soundings.



Map Date: August 25, 2017  
Map Author: O. Working  
File: Data\Ocean\Geoduck

## EXPLANATION OF SURVEY DATA TABLES

The geoduck survey data for each tract is reported in seven computer-generated tables. These tables contain specific information gathered from transect and dig samples and diver observations. The following is an explanation of the headings and codes used in these tables.

### Tract Summary

This table is a general summary of survey information for the geoduck tract including estimates of *Tract Size* in acres, average geoduck *Density* in animals per sq.ft., *Total Tract Biomass* in pounds with statistical confidence, and *Total Number of Geoducks*. Mass estimators are reported in average values for *Whole Weight* and *Siphon Weight* in pounds. Geoduck siphon weights are also reported in *Siphon Weight as a percentage of Whole Weight*. Biomass estimates are adjusted for any harvest that may occur subsequent to the pre-fishing survey.

### Digging Difficulty

This table presents a station-by-station evaluation of the factors contributing to the difficulty of digging geoduck samples with a 5/8" inside nozzle diameter water jet. Codes for the overall subjective summary of the digging difficulty are given in the *Difficulty* column. An explanation of the codes for the dig difficulty follows:

<u>Code</u>	<u>Degree of Difficulty</u>	<u>Description</u>
0	Very Easy	Sediment conducive to quick harvest.
1	Easy	Significant barrier in substrate to inhibit digging.
2	Some difficulty	Substrate may be compact or contain gravel, shell or clay; most geoducks still easy to dig.
3	Difficult	Most geoducks were difficult to dig, but most attempts were successful.
4	Very Difficult	It was laborious to dig each geoduck. Unable to dig some geoducks.
5	Impossible	Divers could not remove geoducks from the substrate.

*Abundance* refers to the relative geoduck abundance; a zero (0) indicates that geoducks were very sparse, a one (1) indicates that they were moderately abundant and a two (2) indicates that they were very abundant. *Depth* refers to the depth that the geoducks were found in the substrate. A zero (0) indicates that they were shallow, a one (1) indicates that they were moderately deep and a two (2) indicates that they were very deep. The columns labeled *Compact*, *Gravel*, *Shell*, *Turbidity* and *Algae* refer to factors that contribute to digging difficulty by interfering with the digging process. A zero (0) in one of these columns indicates that the factor was not a problem, a one (1) indicates that the



factor caused moderate difficulty and a two (2) indicates that the factor caused a significant amount of difficulty when digging. *Compact* refers to the compact or sticky nature of a muddy substrate. *Gravel* and *Shell* refer to the difficulty caused by these substrate types. *Turbidity* refers to the turbidity within the water near the dig hole caused by the digging activity. High turbidity makes it difficult to find the geoduck siphon shows. The difficulty of digging associated with turbidity varies with the amount of tidal current present. Therefore, the turbidity rating refers only to the conditions occurring when the sample was collected. *Algae* refers to algal cover, which also makes it difficult for the diver to find geoduck siphon shows. Because algal cover varies seasonally, this value only applies to the conditions when the sample was collected. The *Commercial* column gives a subjective assessment of whether or not it would be feasible to harvest geoducks on a commercial basis at the given station.

### **Transect Water Depths, Geoduck Densities and Substrate Observations**

This table reports findings for each transect. *Start Depth* and *End Depth* (corrected to MLLW) are given for each transect. *Geoduck Density* is reported as the average number of geoducks per square foot for each 900 square foot transect. *Substrate Type* and *Substrate Rating* refer to evaluations of the substrate surface. A two (2) rating indicates that the substrate type is predominant. A one (1) rating indicates the substrate type was present.

### **Geoduck Weights and Proportion Over 2 Pounds**

This table summarizes the size and quality of the geoducks at each of the stations where dig samples were collected. Weight values for any geoduck dig samples that were damaged during sampling to the extent that water loss occurred, are excluded from calculations. The *Number Dug* column lists the number of geoducks collected. The *Avg. Whole Weight (lbs.)* column gives the average sample weight of whole geoduck clams for each dig station. The *Avg. Siphon Weight (lbs.)* column gives the average weight of the siphons of the geoducks for each dig station. The percentage of geoducks greater than two pounds is given in the *% Greater than 2 lbs.* column.

### **Transect - Corrected Geoduck Count and Position Table**

This table reports the diver *Corrected Count*, the geoduck siphon *Show Factor* used to correct the count, and the *Latitude/Longitude* position of the start point of each survey transect. Raw (observed) siphon counts are “corrected” by dividing diver observed counts for each transect with a siphon “show” factor (See WDFW Tech. Report FPT00-01 for explanation of show factor) to estimate the sample population density. Transect positions are reported in degrees and decimal minutes to the thousandth of a minute, datum WGS84.

### **Most Common and Obvious Animals Observed**

This table summarizes the animals, other than geoducks, that were observed during the geoduck survey, and reports the total number of transects on which they were present (*# of Transects Where Observed*). This is qualitative presence/absence data only, and only animals that can be readily seen by divers at or near the surface of the substrate are noted. The *Group* designation allows for the organization of similar species together in the table.

Whenever possible, the scientific name of the animal is listed in *Taxonomer*, and a generally accepted *Common Name* is also listed. Many variables may make it difficult for divers to notice other animals on the tract, including but not limited to poor visibility, diver skill, animals fleeing the divers, animal size, or cryptic appearance or behavior (in crevasses or under rocks).

### **Most Common and Obvious Algae Observed**

This table summarizes marine algae observed during the geoduck survey, and reports the total number of transects on which they were seen (*# of Transects Where Observed*).

This is qualitative presence/absence data only, and only for macro algae, with the exception of diatoms. At high densities diatoms form a “layer” on or above the substrate surface that is readily visible and obvious to divers. Other types of phytoplankton are not sampled and are rarely noted. Whenever possible, the scientific name or a general taxonomic grouping of each plant is listed in *Taxonomer*.

Last Updated: April 14, 2020

S:\FP\FishMgmt\Geoduck\EnvironmentalAssessmentReports\Forms\EAcodesexplanation\_7tables.doc

**Table 1. GEODUCK TRACT SUMMARY**

Dougall North geoduck tract # 15490.

Tract Name	Dougall North
Tract Number	15490
Tract Size (acres) <sup>a</sup>	56
Density of geoducks/sq.ft. <sup>b</sup>	0.11
Total Tract Biomass (lbs.) <sup>b</sup>	558,171
Total Number of Geoducks on Tract <sup>b</sup>	273,888
Confidence Interval (%)	23.8%
Mean Geoduck Whole Weight (lbs.)	2.04
Mean Geoduck Siphon Weight (lbs.)	N/A*
Siphon Weight as a % of Whole Weight	N/A*
Number of Transect Stations	46
Number of Geoducks Weighed	65

<sup>a</sup> Tract area is between the -18 ft. and -70 ft. (MLLW) water depth contours

<sup>b</sup> Biomass is based on the 2016 Squaxin Tribe Pre-fishing survey biomass of 1,066,594 lbs. minus total harvest of 508,423 lbs. through August 22, 2022

\* No siphon weights taken during this survey

Generation Date: August 22, 2022  
Generated By: O. Working, WDFW  
File: S:\FP\FishMgmt\Geoduck\EAs\2022

Dougall North geoduck tract # 15490, 2016 Squaxin Tribe Pre-fishing geoduck survey

Dig Station	Difficulty (0-5)	Abundance (0-2)	Depth (0-2)	Compact (0-2)	Gravel (0-2)	Shell (0-2)	Turbidity (0-2)	Algae (0-2)	Commercial (Y/N)
DN8-1	0	2	0	0	0	0	0	0	y
DN12-1	3	2	0	0	2	2	0	0	y
DN11-3	3	0	0	0	2	0	0	0	y
DN6-2	1	2	0	0	0	0	0	0	y
DN2-4	0	2	0	0	1	0	1	0	y
DN3-3	0	2	0	0	1	1	0	0	y

Generation Date: August 22, 2022  
 Generated By: O. Working, WDFW  
 File: S\FP\FishMgmt\Geoduck\EAs\2022

**Table 3: TRANSECT WATER DEPTHS, GEODUCK DENSITIES, AND SUBSTRATE OBSERVATIONS**

Dougall North geoduck tract # 15490, 2016 Squaxin Tribe Pre-fishing geoduck survey

Transect	Start Depth (ft) <sup>a</sup>	End Depth (ft) <sup>a</sup>	Geoduck Density (no. / sq ft) <sup>b</sup>	Substrate <sup>c</sup>				
				mud	sand	peagravel	shell	cobble
DN2-1	18	51	0.1244		2			
DN2-2	51	57	0.0104		2			
DN2-3	57	64	0.0133		2		1	
DN2-4	64	63	0.0637	1	2		1	
DN2-5	63	65	0.0563	2			1	
DN2-6	65	48	0.0993	2				
DN2-7	48	22	0.0963	2				
DN3-1	19	33	0.2296		2			
DN3-2	33	39	0.3837		2			
DN3-3	39	70	0.2770		2			
DN4-2	32	45	0.1687	2				
DN4-3	45	49	0.2470	2				
DN4-4	49	63	0.2338	2				
DN4-5	63	68	0.0422	2				
DN4-6	68	66	0.0096	2				
DN4-7	66	45	0.0374		2			
DN4-8	45	21	0.0434		2			
DN5-3	39	51	0.4965		2			
DN5-4	51	63	0.4796		2			
DN5-5	63	70	0.5941		2			
DN6-1	26	34	0.0012	1	2			
DN6-2	34	44	0.0325	1	2			
DN6-3	44	63	0.4483		2			
DN6-4	63	61	0.1916		2			
DN7-1	53	51	0.0181		2			
DN7-2	51	58	0.2892		2			
DN7-3	58	68	0.4134		2			
DN7-4	68	58	0.1904		2			1
DN8-1	59	59	0.2266	1	2			
DN8-2	59	62	0.3169		2			
DN8-3	62	70	0.3917		2			
DN8-4	70	70	0.3688		2			
DN8-5	70	67	0.4387		2			
DN8-6	67	64	0.4326		2			
DN9-1	30	51	0.2756	1	2			1
DN9-2	51	52	0.0593	1	2			1
DN9-3	52	57	0.0415	2	1			1
DN9-4	57	70	0.0044	2	1			1
DN11-1	23	26	0.1393		2	1		
DN11-2	26	44	0.1274		2	1		
DN11-3	44	63	0.1793		2	1		1
DN12-1	18	55	0.3570		2			
DN12-2	55	67	0.4281		2			

**Table 3. Continued**

Transect	Start Depth (ft) <sup>a</sup>	End Depth (ft) <sup>a</sup>	Geoduck Density (no. / sq ft) <sup>b</sup>	Substrate <sup>c</sup>				
				mud	sand	peagravel	shell	cobble
DN12-3	67	54	0.3526	1	2			
DN12-4	64	42	0.3467	1	2			
DN12-5	42	27	0.0830		2	1		

<sup>a</sup> All depths are corrected to mean lower low water (MLLW)

<sup>b</sup> Densities were calculated using a daily siphon show factor or the default 0.75 show factor

<sup>c</sup> Substrate codes: 1 = present ; 2 = dominant

Generation Date: August 22, 2022  
 Generated By: O. Working, WDFW  
 File: S\FP\FishMgmt\Geoduck\EAs\2022

**Table 4: GEODUCK SIZE AND QUALITY**

Dougall North geoduck tract # 15490, 2016 Squaxin Tribe Pre-fishing geoduck survey

Dig Station	Number Dug	Avg. Whole Weight (lbs.)	Avg. Siphon Weight (lbs.)	% of geoducks on station greater than 2 lbs.
DN8-1	11	1.63	N/A*	18%
DN12-1	10	2.57	N/A*	80%
DN11-3	11	2.18	N/A*	70%
DN6-2	11	2.01	N/A*	55%
DN2-4	10	1.66	N/A*	20%
DN3-3	12	2.18	N/A*	58%

\* No siphon weights taken during this survey

Generation Date: August 22, 2022  
Generated By: O. Working, WDFW  
File: S:\FP\FishMgmt\Geoduck\EAs\2022

**Table 5: TRANSECT CORRECTED GEODUCK COUNT AND POSITION TABLE**

Dougall North geoduck tract # 15490, 2016 Squaxin Tribe Pre-fishing geoduck survey

Transect	Corrected Geoduck Count per 900 sq. ft. Transect	Geoduck Siphon Show Factor <sup>a</sup>	Latitude <sup>b</sup>	Longitude <sup>b</sup>
DN2-1	112	0.750	47 18.14	122 51.003
DN2-2	9	0.750		
DN2-3	12	0.750		
DN2-4	57	0.750	47 0.177	122 50.92
DN2-5	51	0.750	9E+05	
DN2-6	89	0.750	4E+05	
DN2-7	87	0.750	47 18.1	122 50.904
DN3-1	207	0.750	47 18.09	122 50.89
DN3-2	345	0.750		
DN3-3	249	0.750	47 18.16	122 50.846
DN4-2	152	0.922		
DN4-3	222	0.922		
DN4-4	210	0.922		
DN4-5	38	0.922	47 18.13	122 50.75
DN4-6	9	0.922		
DN4-7	34	0.922		
DN4-8	39	0.922	47 18.18	122 50.618
DN5-3	447	0.922		
DN5-4	432	0.922		
DN5-5	535	0.922	47 18.2	122 50.447
DN6-1	1	0.922	47 18.08	122 50.498
DN6-2	29	0.922		
DN6-3	403	0.922		
DN6-4	172	0.922	47 18.13	122 50.417
DN7-1	16	0.922	47 18.06	122 50.441
DN7-2	260	0.922		
DN7-3	372	0.922		
DN7-4	171	0.922	47 18.13	122 50.417
DN8-1	204	0.922	47 18.03	122 50.448
DN8-2	285	0.922		
DN8-3	352	0.922		
DN8-4	332	0.922	47 18.04	122 50.389
DN8-5	395	0.922		
DN8-6	389	0.922	47 18.01	122 50.42
DN9-1	248	0.750	47 17.97	122 50.35
DN9-2	53	0.750		
DN9-3	37	0.750		
DN9-4	4	0.750	47 17.98	122 50.217
DN11-1	125	0.750	47 18.07	122 50.657
DN11-2	115	0.750		
DN11-3	161	0.750	47 18.15	122 50.643
DN12-1	321	0.750	47 18.13	122 51.027
DN12-2	385	0.750		
DN12-3	317	0.750	47 18.15	122 51.049



**Table 5. Continued**

Transect	Corrected Geoduck Count per 900 sq. ft. Transect	Geoduck Siphon Show Factor <sup>a</sup>	Latitude <sup>b</sup>	Longitude <sup>b</sup>
DN12-4	312	0.750	47 18.04	122 51.095
DN12-5	75	0.750		

<sup>a</sup>: Densities were calculated using a daily siphon show factor or the default 0.75 show factor

<sup>b</sup>: Latitude and longitude are in WGS84 datum, degrees and decimal minutes

Generation Date: August 22, 2022

Generated By: O. Working, WDFW

File: S\FP\FishMgmt\Geoduck\EAs\2022

**Table 6: MOST COMMON AND OBVIOUS ANIMALS OBSERVED**

Dougall North geoduck tract # 15490, 2017 WDFW Supplemental geoduck survey

# of Transects where Observed	Group	Common Name	Taxonomer
10	ANEMONE	BURROWING ANEMONE	<i>Pachycerianthus fimbriatus</i>
6	ANEMONE	PLUMED ANEMONE	<i>Metridium</i> spp.
3	ANEMONE	STRIPED ANEMONE	<i>Urticina</i> spp.
1	ASCIDIAN	SESSILE TUNICATE	Unspecified Tunicate
12	BIVALVE	HORSE CLAM	<i>Tresus</i> spp.
4	CNIDARIA	SEA PEN	<i>Ptilosarcus gurneyi</i>
10	CNIDARIA	SEA WHIP	<i>Stylatula elongata</i>
4	CRAB	DECORATOR CRAB	<i>Oregonia gracilis</i>
1	CRAB	DUNGENESS CRAB	<i>Cancer magister</i>
11	CRAB	GRACEFUL CRAB	<i>Cancer gracilis</i>
9	CRAB	HERMIT CRAB	Unspecified hermit crab
5	CRAB	RED ROCK CRAB	<i>Cancer productus</i>
4	CUCUMBER	SEA CUCUMBER	<i>Parastichopus californicus</i>
1	FISH	BUFFALO SCULPIN	<i>Enophrys bison</i>
3	FISH	FISH	Unspecified Fish
1	FISH	ROCK SOLE	<i>Lepidopsetta bilineata</i>
11	FISH	SANDDAB	<i>Citharichthys</i> spp.
5	FISH	SCULPIN	Unspecified Cottidae
1	FISH	STARRY FLOUNDER	<i>Platichthys stellatus</i>
10	GASTROPOD	MOON SNAIL EGGS	<i>Polinices lewisii</i> egg case
4	NUDIBRANCH	ARMINA	<i>Armina californica</i>
4	NUDIBRANCH	DENDRONOTUS	<i>Dendronotus</i> spp.
5	NUDIBRANCH	ROSY TRITONIA	<i>Tritonia diomedea</i>
4	SEA STAR	FALSE OCHRE STAR	<i>Evasterias troschelli</i>
3	SEA STAR	LEATHER STAR	<i>Dermasterias imbricata</i>
1	SEA STAR	OCHRE STAR	<i>Pisaster ochraceus</i>
1	SEA STAR	SAND STAR	<i>Luidia foliolata</i>
1	SEA STAR	SUNFLOWER STAR	<i>Pycnopodia helianthoides</i>
2	SHRIMP	GHOST SHRIMP	Unspecified ghost shrimp
8	WORM	ROOTS	Chaetopterid polychaete tubes
11	WORM	SABELLID TUBE WORM	<i>Sabellid</i> spp.
2	WORM	TEREBELLID TUBE WORM	<i>Terebellid</i> spp.

Generation Date: August 22, 2022  
Generated By: O. Working, WDFW  
File: S\FP\FishMgmt\Geoduck\EAs\2022

**Table 7: MOST COMMON AND OBVIOUS ALGAE OBSERVED**

Dougall North geoduck tract # 15490, 2017 WDFW Supplemental

# of Transects Where Observed	Taxonomer
1	<i>Laminaria</i> spp.
8	<i>Ulva</i> spp.
10	Small red algae

Generation Date: August 22, 2022

Generated By: O. Working, WDFW

File: S\FP\FishMgmt\Geoduck\EAs\2022