

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Riparian Function Literature ~~Synthesis~~ Review and Annotated Bibliography

Prepared for the Riparian Scientific Advisory Group
(RSAG) of Washington State

Prepared by:

Benjamin Spei, Brandon Light, Mark Kimsey

March 2024

Commented [T(1): NOTE FOR ALL REVIEWERS: This round of review is specifically to correct factually inaccuracies. Please correct factual inaccuracies in-line in track changes (not as a comment). Provide these corrections by Thursday, January 30.

Commented [AM2]: I see your request for inline edits, but wanted to say that we agreed that this was to not be cited as a "synthesis" I believe the language that was agreed on or at least proposed was "review and annotated bibliography"?

Commented [AT3R2]: I have made that change in track changes here in the title. I also conducted a search for "synthesis" in the document and revised to "review" when referencing this document.

26	Table of Contents	
27	Background	2
28	Focal Questions	2
29	Methods	3
30	Results/Summary of Review	5
31	Discussion of findings relative to FPHCP objectives	0
32	Litter/Organic matter inputs/Nutrients.....	0
33	Large Wood (LW)/wood load/wood recruitment	9
34	Bank Stability and Sediment	19
35	Shade and stream temperature	28
36	Results/discussion by focal question	36
37	Focal Question 1.....	36
38	Focal Question 1a.....	47
39	Focal Question 1b	51
40	Focal Question 1c.....	53
41	Focal Question 1d	56
42	Focal Question 1e.....	57
43	Focal Question 2.....	57
44	Focal Question 3.....	64
45	Focal Question 4.....	77
46	Focal Question 5.....	78
47	Focal Question 6.....	83
48	Focal Question 7.....	87
49	References.....	89
50	Appendix I	99
51	Appendix II	130
52		
53		
54		
55		
56		
57		

58 **Background**

59 Washington State Forest Practices rules and management guidelines covered by the FPHCP
60 (Forest Practices Habitat Conservation Plan, 2006) are strongly influenced by the science of
61 riparian processes articulated in the FPHCP Environmental Impact Statement (EIS Chapter 6
62 References, Appendix A Regional Summaries, Appendix B Riparian Modeling, 2005). The EIS
63 references include the Forest Ecosystem Management Assessment Team (FEMAT) report,
64 “Forest Ecosystem Management: an ecological, economic, and social assessment. Section V:
65 Aquatic Ecosystem Assessment (1993).” Although the Forests and Fish Report and FPHCP and
66 the rules derived from it considered many sources, our scientific understanding of riparian
67 processes has evolved based on additional research that has been completed since then. More
68 recent science has affirmed some aspects of the then-current state of knowledge on riparian
69 processes and the effects of timber harvest on them. Still, some of the scientific conclusions are
70 changing. In addition, riparian management strategies have evolved to address resource
71 objectives. This ~~synthesis-review~~ will look at literature that has been completed since the
72 FEMAT and Forests and Fish report, and the FPHCP EIS. It will inform the Adaptive
73 Management Program (AMP) committees and the Forest Practices Board (FPB) regarding the
74 effects of forest harvest and other management practices on riparian functions and processes.

75 This review will follow a similar but modified format of the riparian literature review developed
76 by Schuett-Hames et al. (2015) for the Cooperative Monitoring Evaluation and Research
77 Committee (CMER) under the Westside Type F Prescription Effectiveness Monitoring project.
78 However, this review will not focus only on Type F (fish-bearing streams) but on the response of
79 riparian functions following harvest in all forests adjacent to rivers and streams. Priority will be
80 given to studies conducted in areas with similar habitat and landscape characteristics as those
81 found in the state of Washington. Further, information extracted from these studies will include
82 the experimental designs used, sampling programs, sampled covariates, the metrics used to
83 quantify covariates, and analytical methods.

84 We summarized the overall findings by key riparian function, and related physical processes, and
85 provide a ~~synthesis-review~~ to support recommendations for future research. The riparian
86 functions specified in the FPHCP include “large woody debris recruitment, sediment filtration,
87 stream bank stability, shade, litterfall and nutrients, in addition to other processes important to
88 riparian and aquatic systems.” (FPHCP, 2006).

89 This literature review ~~and synthesis~~ will address specific questions (listed below) and identify
90 appropriate variables and associated metrics that can be used to quantify and assess timber
91 harvest effects on the riparian functions.

92

93 **Focal Questions**

- 94 1. What are the effects of timber harvest intensities and extent on the riparian functions,
95 with an emphasis on the five key functions listed above, in comparison to conditions
96 before harvest?

- 97 a. What are the effects of thinning (intensity, extent) on the riparian functions, over
98 the short and long-term compared to untreated stands?
99 b. How do buffer widths and adjacent upland timber harvest prescriptions influence
100 impacts of riparian thinning treatments?
101 c. What are the effects of clearcut gaps in riparian stands (intensity, extent) on the
102 riparian functions, over the short and long-term, compared to untreated stands
103 d. How do buffer widths and upland timber harvest influence impacts of clearcut
104 gaps treatments?
105 e. What are the effects of any combinations of the above treatments?
106 2. How and to what degree do specific site conditions (e.g., topography, channel width and
107 orientation, riparian stand age and composition) influence the response of the riparian
108 functions?
109 3. What is the frequency of weather-related effects (e.g., windthrow, ice storms, excessive
110 heat, flood and drought events) on riparian areas? What are the weather-related effects
111 (positive and negative) on the riparian functions, and how are they distinguished from
112 harvest effects? How do these effects differ between treated and untreated riparian
113 forests?
114 4. How do various treatments within riparian buffers relate to forest health and resilience to
115 fire, disease, and other forest disturbances?
116 5. How do the functions provided by riparian stands change over time (e.g., large woody
117 debris recruitment from farther away from the stream)?
118 6. Are there feedback mechanisms (e.g., microclimate changes within the riparian buffer)
119 related to forest management that affect the recovery rates of riparian functions?
120 7. What major data gaps and uncertainties exist relative to effects of timber harvest (both
121 riparian and adjacent upland) on the riparian functions?

122 Methods

123 The riparian function literature ~~synthesis review~~ includes literature pertinent to the effects of
124 timber harvest, management, natural disturbances (e.g., fire, disease, insect infestation, etc.), and
125 channel geomorphology in riparian areas on the “five key riparian functions” as defined in the
126 Forest Practices Habitat Conservation Plan (FPHCP, 2006). Literature searches were primarily
127 conducted using the Web of Science and Google Scholar. Sources were also gathered via
128 personal communication with employees and members of the Washington State Department of
129 Natural Resources’ Cooperative Monitoring Evaluation and Research (CMER) scientific
130 advisory groups. Technical reports on the United States Forest Service website were also
131 investigated for their potential use. Finally, we also considered studies and manuscripts
132 unpublished in formal scientific journals available on ResearchGate and ProQuest, including
133 Ph.D. dissertations and master's theses. Papers returned from the keyword searches were initially
134 screened by title and abstract. Papers were deemed appropriate for inclusion if they fit 3 criteria:
135 (1) utilize experimental designs such as before-after-control-impact (BACI), after-control-impact
136 (ACI), before-after-impact (BAI), after-impact (AI), simulation modeling, or meta-analysis to
137 quantify the effect of riparian forest treatment, harvest, disturbance, site characteristics and
138 conditions, etc. on riparian functions with an emphasis on the five key functions. Observational

139 studies that that substituted space for time (e.g., difference between old-growth and young
 140 regenerating forests) were also included. (2) have been published or completed since the Forest
 141 and Fish report, i.e., 1999, (3) have been conducted in western North America including coastal
 142 Alaska, southern and coastal British Columbia, southern Alberta, the Pacific Northwest, the
 143 Intermountain West, and the Great Basin regions. Studies from outside these areas were included
 144 if they contained generalizable information about riparian functions (e.g., the relationship of
 145 canopy cover with shade and temperature).

146 A list of search terms was developed to capture any studies relevant to the topics of the seven
 147 focal questions (Table. 1). A master list of all returned study titles and abstracts from Web of
 148 Science was also compiled for further analysis of keyword popularity and combinations (Figure
 149 1).

150 Table 1. List of terms used in search of keywords and titles of literature sourced from Web of
 151 Science. Terms in **bold** were used in all searches. Terms were grouped by topic (e.g.,
 152 management, physiography, disturbance, etc.). Results show the number of publications returned
 153 for each combination of search terms.

Key Words/title	Results
(Riparian OR stream OR headwater Or Watershed) AND	
(Function OR sediment OR nutrient OR woody debris OR large wood OR LWD OR woody debris recruitment OR shade OR temperature OR light OR litter OR water quality OR diversity OR wood*) AND/OR	15,138
(Manag* OR harvest OR thin* OR forest* OR forest operation OR buffer OR buffer strips OR gap* OR treat* OR clearcut OR clearcut gap)	12,602
(Topograph* OR physiograph* OR channel width OR stream width OR bankfull width OR valley constraint OR morphology OR diversity OR distance to stream OR Parent material OR soil OR litholo* OR geolog*)	12,381
(Disturbance OR fire OR windthrow OR ice storms OR drought OR flood* OR resilience OR resistance OR microclimate OR site conditions)	12,725
(Climate)	12,588
(feedback OR long-term OR short-term OR time)	12,150
(Forest health OR recovery OR regeneration OR disease OR insect OR fung* OR patho*)	12,328
(Stand structure OR stand age OR composition OR density OR structure OR species OR species composition)	12,214

Total titles and abstracts searched, excluding duplicates	16,750
---	---------------

154

155 From the initial title and topic review of the 16,516 papers sourced in our search, we refined the
 156 list to 528 papers for consideration based on the 3 criteria listed above (e.g., utilize experimental
 157 design with results focusing on at least one of the five key functions; published after 1999; were
 158 conducted in western North America). From these 528 papers we further refined our list to 105
 159 articles based on information gleaned from the abstract, introduction and methods sections
 160 regarding study design and relevant geography. Of these 105 articles 91 provided information on
 161 at least one of the five key functions and were thoroughly read and used to develop an annotated
 162 bibliography (Appendix). The other 14 articles provided information and experimental results
 163 about fire frequency and fire behavior in riparian areas, or effects of fire on one of the five key
 164 functions. These 14 papers about fire were not included in the literature review but were
 165 reviewed and discussed in focal questions 3 and 7. Frequency of the top 8 keywords were
 166 represented in a histogram to express the popularity of topics in the literature since the year 2000
 167 (Figure 1). We organized our review of the relevant literature by (1) FPHCP objective and (2)
 168 focal question. A table was submitted along with this report that gives a more thorough
 169 description of details used to categorize publications in supplemental materials (supplemental
 170 table of references; S1).

171 Table 2. Frequency of keywords in the original 16,516 publications sourced from Web of Science


Keywords	Count
Water quality	1165
Streams	1004
Watershed	1000
Climate change	848
Watershed management	729
Riparian	652
Stream	604
Nitrogen	489

172

173 Results/Summary of Review

174 We conducted our review of the 72 relevant publications to (1) summarize the most current state
 175 of knowledge of how timber harvest affects riparian function and related processes with a focus
 176 on the five key riparian functions defined in the FPHCP, and (2) extract information that has the
 177 potential to provide answers to, or methods and experimental designs that could be used to
 178 answer the 7 focal questions. Our review focused primarily on peer-reviewed journal
 179 publications but included 3 CMER reports and 1 report from the United States Forest Service
 180 website. Of these 72 studies, 33 were conducted on headwater or non-fish-bearing streams, 16 on
 181 fish-bearing streams, and 23 on a combination of fish and non-fish-bearing streams or

182 hypothetical streams in a model simulation (Table 3.). Most of the studies reviewed were
 183 conducted in the Pacific Northwest region but several from just outside this region (British
 184 Columbia, Alberta, Idaho, Montana, Wyoming, Colorado) were also included (Figure 2.). Few
 185 studies could be found that quantify how riparian area treatments directly affect bank stability.
 186 Several CMER studies, however, have investigated the effects of riparian timber management on
 187 soil and streambank disturbance and erosion (Ehinger et al., 2021; McIntyre et al., 2018; Schuett-
 188 Hames et al. 2011). In these studies, soil/bank disturbance and erosion were further analyzed for
 189 their contribution to sediment export and delivery to streams. Because of this relationship
 190 between bank erosion and sediment delivery, bank stability is discussed and reviewed in the
 191 section with sediment. Further, because of the paucity of studies in the literature that provide
 192 experimental evidence of how riparian area treatments affect bank stability, studies that
 193 investigate bank stability or bank erosion based on other factors (e.g., vegetation type, vegetation
 194 coverage) have been included and reviewed in question 7. These studies are provided as
 195 recommendations for methods that could be used in an experimental design comparing changes
 196 in bank stability before and after treatment or between treated and untreated riparian stands.



197
 198 Figure 2. Locations where studies were conducted. References not listed include studies that
 199 sourced data from multiple locations.

Table 3. Characteristics of 72 relevant publications

Reference	Purpose	Sampling Period Study Duration	Sample size (n)	Function / process	Experiment type Study Type	Scale (local, regional, global)	State Location
Anderson & Meleason (2009)	The effects of buffer width in combination with thinned stands, patch openings, and unthinned stands on LWLWD and vegetation cover.	5-6 years	2	LWD, vegetation	BACI experiment	Local, 6 reaches, 2 transects per reach	Oregon
Anderson et al. (2007)	The effects of forest mgmt. on stream shade and stream temperature.	3-6 years	2	SHD	BACI experiment	Local, 6 reaches, 2 transects per reach	Oregon
Bahuguna et al. (2010)	The effect of riparian buffer width on windthrow and LWD recruitment.	8 years	3	LWD	BACI experiment	Local, streams within 1 watershed	British Columbia
Benda et al. (2016)	The effects of forest management on LW large woody debris LW recruitment	100 years in 5-year time steps (modelled)	1	LWD	Simulation modelling from field data	Local, Alsea watershed	Oregon
Bilby & Heffner (2016)	Combination of literature and field experiments to determine factors contributing to litter delivery to streams.	1- year experimental data	4 mature sites, 3 young sites	LIT	Mixed effects modelling from field experiments	Local, data on windspeed collected from Humphrey Creek	Washington
Bladon et al. (2016)	Effects of buffers vs. no buffers on stream temperature.	6 years	6	SHD, stream temperature	BACI experiment	Local, Alsea watershed	Oregon

Commented [AM4]: Column headers should be defined as needed. E.g., how can study duration be a range of years? And what is "sample size". For some entries seems to be the total number of sites but then for some there is a "disclaimer" of "unbalanced" and it is not clear what that means? For function/process, these should be clearly defined (e.g., LWD = large wood) - also, many are steering away from using LWD and using just LW (since "debris" can have a negative connotation - suggest change to LW). For experiment type define BACI table header.

Commented [bs5R4]: Adjusted

Commented [AM6]: Not all of these are experiments.

Commented [AM7]: Not clear what "local" is. Define in header.

Commented [bs8R7]: Okay, 3 categories added to header.

Commented [AM9]: I would think this belong sin the "experiment type" except that a model is not per se an experiment.

Commented [bs10R9]: Okay

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Bladon et al. (2018)	The effects of a variety of contemporary forest mgmt. prescriptions on small, headwater streams.	14 years	7	SHD	ACI experiment	Local, 3 watersheds: Alsea, Trask, and Hinkle watersheds	Oregon
Burton et al. (2016)	Instream wood loading at different buffer widths, basin geomorphologies, and harvest intensities.	15 years	6	LWD	BACI experiment	Regional, along Oregon coast and Cascade Range	Oregon
Bywater-Reyes et al. (2018)	Variability in suspended sediment yield over half-century.	60 years	10	SED	Modeling, regression analysis of historical data	Local, H.J. Andrews Experimental Watershed	Oregon
Bywater-Reyes et al. (2017)	Effect of forest mgmt. on stream sediment delivery.	6 years	10	SED	ACI experiment	Local, Trask River Watershed	Oregon
Chen et al. (2005)	Compares the LWLWD biomass between 4 different mgmt. strategies.	1 year+ year data collection, 4 disturbance histories	4-5	LIT, LWLWD, NUT	ACI experiment	Local, Okanagan Valley, Kelowna	British Columbia, Canada
Chen et al. (2006)	Assesses the amount, distribution, dynamics, and function of LWLWD in forest streams	2 years field data	35	LWLWD	ACI experiment	Local, Okanagan Valley, Kelowna	British Columbia, Canada
Cole & Newton (2013)	Effect of 3 different retention buffer prescriptions on stream temperature.	6-7 years	4	SHD	BACI experiment	Local, within a radius of 200 km of Corvallis	Oregon

Commented [AM6]: Not all of these are experiments.

Commented [bs11]: These changes were not specifically requested, but clarify Aimee's points above.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Deval et al. (2021)	Disturbance effects on stream chemistry.	13 years	7	NUT	BACI experiment	Local, Mica Creek Experimental Watershed	Idaho
Ehinger et al. (2021)	Effectiveness of riparian mgmt. in maintaining function in for non-fish-bearing headwater streams on incompetent lithologies.	9 years	11 (two treatments)	LW, NUT, SED, SHD, Stream Temperature	BACI experiment	Regional	southwestern Washington
Fox & Bolton (2007)	observational study that categorizes the effects of riparian site geomorphology on LWLWD recruitment.	1 year data collection, multiple age classes, coverts and disturbance histories	150	LWLWD	Descriptive, spatial modeling on historical data	Regional, Coastal, West and east Cascade Range of Washington State	Washington
Gomi et al. (2001)	LWLWD recruitment in the short and long-term under 5 different mgmt. strategies.	1 year, 4-year data collection, 5 management histories	3	LWLWD, SED	ACI experiment	Local, Maybeso Experimental Forests	Alaska
Gravelle & Link (2007)	The impacts of timber harvest practices on stream temperature.	13 years	3	SHD	BACI experiment	Local, Mica creek Experimental Watershed	Idaho
Gravelle et al. (2009)	The effects of contemporary forest practices on the chemical properties of headwater streams and downstream locations.	14 years	3	NUT, SED	BACI experiment	Local, Mica creek Experimental Watershed	Idaho

Commented [AM6]: Not all of these are experiments.

Commented [AM12]: Not clear why these are not in alphabetical order. Should be made so.

Commented [bs13R12]: Okay

Commented [bs14R12]: Alphabetized

Commented [AM15]: This is mixing approach with duration, which is added inconsistently.

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Groom et al. (2011a/2011b)	The effectiveness of new riparian management protocols in preserving stream side shade and in-stream temperature under Oregon forest practice rules.	7 years	Unbalanced (15 state-owned and 18 private-owned)	SHD, stream temperature	BACI experiment	Regional, Oregon Coast Range	Oregon
Groom et al. (2011b/2011a)	The efficacy effect of new riparian forest management protocols in preserving stream side shade and in-stream temperatures. temperature under Oregon forest practice rules	7 years	Unbalanced (15 state-owned and 18 private-owned)	SHD, stream temperature	BACI experiment	Regional, Oregon Coast Range	Oregon
Guenther et al. (2014)	Differences in surface/sub-surface variability as well as influences of partial retention harvesting on stream temperature.	2 years	3	SHD	BACI experiment	Local, Malcolm Knapp Research Forest	British Columbia, Canada
Hart et al. (2013)	What riparian forest characteristics influence litter input to streams.	2 years	5	LIT, NUT	ACI experiment	Local, 5 contiguous watersheds in Oregon Coast range	Oregon

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Hatten et al. (2018)	The effect of contemporary and historical forest harvesting practices on suspended stream sediment.	12 years	3	SED	ACI experiment	Local, Central Oregon Coast Range	Oregon
Hough-Snee et al. (2016)	Evaluates which riparian, geomorphic, and hydrologic attributes are most strongly correlated to instream wood loads.	2 years of data	7	LWLWD, SHD	Modeling, correlative analysis	Regional, interior Columbia River basin	Canada, Oregon, Washington, Idaho
Hunter & Quinn (2009)	How differences in stream geomorphology affect water temperature.	2 years of data	2	stream temperature	AI experiment	Local, Olympic Peninsula	Washington
Hyatt & Naiman (2001)	The depletion rate of LWLWD in streams by size and species.	1 year; of data collection; Dendrochronology to estimate up to 50 years.	4	LWLWD	AI	Local, Queets River	Washington
Jackson et al. (2001)	Effect of forest mgmt. on stream temperature, large woody debris, LW, and stream sediment, between clearcut, thinned, and buffered treatments.	2 years	unbalanced: 4-6	LWD, SED	BACI experiment	Local, northwestern Washington Coast Range	Washington

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period Study Duration	Sample size (n)	Function / process		Experiment type Study Type	Scale (local, regional, global)	State/Province
Jackson & Wohl (2015)	Instream wood loads and geomorphic effects between streams draining montane forests of different ages.	1 year of data	10 sites > 200 years old, 23 young sites <200 years old	LWL	WD	CI, regression analysis	Local, Arapaho and Roosevelt National Forests	Colorado
Jackson et al. (2001)	Effect of forest mgmt. on stream temperature, LW, and stream sediment, between clearcut, thinned, and buffered treatments.	2 years	unbalanced: 4 -6	LW	SED	BACI experiment	Local, northwestern Washington Coast Range	Washington
Janisch et al. (2012)	The response of stream temperature to forest harvest, testing differences in continuous vs. patch buffers.	4-5 years	unbalanced: 5-6	SHD	SHD	BACI experiment	Local, southwestern Washington Coast Range	Washington
Johnson & Jones (2000)	Short-term and long-term effects of forest harvest on stream temperatures.	Historical dataset 1959-1982	3	SHD		BACI experiment	Local, H.J. Andrews Experimental Watershed	Oregon
Karwan et al. (2007)	Effects of timber harvest on suspended sediments in streams following timber harvest.	3 years	2	SED		BACI experiment	Local, Mica creek Experimental Watershed	Idaho

Reference	Purpose	Sampling Period Study Duration	Sample size (n)	Function / process	Experiment type Study Type	Scale (local, regional, global)	State/Province
Kaylor et al. (2017)	Examines the effects of riparian forest harvest and varying stages of stand recovery on light availability 50-60 years post treatment.	1 year data collection, 50 - 60 years post treatment	14	SHD	AI	Local, H.J. Andrews Experimental Watershed	Oregon
Kibler et al. (2013)	Examined the effects of contemporary forest practices on warm-season stream temperature regimes in headwater streams.	3.5 years	8	SHD	BACI experiment	Local, Hinkle Creek	Oregon
Kiffney & Richardson. (2010)	Evaluates the effects of forest mgmt. on organic matter/ litterfall recruitment.	8 years	Unbalanced: 2-3	LIT	ACI experiment	Local, southwestern British Columbia	British Columbia, Canada
Liquori (2006)	Examines differences in post-harvest ecological and geomorphic processes in buffered forest sites	1 year data collection	Unbalanced: 4-9	Other processes, disturbance post-harvest	AI	Local, managed tree farm in Cascade Mountains of western Washington	Washington
Litschert & MacDonald (2009)	Assessed streamside management zones to understand characteristics of the sediment delivery pathways following upland harvest.	1 year data collection	200	SED	AI	Regional, National Forests in the Sierra and Cascade mountains.	California

Commented [AM6]: Not all of these are experiments.

Commented [bs16]: Changes made in response to Aimee's comment

Reference	Purpose	Sampling Period Study Duration	Sample size (n)	Function / process	Experiment type Study Type	Scale (local, regional, global)	State/Province
Macdonald et al. (2003b) (2003a)	Examined the effects of three different retention harvesting prescriptions on stream temperature, suspended sediment concentrations.	76 years	52	SHD SED	ACI BACI experiment	Local, Baptiste and Galuski watersheds	British Columbia, Canada
Macdonald et al. (2003a) (2003b)	Evaluates the effects of three different harvest variable retention harvesting prescriptions on suspended sediment concentrations, stream temperature	67 years	25	SED SHD	BACI ACI experiment	Local, Baptiste watershed and Galuski watersheds	British Columbia, Canada
Martin & Grotefendt (2007)	Compared site conditions between riparian buffer strips and unlogged riparian stands using aerial photography to determine mortality and LWLWD recruitment	1 year data collection	9	LWLWD	ACI experiment	Regional, northern and southern portions of southeast Alaska	Alaska
May & Gresswell (2003)	Investigates the mechanisms responsible for LWLWD recruitment into streams.	2-years data collection	4	LWLWD, SED	modeling, Regression analysis	Local, North Fork of Cherry Creek Research Natural Area	Oregon

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
McIntyre et al. (2018)	Effectiveness of forest mgmt. in maintaining function for non-fish-bearing headwater streams on competent lithologies.	6 years	17 (four treatments)	SHD, SED, NUT, LW, LIT, stream temperature	BACI experiment	Regional	western Washington
McIntyre et al. (2021)	Continuation of McIntyre et al., 2018 to assess changes over a longer time period (up to 9 years post-harvest).	12 years	17 (four treatments)	LW, NUT, SED, SHD, stream temperature	BACI experiment	Regional	western Washington
Meleason et al. (2003)	Evaluate of the potential effects of different riparian mgmt. strategies on the standing stock of wood.	Simulation modeling of 720 years	1	LW, LW, D	Modeling	simulation of stream types common in PNW	PNW, hypothetical stream
Mueller & Pitlick (2013)	Examines the relative importance of lithology as a driver of sediment delivery into streams.	multiple datasets ranging 5-90 years	83	SED	spatial modeling, correlative analysis of historical data	Regional, Northern Rocky Mountains	ID, WY, MT
Murray et al. (2000)	Examined the influence of partial harvesting on stream temperature, chemistry, and turbidity 10-15 years post treatment.	2 years data collection, 10-15 years after treatment	1	NUT, SED, SHD	ACI experiment	Local, Rock and Tower Creek watersheds	Washington

Commented [AM6]: Not all of these are experiments.

Commented [AM17]: If you want to address unbalance across treatments that should be a different column. This is the total sample size. You could add number of treatments and numbers of sites within each of those treatments, but as this stands this should be the total N.

Commented [bs18R17]: Okay

Commented [AM19]: This is the state/provence. Not needed here?

Commented [bs20R19]: Okay

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Nowakowski & Wohl (2008)	Examined differences in wood load and valley/channel characteristics between managed and unmanaged riparian areas.	1 year data collection	19	LWLWD	ACI experiment	Local, Upper Tongue River and North Rock Creek watersheds	Wyoming
Pollock et al. (2009)	The influence of forest harvests on stream temperature.	2 months	33	SHD	ACI experiment	Local, Hoh river Basin, and Clearwater River Basin	Washington
Puntenney-Desmond et al. (2020)	The potential effect of climate change on sediment yield and concentrations in riparian area run-offs.	1 month	15	SED	BACI, simulated rainfall in field plots	Local, Star Creek headwater catchment	Alberta, Canada
Rachels et al. (2020)	Investigates the source of suspended sediment to a stream draining a recent harvested catchment.	1 summer	1	SED	ACI experiment	Local, Enos Creek	Oregon
Reid & Hassan (2020)	Combines a wood budget model and a 45-year record of LWLWD to examine changes in LWLWD characteristics.	Long-term dataset from 1973-2017, simulated 300 years	8	LWLWD	Simulation Modeling for framework development	Local, Carnation Creek	BC, Canada
Reiter et al. (2009)	Effects of forest practices on sediment production at the watershed-scale with 30 years of water quality data.	Long-term dataset from 1975-2005	4	SED	AI	Local, Deschutes River watershed	Washington

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Reiter et al. (2015)	Long-term combined effects of hydro-climatic factors and intensively managed forests with buffers on stream temperature.	Long-term dataset from 1975-2009	4	SHD	BAI experiment	Local, Deschutes River watershed	Washington
Reiter et al. (2009)	Effects of forest practices on sediment production at the watershed scale with 30 years of water quality data.	Long-term dataset from 1975-2005	4	SED	AI	Local, Deschutes River watershed	Washington
Reiter et al. (2020)	Effects of harvesting and variable buffer widths on stream temperature	10 years	Unbalanced: 3-7	SHD	BACI experiment	Local, Trask River Watershed	Oregon
Roon et al. (2021a)	Thinning effects of second growth redwood forests in northwestern California.	2 years	3	SHD	BACI experiment	Local, Tectah and Lost Man watersheds	California
Roon et al. (2021b).	Investigation of how different thinning intensities affect stream temperature via loss of canopy cover at local and watershed scales.	2 years	3	SHD, stream temperature	BACI experiment	Local, Tectah and Lost Man watersheds	California

Commented [AM6]: Not all of these are experiments.

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Safaeq et al. (2020)	Presents an approach at isolating the streamflow effect on sediment delivery post-harvest.	Long-term dataset, 1952-2016	2	SED	BACI experiment	Local, H.J. Andrews Experimental Watershed	Oregon
Schuett-Hames & Stewart (2019a)	comparison of LW inputs, tree fall, and stand structure 5 years post-harvest.	5 years	Unbalanced: 8-9	LW	ACI experiment	Regional, northeastern Washington, 1 site in East Cascades	Washington
Schuett-Hames & Stewart (BCIF), (2019b)	The study analyzes the changes in stand structure, buffer tree mortality, and riparian functions 10 years after upland timber harvest.	10 years	Unbalanced: 3-14	LW, LWD, SED, SHD	ACI experiment	Regional, western Washington Coast and Cascade Range	Washington
Schuett-Hames & Stewart (2019a)	comparison of LWD inputs, tree fall, and stand structure 5 years post-harvest.	5 years	Unbalanced: 8-9	LWD	ACI experiment	Regional, northeastern Washington, 1 site in East Cascades	Washington
Schuett-Hames et al. (2011)	Evaluates the effects of forest mgmt. on stream shade, LW, large woody debris, LW recruitment, and sediment delivery.	5 years	Unbalanced: 3-15	LW, LWD, SED, SHD	ACI experiment	Regional, western Washington Coast and Cascade Range	Washington
Six et al. (2022)	Assessed differences in levels of riparian buffer retention at mitigating changes to organic matter dynamics.	2 years	3	LIT, LW, LWD	BACI experiment	Local, Trask River Watershed	Oregon

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Sobota et al. (2006)	Study of riparian characteristics and their effects on tree fall direction and in-stream recruitment.	3 years	21	LWLWD	model with field data	Regional, Pacific Northwest and Intermountain West	Idaho, Washington, Oregon, Montana
Sugden et al. (2019)	Assessed the efficacy of Montana SMZ guidelines for controlling stream temperature.	2 years	30	SHD	BACI experiment	Regional, Western Montana	Montana
Swartz, et al. (2020)	Assessed whether experimental canopy gaps meant to mimic natural disturbances affect stream temperature	2 years	6	SHD	BACI experiment	Local, Mckenzie River Basin	Oregon
Teply et al. (2007)	Compares the effects of mgmt. harvest prescriptions and no-harvest RMZs on LWLWD recruitment in streams.	1 year data collection, 100 years simulated	58	LWLWD	Simulation Modeling	Local, Priest Lake Watershed	Idaho
Vanderbilt et al. (2003)	Correlation of nutrient inputs with weather events (mainly precipitation).	long-term datasets, ranging from 20-30 years	6	NUT	ACI experiment	Local, H.J. Andrews Experimental Watershed	Oregon
Warren et al. (2013)	Evaluates stand age and associated canopy structural differences on stream light in second-order streams.	1 year+ year data collection	2	SHD	ACI experiment	Local, H.J. Andrews Experimental Watershed	Oregon

Commented [AM6]: Not all of these are experiments.

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process	Experiment type/Study Type	Scale (local, regional, global)	State/Province
Wing & Skaugset (2002)	Examines the relationship between channel characteristics and LWLWD in streams.	Extensive spatial dataset from 1990-1996	3793	LWLWD	modeling, regression analysis	Regional, Western Cascade and Coast Range of Oregon	Oregon
Wise et al. (2010)	Uses tree rings to augment previous records to reconstruct multi-century data for the Snake River.	Dendrochronology records from 1600-2005	3	Drought Frequency	Climate reconstruction from dendrochronology records	Local, 3 sites in western Wyoming	Wyoming
Yang et al. (2021)	Examined the temporal variation in response of downstream water chemistry to prolonged drought and forest thinning.	5 years	2	NUT	BACI experiment	Local, The Kings River Experimental Watershed	California
Yeung et al. (2019)	Modelled the post-harvest response of leaf litter coarse particulate organic matter quantity in a coastal stream	Published data spanning 4-5 years	Total n not reported	LIT	Heuristic modeling	CPOM data from local streams in coastal BC	Model developed from multiple North American sites
Ehinger et al. (2021)	Effectiveness of riparian mgmt. in maintaining function in for non-fish-bearing headwater streams on incompetent lithologies.	95-6 years	Unbalanced ±6-11 (two treatments)§	LWD, NUT, SED, SHD, Stream Temperature	BACI experiment	Regional, southwestern Washington	southwestern Washington

Reference	Purpose	Sampling Period/Study Duration	Sample size (n)	Function / process		Experiment type/Study Type	Scale (local, regional, global)	State/Province
McIntyre et al. (2018)	Effectiveness of forest mgmt. in maintaining function for non-fish-bearing headwater streams on competent lithologies.		6 years	17 (four treatments)	SHD, SED, NUT, LW, LIT, stream temperature	BACI experiment	Regional	western Washington
McIntyre et al. (2021)	Follow-up study to the Continuation of McIntyre et al., 2018 to assess changes over a longer time periods (up to 9 years post-harvest).	12-5 years	Unbalanced: 3-617 (four treatments)	LWD, NUT, SED, SHD, stream temperature	BACI experiment	Regional, western Washington	western Washington	
McIntyre et al. (2018)	Effectiveness of forest mgmt. in maintaining function for small headwater streams on competent lithologies.		5-11 years	Unbalanced: 3-7	SHD, SED, NUT, LW, LIT	BACI	Regional, western Washington	western Washington
Deval et al. (2021)	Disturbance effects on stream chemistry.		13 years	7	NUT	BACI experiment	Local, Mica creek-Creek Experimental Watershed	Idaho

Commented [AM6]: Not all of these are experiments.

Commented [AM21]: If you want to address unbalance across treatments that should be a different column. This is the total sample size. You could add number of treatments and numbers of sites within each of those treatments, but as this stands this should be the total N.

Commented [AM22]: This is the state/province. Not needed here?

Commented [AM23]: Not clear why these are not in alphabetical order. Should be made so.

204 Discussion of findings relative to FPHCP objectives

205 Litter/Organic matter inputs/Nutrients

206 Prior to the Forest and Fish Report (1999), studies that directly quantify the effects of timber
207 harvest within riparian areas on litter and organic matter (OM) input into streams in managed
208 watersheds of western north America are sparse. Two seminal studies, one from the H.J.
209 Andrews experimental watershed studies (Gregory et al., 1987) and one from the Carnation
210 Creek experimental watershed (Hartman & Scrivener, 1990) present results that estimate loss of
211 litter input following harvest. Gregory et al., (1987) which was part of the Streamside
212 Management: Forestry and Fishery Management collection produced by Salo & Cundy (1987)
213 noted that removal of the forest canopy from timber harvesting resulted in decreases in annual
214 litter fall from 300-400 g/m² in the mature forests to less than 100 g/m². Further, they posit that
215 decreased litter inputs after logging can persist for 10 – 20 years before recovering. Results from
216 Hartman & Scrivener, (1990) showed that litter inputs post-logging were 25-50% of pre-logging
217 levels with about 50% of the loss recovering within a decade (note: buffer widths varied from 1-
218 70 m, litter input loss was not analyzed by buffer width).

219 Experimental studies published after 1999 that investigate the factors affecting litter and organic
220 matter (OM) input (not including LW) into streams in western North America are still relatively
221 few. In our search we found six papers that quantify the effects of timber harvest or the effects of
222 site factors (e.g., topography, vegetation characteristics) Four of these studies focus on headwater
223 streams and two of the studies reviewed here extend into larger fish-bearing streams (Bilby &
224 Heffner, 2016; Hart et al., 2013; Kiffney & Richardson, 2010; McIntyre et al., 2018; Six et al.,
225 2022; Yeung et al., 2019).

226 Studies specifically investigating controls on litter inputs used litter traps for sample collection
227 and quantify changes in litter delivery from dry weight. Before litter quantification, it is
228 commonly separated by type (e.g., leaves, twigs, cones, etc.), species (e.g., hardwood, conifer),
229 season, and distance from stream. Litter weights are usually compared with treatment (e.g.,
230 harvest intensities, buffer widths), site factors (e.g., slope, species composition, stand density,
231 distance to stream), and local weather conditions (e.g., precipitation, wind speed) with statistical
232 or simulation modeling.

233 In terms of site factors, Bilby & Heffner (2016) used a combination of field experiments,
234 literature review, and modeling to estimate the relative importance of factors affecting litter
235 delivery from riparian areas into streams of western Washington in the Cascade mountains at
236 high and low elevations. Their results showed that under the wind conditions recorded at
237 Humphrey Creek, most litter recruited into the stream originated from within 10 m of the stream
238 regardless of litter or stand type. No difference was found in delivery distance and litter type
239 (needles or broadleaf) at young sites. However, needles released at mature sites had a higher
240 proportion of cumulative input from greater distances than needles or alder leaves released at
241 younger sites. Litter travel distance was linearly related to wind speed ($p < 0.0001$). Doubling
242 wind speed at one site led to a 67-87% expansion of the riparian litter contribution zone in the

243 study area. The results also reveal a trend that suggests slope affects the width of the litter
244 contributing area. However, the authors did not apply statistical analysis to these values and only
245 speculate that increasing the slope from 0-45% would increase the width of the litter contributing
246 area by up to 71% for needles and 95% for leaves. From these results, Bilby & Heffner (2016)
247 suggest that wind speed has a strong effect on the width of litter delivery areas within riparian
248 areas, but that relationship is also affected by stand age (suggesting that tree height was a factor)
249 and litter type (deciduous vs. conifer). Other than stand structure and topography, another study
250 shows evidence of species composition affecting litter delivery into streams. Hart et al. (2013)
251 compared litter delivery into streams between riparian zones dominated by deciduous (red alder)
252 and coniferous (Douglas-fir) tree species in western Oregon. Results from this study show that
253 deciduous forests dominated by red alder delivered significantly greater vertical and lateral
254 inputs ($\text{g m}^{-2} \text{y}^{-1}$) to adjacent streams than did coniferous forests dominated by Douglas-fir.
255 Deciduous-site vertical litter input (mean = $504 \text{ g m}^{-2} \text{ y}^{-1}$) exceeded that from coniferous sites
256 ($394 \text{ g m}^{-2} \text{ y}^{-1}$) by 110 g/m^2 over the full year. Annual lateral inputs at deciduous sites (109 g
257 $\text{m}^{-2} \text{ y}^{-1}$) were $46 \text{ g m}^{-2} \text{ y}^{-1}$ more than at coniferous sites ($63 \text{ g m}^{-2} \text{ y}^{-1}$). The timing of the
258 inputs also differed, with the greatest differences occurring in November during autumn peak
259 inputs for the deciduous forests. Further, annual lateral litter input increased with slope at
260 deciduous sites ($R^2 = 0.41$, $p = 0.0771$), but showed no strong relationship at coniferous sites (R^2
261 $= 0.1863$, $p = 0.2855$). These results were partially consistent with Bilby & Heffner (2016) in
262 that they suggest litter type, and topography (slope) can affect the litter input rates. Lateral litter
263 movement in the riparian area increased with slope for deciduous riparian forests throughout the
264 year and for coniferous forests only in the spring and summer months.

265 In terms of the effects of timber harvest on litter and OM quantity in streams, 4 studies in our
266 review were found that provide experimental results that have been conducted since 2000 and
267 focus on western North America. Of these ~~4~~four studies, ~~1~~one used simulation modeling
268 (Yeung et al., 2019), and the other 3 (Kiffney & Richardson, 2010; McIntyre et al., 2018; Six et
269 al., 2022) used field-based experiments to estimate the effects of timber harvest within riparian
270 forests on OM inputs and dynamics in streams. Yeung et al. (2019) simulated post-harvest
271 responses to leaf-litter derived coarse particulate organic matter (CPOM) quantity in a coastal
272 rainforest stream in British Columbia, Canada. For this study, Yeung et al. (2019) used published
273 empirical data from representative small, forested streams in coastal British Columbia to
274 calibrate and set parameters for their CPOM model. The model compared the effects litterfall
275 reduction, increase in peak flows, and increase in stream temperature (estimated for 4 harvesting
276 intensities based on available data) on in stream CPOM standing stocks. Results showed
277 evidence that litterfall reductions from timber harvest was the strongest control on in-stream
278 CPOM quantity for 4 years post-harvest. However, when litterfall reductions were below 30%,
279 the effect size varied with relative changes to peak flows and stream temperature. Stream
280 temperature increases specifically showed a significant interaction with litterfall reductions. The
281 authors propose that the decreased activity of CPOM consumers caused by increasing stream
282 temperatures by $4 \text{ }^\circ\text{C}$ or more, may be enough to offset the loss of litterfall inputs of CPOM
283 stocks. This speculation was made based on the temperature dependent function of leaf litter
284 consumption by common shredder species and temperature ranges modeled by Stenroth et al.
285 (2014). This model predicts shredder activity is optimized at $\sim 15 \text{ }^\circ\text{C}$ (ranging between $13.7 -$

286 16,7°C) but begins to quickly decline at temperatures above 16 °C. The caveat of this study is
287 that it did not include LW dynamics in preserving CPOM post-harvest.

288 All four studies that applied an experimental design to assess the changes in litter and OM
289 delivery into streams used a Before-After Impact-Control (BACI) design. Also, all these studies
290 compared changes in litter and OM inputs into streams for two or more riparian forest harvest
291 prescriptions (Table A1; Appendix I). Kiffney & Richardson (2010) compared changes in litter
292 input between riparian harvest prescriptions that included clear-cut to stream edge, 10 m wide
293 buffer reserve, 30 m buffer reserves, and an uncut control over the course of 8 years. No thinning
294 was applied within the reserves. Upland treatment at all sites applied clearcut. Results showed
295 differences in litter flux relative to riparian treatment persisted through year 7, while a positive
296 trend between reserve width and litter flux remained through year 8. Needle inputs remained 6x
297 higher in the buffer and control sites through year 7, and 3-6x higher in year 8 than in the
298 clearcut sites. Twig inputs into the control and buffered sites were ~25x higher than in the
299 clearcut sites in the first year after treatment. The linear relationship between reserve width and
300 litter inputs was strongest in the first year after treatment, explaining ~57% of the variation, but
301 the relationship could only explain ~17% of the variation in litter input by buffer width by year 8
302 (i.e., the relationship degraded over time). The authors interpret these results as evidence that
303 litter flux from riparian plants to streams, was affected by riparian reserve width and time since
304 logging.

305 McIntyre et al. (2018) also [experimentally](#) assessed the difference in the changes in litterfall
306 inputs into streams following ~~three-four experimental~~ treatments: an unharvested control
307 (Reference), current Forest Practices that apply a two-sided 50-ft riparian buffer along at least
308 50% of the stream (FP; with clearcut to stream's edge outside of the buffer), a two-sided 50-ft
309 buffer along the entire stream (100%), and a clearcut to stream without a buffer (0%). The upland
310 forests of all treatments were clearcut harvested. Results for litterfall input showed a significant
311 decrease in total litterfall (includes leaves/needles, twigs, cones etc.) input in the FP and 0%
312 treatments between pre- and post-treatment periods (2 years of pre-, and 2 years of post-harvest
313 data). However, compared to the Reference streams, only the 0% treatment (unbuffered) showed
314 a significantly lower litterfall input post-harvest and only for deciduous leaves, and combined
315 total of deciduous leaves and conifer needles. The 100% buffer showed a non-significant
316 increase in litterfall inputs relative to the reference streams. The authors interpret these results as
317 evidence that the riparian vegetation community in the unbuffered treatment had not recovered
318 by the end of year 2 post-harvest.

319 Six et al. (2022) also investigated the effects of timber harvest on litter inputs. However, this
320 study had no replication in their design for each treatment and only 2 control sites (i.e., n = 1 for
321 each treatment). The results are presented here because there is a general lack of studies available
322 in the literature after 2000 that provide experimental evidence of the effects of riparian timber
323 harvest on litterfall inputs into streams. Six et al. (2022) compared changes in litterfall pre- and
324 post-treatment between sites with a complete clearcut to stream, a clear cut with leave trees
325 (retention of 5 trees per hectare), clearcut with a 15 m no-cut retention buffer, and an uncut
326 control. Because of the small sample sizes, no tests for significance could be applied. However,

327 the authors interpreted the data with descriptive statistics and graphical summaries. Their results
328 showed post-harvest litter delivery decreased for the clearcut with no leave trees but increased
329 for both the clearcut with leave tree and clear cut with retention buffer. These results are
330 somewhat consistent with those of McIntyre et al., (2018) which showed significant decreases in
331 litter delivery only in sites with no retention buffer.

332 The objective of the study from Wooton (2012) was to assess how riparian area treatments
333 impact river food webs with an emphasis on economically important salmonid species in an
334 Olympic Peninsula River in Washington state. However, they present results and statistical
335 analysis for differences in litter inputs ($\text{g m}^{-1} \text{hr}^{-1}$) between treated and untreated reaches.
336 Because of the lack of litter input studies in literature, their results are presented here. Wooton
337 (2012) removed the dominant tree species, red alder (*Alnus rubra*), from one bank along five
338 treatment reaches ranging from 100-300 m long and replaced them with conifer seedlings. Paired
339 control reaches were interspersed between treated reaches along the stream. Specific methods for
340 tree removal or width of buffer in treatment reaches were not reported. Leaf litter decreased
341 significantly ($p = 0.04$) in the treatment reaches compared to the control reaches ($4.92 + 2.55$ vs.
342 $14.12 + 5.70 \text{ g m}^{-1} \text{hr}^{-1}$).

343 *Nutrients*

344 Riparian timber management practices in the 1970s were developed for water quality standards
345 with the development of the Clean Water Act of 1972, based on nutrient concentrations and
346 water clarity. Before implementing these BMPs, timber harvest practices included clearcut to the
347 stream edge, burning of slash, and application of pesticides which resulted in large and
348 immediate increases in stream water nutrient concentrations that remained higher than pre-
349 harvest or reference stream values for months and even years (Brown, 1973; Fredriksen, 1975).
350 However, BMP development and implementation over the past several decades have shown
351 evidence of their effectiveness in minimizing these effects both in magnitude and across time
352 (Deval et al., 2021; Shah et al., 2022; Stednick, 2008). For example, Shah et al. (2022) in their
353 global review of the effects of forest management on water quality under contemporary
354 management practices concluded that the development of BMPs across the world has resulted in
355 reduced or in some cases, undetectable impacts on water quality. However, they also report that
356 harvest impacts on nutrient concentrations can be complex and depending on the management
357 practices implemented, their effects may manifest many years after the work has been completed
358 (e.g., slow decomposition of slash, regrowth of vegetation, changes in land use). Indeed,
359 Sweeney & Newbold (2014) in their literature review and synthesis on the efficacy of forest
360 buffers in protecting water quality based on buffer width, remark on the high variability of
361 responses across studies. They report that removal of nitrogen from upland sources per unit
362 width of a forested buffer varied inversely with subsurface water flux. This suggests factors that
363 influence water flux through the buffer (e.g., hillslope gradient, soil porosity, vegetation type and
364 composition, precipitation) also impact buffer efficacy in removing nutrients and pollutants.

365 Zhang et al. (2010) in a review and meta-analysis of the effectiveness of buffers in reducing
366 nonpoint source pollution found comparable results. They reported slope (hillslope gradient) as
367 having a linear relationship with buffer pollutant removal efficacy that switched from positive to

368 negative when slope increased beyond 10% (i.e., hillslope gradients of ~10% were optimal for
369 buffer efficacy in removing pollutants). However, there may be some variation in these
370 relationships based on the nutrient or pollutant observed (e.g. form of nitrogen, phosphorus, etc.).
371 For example, Vanderbilt et al. (2003) analyzed long-term datasets (ranging 20-30 years for each
372 watershed) to investigate patterns in dissolved organic nitrogen (DON) and dissolved inorganic
373 nitrogen (DIN) export with watershed hydrology. Their results showed that total annual
374 discharge was a positive predictor of annual DON export in all watersheds with R^2 values
375 ranging between 0.42 to 0.79. In contrast, relationships between total annual discharge and
376 annual export of nitrate ($\text{NO}_3\text{-N}$), ammonium ($\text{NH}_4\text{-N}$), and particulate organic nitrogen (PON)
377 were variable and inconsistent across watersheds. The authors speculate that different factors
378 may control organic vs. inorganic N export.

379 In our search of the literature, four studies were found that provide experimental evidence of the
380 effects of riparian timber harvest on nutrient flux in western north America and were published
381 since 2000. Gravelle et al., 2009 compared the effects of contemporary forest harvesting
382 practices in Idaho on nutrient cycling and in stream concentrations. This study followed the
383 BACI design and featured a pre-treatment measurement phase (5 years), a post-road construction
384 phase (5 years), and a post-harvest phase (5 years). Treatments imposed included a clearcut to
385 stream with 30-foot equipment exclusion zone (non-fish-bearing), a target reduction of 50% of
386 the canopy removal over 50% of the area, equating to 25% removal of existing shade (fish-
387 bearing streams), and was compared to an uncut reference. Results for the post-road construction
388 period showed no significant changes in any analyzed nutrient concentrations. Results for the
389 post-harvest period showed significant increases in monthly mean nitrate and nitrite (NO_3 and
390 NO_2) at sites immediately downstream from the clearcut, the partial harvest, and at sites
391 downstream from both treatments in the stream network (cumulative). The changes in monthly
392 mean NO_3 and NO_2 during the five years post-harvest were greatest for the clearcut treatment
393 ($+0.29 \text{ mg L}^{-1}$), followed by the cumulative ($+0.07$ and $+0.05 \text{ mg L}^{-1}$) and partial harvest ($+0.03$
394 mg L^{-1}). NO_3 showed progressively increasing monthly concentrations for 3 years after harvest
395 before declining. None of the other nutrients analyzed in this study (Kjeldahl nitrogen (TKN),
396 total phosphorus (TP), total ammonia nitrogen (TAN) consisting of un-ionized (NH_3) and ionized
397 (NH_4^+) ammonia, and unfiltered orthophosphate (OP) samples) showed significant changes
398 during the post-harvest period.

399 In a follow up study, Deval et al. (2021) compared changes to nutrient concentrations 8 years
400 after Gravelle et al. (2009) completed their study. During these 8 years (extended harvest period)
401 the extent and frequency of harvest operations increased. Treatments consisted of additional road
402 construction and timber harvest (clearcut), with site management operations including pile
403 burning and competition release herbicide application. Following these treatments, streams in all
404 harvested watersheds again experienced significant increases in $\text{NO}_3 + \text{NO}_2$ concentrations of
405 even higher magnitude than during the first post-harvest period. Further, there were also small
406 but significant increases in mean monthly total phosphorus (TP) concentrations at all treatment
407 sites, including the downstream cumulative site. Cumulative $\text{NO}_3 + \text{NO}_2$ concentrations increased
408 throughout the study but showed signs of recovery in one watershed approximately 3 years after
409 the last treatment (clearcut, broadcast burn, herbicide). The authors attribute the increase in

410 NO³+NO² and TP during the extended harvest periods (i.e., beyond what was observed in the
411 first post-harvest period) to the application of herbicides and broadcast burning.

412 In general, the authors of both these studies (Deval et al 2021; Gravelle et al., 2009) concluded
413 that Idaho BMPs for riparian forest harvest are effective in reducing sediment and pollutants into
414 streams. While there were significant increases in nitrate and nitrite concentrations following
415 management operations, levels never increased above acceptable values for water quality
416 standards and there was evidence of nitrogen recovery to pre-harvest (or unharvested) levels
417 after 3 years.

418 Considering the interaction between climate and forest harvest on nutrient transport, Yang et al.
419 (2021) investigated the effects of drought and forest thinning operations (independently and
420 combined) on stream and soil water chemistry in the Mediterranean climate headwater basins of
421 the Sierra National Forest. Data on water chemistry were taken 2 years prior and 3 years
422 following drought and thinning operations in two watersheds, each with thinned and control
423 stands. Young stands with high shrub cover (> 50%) were masticated to < 10% shrub cover. The
424 thinning prescription in mature stands removed trees across all diameter classes to a target basal
425 area range of 27–55 m² ha⁻¹ with target basal areas varying based on tree density. Thinning
426 extended into the riparian management zone. Trees within 15 m of the stream could be chainsaw-
427 felled and skidded, but mechanical equipment was excluded within 30 m of the stream. Results
428 showed that drought alone altered dissolved organic carbon (DOC) in stream water, as well as
429 altered the proportion of dissolved organic carbon to nitrogen (DOC: DON) in soil solution in
430 unthinned (control) watersheds. Volume-weighted concentration of DOC was 62% lower ($p <$
431 0.01) and DOC:DON was 82% lower ($p = 0.004$) in stream water and soil solution, respectively,
432 during years of drought than in years prior to drought. Drought combined with thinning altered
433 DOC and dissolved inorganic nitrogen (DIN) in stream water, and DON and total dissolved
434 nitrogen (TDN) in soil solution. For stream water, volume-weighted concentrations of DOC were
435 66- 94% higher in thinned watersheds than in control watersheds for all three consecutive
436 drought years following thinning. No differences in DOC concentrations were found between
437 thinned and control watersheds before thinning. The authors conclude that their results provide
438 evidence that the influences of drought and thinning are more pronounced for DOC than for
439 nitrogen in streams. They also speculate that the periodic changes in climate (e.g., seasonal,
440 drought) contribute to the high variability in carbon and nitrogen concentration in streams in
441 Mediterranean climates following harvest.

442 Specific to Washington, the Hard Rock (McIntyre et al., 2021) and the Soft Rock (Ehinger et al.,
443 2021) ~~studies~~ [Studies](#) also reported on changes in nutrient concentrations and nutrient export in
444 streams following riparian timber harvest along headwater streams of western Washington. [Hard](#)
445 [Rock](#) ~~t~~ Treatments included a 50 ft buffer along both sides of the stream for the entire RMZ
446 (“100%”), 50 ft buffer along at least 50% of the RMZ (“FP”), clearcut to stream (“0%”), and an
447 unharvested reference (Ref). Results for nitrogen and phosphorus concentrations in streams
448 showed that post-harvest changes for total-N or total-P were not significant for any of the
449 treatments relative to the Reference. The only significant difference detected post-harvest was for
450 nitrate-N concentration between the 0% buffer treatment and all other treatments. However, for

451 annual export (kg ha⁻¹ yr⁻¹), total-N and nitrate-N export increased post-harvest at all sites, with
452 the smallest increase in the 100% treatment and the largest in the 0% treatment. Compared to the
453 reference sites, analysis showed an increase in total-N export of 5.52 (P = 0.051), 11.52 (P =
454 0.0007), and 17.16 (P < 0.0001) kg ha⁻¹ yr⁻¹ in the 100%, FP, and 0% treatments, respectively, in
455 the first 2 years post-harvest. In the extended period (7-8 years post-harvest) export for total-N
456 remained higher in all treatments compared to the reference by 6.20 (P = 0.095), 5.34 (P =
457 0.147), and 8.49 (P = 0.026) kg ha⁻¹ yr⁻¹ for the 100%, FP, and 0% treatments, respectively.
458 Nitrate-N showed the same pattern with slightly lower values than total-N. The increase in total-
459 N and nitrate-N export from the treatment watersheds post-harvest was strongly correlated with
460 the increase in annual runoff (R² = 0.970 and 0.971; P = 0.001 and 0.001) and with the
461 proportion of the basin harvested (R² = 0.854 and 0.852; P = 0.031 and 0.031). The authors note
462 that there was high variability in the data for the extended period and nitrate-N export only
463 returned to pre-harvest levels in one watershed. Total-P export increased post-harvest by a
464 similar magnitude in all treatments: 0.10 (P = 0.006), 0.13 (P = 0.001), and 0.09 (P = 0.010) kg
465 ha⁻¹ yr⁻¹ in the 100%, FP, and 0% treatments (only analyzed during the 2-year post-harvest
466 period). The authors conclude that the 100% treatment was generally the most effective in
467 minimizing changes from pre-harvest conditions, the FP was intermediate, and the 0% treatment
468 was least effective. Thus, similar to the results of other studies reviewed, these results provide
469 evidence that the effects of timber harvest on nutrient export is proportional to the intensity of
470 the treatment (e.g. percent of basin harvested, presence of protective buffer).

471 *Summary of Factors Impacting Nutrient Concentrations and Export*

472 Similar to instream sediment concentrations and export, there is evidence from the studies
473 reviewed that nutrient dynamics are affected by the intensity of riparian timber harvest (e.g.,
474 presence of buffer widths, percent of basin harvested), changes in streamflow (either seasonally
475 or from harvest), climatic events (e.g., drought, heavy precipitation), physiography (e.g.,
476 hillslope gradient), and soil disturbance. The Soft Rock study (Ehinger et al., 2021) did analyze
477 changes in both sediment and nutrient flux following harvest for comparison with the Hard Rock
478 study. While the authors of this study report that the softer lithologies were more erodible than
479 the sites sampled for the Hard Rock study and that nutrient flux was within the range of results
480 for the Hard Rock study, effects of treatment and significant differences between studies could
481 not be detected because of limited sample sizes, inconsistent buffer widths, and timing of
482 harvestthe inability to statistically detect a change in export after harvest was likely due to the
483 extent of the riparian buffer (58% and 92% buffered), the unusually dry pre-harvest period, and
484 small number of sites.

485 In contrast to the results for sediment, there is evidence that changes in nutrient flux following
486 harvest can persist for considerably longer periods. This has been attributed to management
487 operations such as slash burning, herbicide or fertilizer application that directly affect nutrient
488 loads, and from decomposition of unburned downed wood and litter (Deval et al., 2021; Shah et
489 al., 2022). Results showed that instream dissolved organic carbon (DOC) concentrations of un-
490 thinned stands during drought years were lower, and aromatic DOC was higher than in non-

491 drought years. In-stream DOC concentrations were higher for three consecutive years following
492 thinning, than un-thinned stands.

493

495 [Large Wood \(LW\)/wood load/wood recruitment](#)

496 Large wood in streams is essential to create pools, regulate flow, and provide a slow pulse of
497 nutrients that help create and maintain salmonid habitat (Harmon et al., 1986). Sievers et al.
498 (2017), in a global meta-analysis of the effects of riparian alteration on trout populations, found
499 the most positive response of trout populations was with increasing in-stream wood and livestock
500 exclusion (+87.7% and +66.6%, respectively) from the riparian area. However, while most
501 studies show a positive relationship between increasing LW and salmonid populations, few have
502 examined long-term watershed-scale responses of increasing LW or studied a wide range of
503 species (oni et al. 2014). Large wood ~~debris~~ production and recruitment into streams can vary
504 between watersheds, and multiple studies have attempted to identify the drivers of LW
505 production and recruitment with varying results. For example, Benda et al. (2003) present a
506 wood budgeting framework, developed from 20 years of LW research based in the Pacific
507 Northwest, for riparian zones that includes numerical expressions for punctuated forest mortality
508 by important drivers they identify as fire, chronic mortality and tree fall, bank erosion and mass
509 wasting, decay, and stream transport. This framework can be applied to different regions by
510 adjusting parameter values to make predictions of the importance of landscape factors (e.g.,
511 climate, topography, basin size) on wood recruitment and abundance in streams for any area.
512 Depending on the region or landscape for which the framework is being applied, less common
513 but more locally important disturbances such as ice storms, ice breakage, and wind throw can
514 also be incorporated. This study and the framework it developed illustrate the diversity of the
515 wood recruitment, transport, and decay processes. The relative importance of each wood
516 recruitment mechanism, and the fate and transport of the in-stream wood depends on the
517 variation observed in the environmental, management, and vegetation factors of a site. Thus,
518 frameworks such as the one developed by Benda et al. (2003) help identify the relative
519 importance of these recruitment processes and their relationship with local landscape factors.

520 A Review of the Available Literature Related to Wood Loading Dynamics in and around Streams
521 in Eastern Washington Forests, was developed for CMER in October of 2004 (CMER 03-308,
522 2004). In this review, the researchers sourced 14 references with quantitative and descriptive
523 information relating to the correlation between wood volume and pieces of wood in streams and
524 the adjacent riparian community. The authors conclude that while the literature was incomplete,
525 several significant correlations existed between LW in streams and riparian zone stand
526 characteristics. For unmanaged (defined as unlogged and un-roaded) sites in Washington,
527 researchers reported positive correlations between the volume of LW in streams with adjacent
528 riparian zone mean tree height ($P < 0.001$), mean tree diameter ($P < 0.001$), and mean basal area
529 ($P < 0.001$). For numbers of LW pieces, positive correlations were found with the basal area
530 ($P < 0.007$) but no other vegetation characteristic of the adjacent riparian area. However,
531 regression analysis showed a significant positive correlation of LW piece quantity with core zone
532 trees/acre ($P < 0.001$, $R^2 = 0.45$) and core zone basal area/acre ($p = 0.004$, $R^2 = 0.29$). Relative to
533 managed riparian areas, streams adjacent to unmanaged riparian areas had significantly higher
534 LW volume. The most relevant sources of these results listed in this review were from Fox
535 (2001), Chesney (2000), Camp et al. (1997), and Knight (1990). Two other studies named in this
536 review (McDade et al., 1990; Fox, 2003) show evidence that as much as half of the wood found

537 in the streams could not be attributed to the adjacent designated riparian areas which indicates
538 the importance of scale when investigating in stream LW source.

539 In the western United States, several notable studies since 2000 have continued to investigate
540 and refine the factors important for LW recruitment. For example, Wing & Skaugset (2002)
541 investigated the relationships between land use, land ownership, and channel and habitat
542 characteristics with LW quantity and volume in stream reaches in western Oregon. The relevant
543 results (those derived for forested streams only) showed that stream gradient was the most
544 important explanatory variable for in-stream LW volume with the split in the regression analysis
545 occurring at 4.7%. Stream reaches with gradients less than 4.7% had on average less than half
546 the in-stream LW volume (11.3 m³ vs. 25.2 m³ per reach) than reaches with gradients >4.7%.
547 Results for LW pieces (logs at least 0.15 m diameter, and 3 m long) per 100 m length showed
548 bankfull width (BFW) as the most important explanatory variable with a split in the regression
549 analysis occurring at 12.2 m BFW. Reaches with a BFW <12.2 m averaged 11.1 LW pieces per
550 100 m compared to wider streams which averaged 4.9 pieces per 100 m. When the analysis was
551 constrained to “key” LW pieces (logs at least 0.6 m diameter and 10 m long), stream gradient
552 again emerged as the most important explanatory variable with the split in the regression
553 occurring at 4.9% stream gradient (mean key pieces per 100 m were 0.5 and 0.9 for gradients <,
554 and >4.9%, respectively). Following stream gradient and BFW, lithology was also an important
555 explanatory variable showing splits for Mesozoic and sedimentary lithologies (in 3 out of 4
556 analyses) grouped as containing half the LW quantity (pieces, key pieces, volume) on average
557 than all other geologies (basalt, cascade, and marine sedimentary geologies). Wing & Skaugset
558 (2002) suggests that geomorphic characteristics, in particular stream gradient and bankfull width,
559 but also underlying lithology in forested areas correlate best with LW presence in headwater
560 streams of western Oregon.

561 Another study from the Oregon Coast Range, May & Gresswell (2003), compared LW
562 recruitment processes between small colluvial channels and larger alluvial channels. Results
563 from this study showed that LW derived from local hillslopes and riparian areas accounted for
564 the majority of pieces (63%) in small colluvial channels. In contrast, the larger alluvial channel
565 received wood from a greater variety of sources, including recruitment from local hillslopes and
566 riparian areas (36%), fluvial redistribution (9%), and debris flow transported wood (33%).
567 Further, distributions of the source distance of wood pieces were significantly different between
568 colluvial and alluvial channels. In colluvial streams, 80% of total wood and 80% of total wood
569 volume recruited to colluvial streams originated from trees rooted within 50 m of the channel. In
570 the alluvial channel, 80% of the pieces of wood and 50% of the total volume originated from
571 trees which came from within 30 m of the channel. Considering the mechanisms responsible for
572 recruitment, for both colluvial and alluvial stream channels, slope instability exhibited the
573 longest source distance (median source distance = 40 m), followed by windthrow (median source
574 distance = 20 m), then natural mortality (median source distance = 18 m), and for obvious
575 reasons, bank erosion had the shortest median source distance (2 m). Compared between channel
576 types (colluvial vs. alluvial), the median source distance of wood recruited by windthrow was
577 significantly greater in colluvial channels than in the alluvial channel ($p < 0.05$). Source
578 distances for all other processes did not differ significantly between channel types. May &

579 Gresswell (2003) interpret these results as evidence that stream size and topographic position
580 strongly influence processes that recruit and redistribute wood in channels. Processes of slope
581 instability were shown to be important conveyors of wood from upland forests to small colluvial
582 channels. In the larger alluvial channels, windthrow was found to be the dominant recruitment
583 process from adjacent riparian area.

584 Three larger scale studies from Washington (Fox & Bolton, 2007), the northwestern United
585 States (Sobota et al., 2006), and the Columbia River Basin (Hough-Snee et al., 2016) present
586 results from simulation modeling or statistical modeling for site and physiographic factors
587 influencing LW recruitment and in stream loading. Sobota et al. (2006), in a landscape-wide
588 study of factors affecting tree fall direction and LW recruitment in watersheds of the Pacific
589 Northwest (data sourced from Washington, Oregon, Idaho, and Montana), found valley
590 constraint to have the strongest correlation with in-stream woody debris. Outputs from their
591 model showed that riparian areas in channels with >40% valley side slopes had the highest
592 tendency for tree fall towards streams; in these steep slope valleys, recruitment of large wood in
593 streams was 1.5-2.4 times greater than on moderately sloped landforms (< 40%).

594
595 Fox & Bolton (2007) modeled LW values from 150 stream segments located in unmanaged
596 watersheds, across Washington, with landscape, reach, and stand characteristics to understand the
597 central tendency of instream LW values in “natural” fish-bearing streams. Outputs from their
598 models show evidence that in-stream wood volume (m³ per 100 m stream length) and LW piece
599 count for streams up to 20 m in bankfull width (BFW) increased with drainage area and as
600 streams became less confined with BFW being a significantly better predictor of wood
601 parameters than basin size. Also, in-stream wood volume increased with adjacent riparian timber
602 age as determined by the last stand replacing fire. In this study (Fox & Bolton, 2007), the authors
603 noted that other predictor variables (e.g., gradient, bedform) also showed some evidence of an
604 effect but the variability of these variables were too great to evaluate with confidence.

605
606 Hough-Snee et al. (2016) reported similar issues with their results using Random Forest (RF)
607 models developed from field data to identify relationships between hydrogeomorphic and
608 ecological attributes that influence instream wood accumulation. Final RF models explained
609 43.5% of the variance in volume and 42.0% of the variance in frequency of in stream wood
610 loads. Mean annual precipitation, riparian large tree cover, and watershed area were estimated as
611 the most important predictors of in stream wood loads. However, so did individual watershed
612 which showed there was an interaction with site (i.e., site conditions unaccounted for may be
613 affecting the response). Given the heterogeneous results across all sub-basins studied, the authors
614 conclude by emphasizing the importance of incorporating local data and context when building
615 wood models to inform future management decisions.

616
617 Multiple studies have also investigated the effects of timber harvest under varying riparian
618 management zone prescriptions on LW recruitment. Specific to Washington, Schuett-Hames and
619 Stewart (2019a) compared in stand structure, tree fall rates, and LW recruitment between riparian
620 management zones harvested under the current standard Shade Rules (SR), the All-Available
621 Shade Rule (AAS), and unharvested references for fish-bearing streams in the mixed conifer

622 habitat type (2500 - 5000 feet elevation) for eastern Washington. Both shade rules have a 30-ft
623 no-cut buffer (core zone) immediately adjacent to the stream. The SR prescription allows
624 thinning in the buffer zone 30-75 feet (inner zone) from the stream while the AAS prescription
625 requires retention of all trees providing shade in this area. Results showed that cumulative wood
626 recruitment from tree fall after the five-year post-harvest interval was highest in the SR group,
627 lower in the AAS group and lowest in the REF group. The SR and AAS LW recruitment rates by
628 volume were nearly 300% and 50% higher than the REF rates, respectively. Wood recruitment in
629 the SR sites was significantly greater than in the AAS and reference sites. Conversely, wood
630 recruitment did not differ significantly between the AAS and reference sites. Considering the
631 source distance of post-harvest recruited LW, most recruited fallen trees originated in the core
632 zone (76%, 72%, and 64% for the REF, AAS and SR groups, respectively), while the proportion
633 from the inner zone (30–75 feet from the stream) was ~10% greater for the SR group compared
634 to the AAS and REF groups. These results suggest that while treatment of SR sites is intended to
635 increase resistance to disturbances such as fire and disease, it also provides evidence that these
636 treatments increase the susceptibility to windthrow and thus increases mortality relative to
637 reference sites five years post-harvest. Further, thinning treatments in the inner zone appeared to
638 change the spatial pattern (source distance) of wood recruitment from fallen trees. It is important
639 to note that this was a short-term study (5 years). The authors remark that LW recruitment is a
640 process that can change over decadal time scales, and follow-up monitoring is recommended.

641 Four similar studies conducted for [non-fish-bearing](#) streams in western
642 Washington compared changes in LW recruitment and stand mortality following harvest
643 (Ehinger et al., 2021; McIntyre et al., 2021; Schuett-Hames et al., 2011; Schuett-Hames et al.,
644 2019b. Schuett-Hames et al., (2011) and Schuett-Hames & Stewart (2019b) investigated
645 changes in riparian stand mortality and LW recruitment into the bankfull channel 5- and 10-years
646 post-harvest, respectively. Treatments for riparian forests adjacent to non-fish-bearing streams
647 evaluated in these studies include clearcut to stream edge, upland clearcut with a 50-foot no cut
648 buffer, and these were compared to unharvested reference streams. Results showed that tree fall
649 rates (annual fall rates of live and dead standing stems combined) was over 8 times and 5 times
650 higher in the 50-foot buffers than in the reference buffers 3 years after treatment when compared
651 as a percentage of standing trees and as trees/acre/yr, respectively. These differences were
652 significant for both metrics ($p \leq 0.001$). Total tree-fall rates in the period 4-5 years after
653 treatment, while still higher in the 50-foot buffers was not significant.

654 Over the entire five-year period, the percentages of standing trees that were uprooted and broken
655 (as well as the combined total) were significantly greater in the 50-foot buffer than in the
656 reference. Differences in mortality followed a similar pattern to tree fall rates. In the 50-foot
657 buffer sites, mortality rates were significantly higher (3.5 times higher) than in the reference sites
658 for the first three years following harvest. However, in years 4-5 mortality rates increased in the
659 reference buffers after high-intensity storms resulting in non-significant differences in mortality
660 during this period. The cumulative percentage of live trees that died over the entire five-year
661 period was 27.3% in the 50-ft buffers compared to 13.6% in the reference reaches, but the
662 difference was not statistically significant. This was likely because of the high variability in
663 mortality between sites in the 50-foot buffers. The data for mortality rates in the 50-foot buffers

664 had a bimodal distribution with most sites exhibiting less than 30% mortality, although three
665 sites (of 13) exhibited mortality rates greater than 50%.

666 For LW recruitment into the bankfull channel, results showed during the first three years after
667 treatment recruitment rates were 8 times and 14 times higher in the 50-foot buffers than in the
668 reference buffers respectively. The differences in pieces/acre/year and volume/acre/year between
669 reference and 50-foot buffers were significant. In years 4-5 after harvest LW recruitment
670 decreased in the 50-ft buffers and increased in the reference patches, and the number of recruited
671 LW pieces/acre/yr was greater in the reference patches, although the volume of LW recruited was
672 greater in the 50-ft buffers. Differences in recruitment rates between the 50-foot buffer and the
673 reference buffers for the 4–5-year period were not significant. For the entire first 5 years after
674 harvest, the 50-ft buffers recruited about twice the number of LW pieces recruited in the
675 reference patches, and over 3 times the volume; differences were marginally significant.

676 The results of the 10-year follow-up study for these sites (Schuett-Hames & Stewart, 2019b)
677 showed that stand mortality in the 50-foot buffer sites had stabilized and showed a cumulative
678 14.1% reduction in live basal area, while the reference stands showed a 2.7% increase in live
679 basal area. The differences in these values were not significant. Cumulative LW recruited into the
680 stream channel over the 10-period was double in the 50-ft treatment streams compared to the
681 reference streams. However, the majority of the LW recruited in the 50-ft treatment streams came
682 to rest above the streams, providing shade but not affecting streamflow, pool formation, or
683 sediment storage. Further, while the 50-ft buffer treatment provided more LW recruitment in the
684 short-term (10-years), the authors speculate there is a reduction in future LW recruitment
685 potential given the removal of trees outside the 50-ft buffer.

686 Two other studies which evaluated changes in LW following riparian forest harvest along non-
687 fish-bearing streams in western Washington were complimentary studies. Treatment sites in these
688 studies were underlain by either competent (McIntyre et al., 2018, 2021; also referred to as Phase
689 2 of the “Hard Rock” study), or incompetent (easily eroded) marine sedimentary lithologies
690 (Ehinger et al., 2021; also referred to as the “Soft Rock” study). The buffer treatments evaluated
691 for these studies were compared against unharvested reference sites (“REF”) and included a
692 minimum two-sided 50-ft wide riparian buffer along the entire reach (“100%”) ~~and, and~~ the
693 standard Forest Practices treatment (FP), a minimum two-sided 50-ft wide riparian buffer along
694 at least 50% of the RMZ (buffered and unbuffered portions were analyzed separately; hereafter
695 referred to as FPB for the buffered portion, and 0% for the unbuffered portion). However,
696 because of unstable slopes in some of the sites in the Soft Rock study (Ehinger et al., 2021),
697 many of the buffers were required to be wider than 50-feet (ranging from 18 –160% wider than
698 50-feet). Conversely, some of the sites ~~treated ended up with~~ had additional buffers that were
699 narrower than 50 feet. Further, there was limited availability of sites that fit the criteria (marine
700 sediment lithology, timing of treatment). Because of these limitations, statistical analysis and
701 comparison of LW response between treatments and references ~~could was not be performed~~ as a
702 part of the Soft Rock study. Thus, the results are only descriptive, but they provide an useful
703 information for informative comparison to the Hard Rock study.

704 Results from the Soft Rock study showed mean cumulative post-harvest mortality during the 3-
705 year post-harvest interval was only 6.5% of live density (trees/ha) in the reference sites. In
706 contrast, mean post-harvest mortality in the full buffer sites and the <50 ft buffer sites were 31
707 and 25% of density, respectively. However, there was considerable variation in mortality among
708 sites, exceeding 65% in two full buffer treatment sites. Windthrow and physical damage from
709 falling trees accounted for ~75% of mortality in the full and <50 ft buffers. In contrast to the
710 treated sites, <10% of trees died due to wind or physical damage in the reference sites. For LW
711 recruitment, there was an increase in pieces of LW per 100 m length of stream in the full buffers
712 (8%) and the unbuffered treatments (13%) and a decrease in the streams adjacent to buffers < 50
713 feet wide (-15%) 3 years after harvest. The Hard Rock study did not require changes to the
714 grouping of treatments (i.e., all treatment buffers were harvested as described above; e.g.,
715 Reference, 100%, FPB, 0%). Also, the Hard Rock study collected up to 9 years of post-harvest
716 data that allowed for the comparison of LW changes over time pre- to post-harvest, and between
717 treatments.

718 Results for the Hard Rock study showed that by year 8 post-harvest mortality as a percentage of
719 pre-harvest basal area was lower in the reference (16.1%) than in the 100% (24.3%) and FPB
720 (50.8%) treatments. The FPB–Reference contrast in mortality was not significant 2 years post-
721 harvest, but it was at 5- and 8-years post-harvest as mortality in FPB increased relative to the
722 Reference over time. The contrast in mortality between the 100% and Reference were not
723 significant for any time interval 8 years post-harvest. Wind/physical damage was the primary
724 cause of mortality for all treatments, including the Reference. In the 100% treatment it accounted
725 for 78% and 90% of the loss of basal area and density (trees/ha), respectively; in FPB it
726 accounted for 78% and 65% of the loss. Wind accounted for a smaller proportion of mortality in
727 the Reference RMZ (52% and 43%, respectively). LW–Large wood recruitment to the channel
728 was greater in the 100% and FPB RMZs than in the reference for each pre- to post-harvest time
729 interval. Eight years post-harvest mean recruitment of large wood volume was two to nearly
730 three times greater in 100% and FPB RMZs than in the references. Annual LW recruitment rates
731 were greatest during the first two years, then decreased. However, there was a great deal of
732 variability in recruitment rates within treatment sites and the differences between treatments
733 were not significant. Mean LW loading into the channel (pieces/m of channel length) differed
734 significantly between treatments in the magnitude of change over time. There was a 66%, 44%
735 and 47% increase in mean large wood density in the 100%, FP and 0% treatments, respectively,
736 in the first 2 years post-harvest compared with the pre-harvest period and after controlling for
737 temporal changes in the references. By year 8, only the FP treatment showed a significantly
738 higher proportional increase (41%) in wood loading when compared to the reference. In the time
739 interval 2-8 years post-harvest wood loading in the 100% treatment stabilized and began to
740 decrease in the 0% treatment.

741 The Hard Rock and Soft Rock studies showed similar results. Both studies showed an increase in
742 stand mortality that ~~also led to~~resulted in an increase in LW recruitment into the channels
743 adjacent to 50-foot (and greater in the Soft Rock) riparian ~~buffer treatments~~ relative to
744 unharvested reference sites. However, the longer time period of study in the Hard Rock
745 study ~~showed~~demonstrated that both mortality and ~~thus~~ LW recruitment began to stabilize after

746 ~~year~~ five years post-harvest. The results presented by Schuett-Hames (2012, 2019b) showed a
747 similar pattern of an initial increase in mortality rates and LW recruitment rates in treated stands
748 relative to untreated stands within three years of treatment, but stabilization within 5-10 years.
749 Unfortunately, because of the limitations in sample size and buffer width consistency in the Soft
750 Rock study, confident conclusions on the effects of lithological competency on LW recruitment
751 post-harvest cannot be drawn.

752 All studies reviewed above, which investigate the effect of timber harvest with riparian buffers
753 on tree mortality and large wood recruitment, show that the initial increase in mortality within
754 treatment buffers relative to reference buffers is primarily a result of increased windthrow
755 mortality. Liquori (2006) found similar results in an investigation of treefall characteristics
756 within riparian buffer sites ranging in width from 25-100 feet along ~~non-fish-bearing~~non-fish-
757 bearing and ~~fish-bearing~~fish-bearing streams. Within no-cut buffers, windthrow caused mortality
758 was up to 3 times greater than competition induced mortality for 3 years following treatment
759 with tree fall probability highest in the outer areas (closest to upland clearcuts) of the buffers.
760 Their results showed that treefall was generally highest at the outside edges of buffers (50+ feet),
761 representing about 60% of the total observed treefall, while the 0-25-foot zone represented
762 ~18%, and the 25-50-foot zone represented ~22%. This suggests an increase in windthrow
763 susceptibility within riparian buffers with increasing distance from the stream. Liquori (2006),
764 however, did not differentiate thinning treatments applied to the outer zones of the buffer in their
765 analysis mentioning “very modest” thinning was applied to some buffers. They suggest in their
766 interpretation of the results that buffer thinning may influence the depth to which wind forces can
767 penetrate into the buffer. The results from Schuett-Hames & Stewart (2019a), discussed above,
768 show evidence that thinning in the outer area (30-75 feet from bankfull width) changed the
769 source distance curve of wood recruitment from fallen trees with thinned buffers (SR
770 treatments). The results exhibited statistically higher overall treefall rates with a larger
771 percentage coming from the outer area in the SR treatments than in the reference and more
772 lightly thinned (AAS) treatment buffers.

773 Outside of Washington, but in areas with similar habitats (Oregon, British Columbia) several
774 experimental studies that have investigated the effects of timber harvest on treefall, mortality,
775 LW recruitment, and LW source distance have found comparable results to those conducted in
776 Washington. For example, Martin & Grotefendt (2007) compared riparian stand mortality and in-
777 stream LW recruitment characteristics between riparian buffer strips with upland timber harvest
778 and riparian stands of unharvested watersheds using aerial photography in the northern and
779 southern portions of Southeast Alaska. All buffer strips in this study were a minimum of 20 m
780 wide and included selective harvest within the 20 m zone (thinning intensity not specified or
781 included in the analyses as an effect). The results from this study showed significantly higher
782 mortality (based on cumulative stand mortality: downed tree counts divided by standing tree
783 counts + downed tree counts), significantly lower stand density (269 trees/ha in buffer units and
784 328 trees/ha in reference units), and a significantly higher proportion of LW recruitment from the
785 buffer zones of the treatment sites than in the reference sites. Also, results showed that mortality
786 varied with distance to the stream. Differences in mortality for the treatment sites were similar to
787 the reference sites for the first 0-10 m from the stream (only a 22% increase in the treated sites).

788 However, mortality in the outer half of the buffers (10-20 m) from the stream in the treatment
789 sites was more than double (120% increase) what was observed in the reference sites. The
790 authors attribute the difference in cumulative stand mortality to the increase in windthrow
791 susceptibility. Mortality attributed to windthrow was twofold and fivefold greater in the inner
792 and outer halves of the treatment buffers than in the reference buffers, respectively.

793

794 Bahuguna et al. (2010) evaluated the difference in windthrow caused mortality between 10 m, 30
795 m buffer widths (neither had thinning within the buffer and both had upland clear-cuts) and
796 unharvested controls in the Coast Mountains, British Columbia. Following harvest, 11% of
797 initially standing timber was blown down in the first and second years in the 10 m buffer,
798 compared to 4% in the 30 m buffer, and 1% in the unharvested controls. However, after 8 years
799 post-harvest, a significant amount of annual mortality occurred when winter storms brought
800 down multiple trees in the unharvested control at 30%, compared to 15% in both 30 m and 10 m
801 buffers. These results show evidence that timber harvest can increase windthrow caused
802 mortality within protective buffers in the short term but can stabilize within a decade. Further,
803 this study shows evidence that windthrow caused mortality is stochastic and large storm events
804 can cause just as much if not higher mortality within untreated riparian forests.

805 Burton et al. (2016) examined the relationship between annual in-stream wood loading and
806 riparian buffer widths adjacent to upland thinning operations. No-cut buffer widths were 6, 15, or
807 70 meters, and upland thinning was to 200 trees per ha (tph), with a second thinning (~10 years
808 later) to ~85 tph, alongside an unthinned reference stand ~400 tph. Their results showed that
809 slightly higher volumes of wood were found in sites with a narrow 6-m buffer, as compared with
810 the 15-m and 70-m buffer sites in the first 5 years after the first harvest and maintained through
811 year 1 of the second harvest (end of study). The authors attributed this difference to a higher
812 likelihood of logging debris and/or windthrow, but these factors were not analyzed. Considering
813 source distance, the authors used a mixed modeling approach to assess the relationship between
814 wood volume and source distance for in-stream wood with an identifiable source. This model
815 was only applied to the 70-meter buffer. The results showed that 82-85% of the wood with
816 discernable sources (90% for wood in early stages of decay; 45% of wood in late stages of
817 decay) came from within 15 m of the stream, and the relative contribution of wood to streams
818 declined rapidly with increasing distance. Still, these results are similar to those presented by
819 Schuett-Hames & Stewart (2019a) which showed the majority of the LW recruited (72-76% for
820 treated stands) into the channel were from within the first 30 feet (~9.1 m) of the stream even
821 though upland harvest prescriptions in this study differed from those evaluated by Burton et al.
822 (2016) (e.g., clearcut vs thinning).

823

824 *Summary of Factors Impacting LW Loads and Recruitment*

825 In general, the studies reviewed above show evidence that upland timber harvest with riparian
826 retention buffers initially increases stand mortality within the buffers and increases LW
827 recruitment relative to unharvested reference stands in the short-term. This increase in mortality

828 and LW recruitment is attributed to an increase in the susceptibility to windthrow within the
829 riparian buffers relative to the unharvested controls. Further, multiple studies (Liquori, 2006;
830 Martin & Grotefendt, 2007, Schuett-Hames & Stewart 2019a) showed evidence that the increase
831 in windthrow caused mortality is highest in the outer area of the riparian buffers (area closest to
832 upland treatments). There is some evidence that thinning within the buffer can also affect
833 mortality rates, but these studies are few. In the three studies that collected post-harvest data for 8
834 or more years (Bahuguna et al., 2010; McIntyre et al., 2021; Schuett-Hames & Stewart 2019b),
835 there is indication that mortality in the riparian buffers and annual LW recruitment into adjacent
836 streams stabilizes within 5-10 years. However, in the subsequent decades following treatments
837 with upland clearcuts there is evidence that LW recruitment rates can continue to decrease and in
838 stream wood loads may become depleted before recruitment rates can recover (Nowakowski &
839 Wohl, 2008; Reid & Hassan, 2020) depending on applied management practices (e.g., buffer
840 widths, road construction, etc.). For example, Teply et al. (2007) used simulation modeling to
841 estimate the effectiveness of Idaho Forest Practices for riparian buffers and found no significant
842 difference between predicted LW loads for harvested and unharvested sites 30-, 60-, or 100-years
843 post-harvest.

844 While the general conclusions of short-term increase in LW and long-term reduction of LW
845 following treatment are similar among studies it is more apparent that LW recruitment dynamics
846 are complex and highly variable even within treatment groups; and local site and landscape
847 factors may interact with treatments making it difficult to generalize the effectiveness of different
848 protective buffer treatments on preserving LW recruitment and in-stream wood loads. Indeed, the
849 LW budget framework created by Benda et al. (2003) emphasizes the importance of including
850 local physiographic, site, and disturbance factors. Additionally, the studies reviewed above
851 present results from experimental studies that vary greatly in their design. Buffer widths, riparian
852 and upland treatment prescriptions differ by region, state, and local regulations that can differ
853 further by stream type and size, and location within the landscape (e.g., elevation). Thus, general
854 global conclusions about the effect of riparian forest treatment on LW dynamics are difficult to
855 discern.

856

857

858

859 Bank Stability and Sediment

860 *Bank Stability*

861 Few studies could be found that quantify how riparian area harvest directly affects bank stability
862 or bank erosion based on our search criteria. Many studies published since 1999 that investigate
863 bank stability and bank erosion compare relative rates of erosion based on the presence/absence
864 of vegetation, type of vegetation (e.g., grassland vs. forest cover), and soil types or lithology
865 (Konsoer et al., 2015; Micheli et al., 2004; Simon & Collision, 2001; Wynn & Mostaghimi,
866 2006). Also, many studies have investigated the relative effects of different types of land use
867 (e.g., agricultural, urban, forested) as well as cattle grazing intensity (McInnis & McIver, 2009;
868 Zaines & Schultz, 2014). The only studies that could be found that provide some experimental
869 evidence as to how timber harvest within the riparian area affects bank stability or erosion come
870 from ~~3~~three CMER reports (Ehinger et al. 2021; McIntyre et al. 2018, Schuett-Hames et al.,
871 2011; Schuett-Hames & Stewart, 2019).

872 Schuett-Hames et al. (2011) investigated how soils and streambanks were disturbed following
873 harvest within the riparian area along perennial ~~non-fish-bearing~~non-fish-bearing streams (Type
874 Np) in western Washington. To evaluate post-harvest soil and stream bank disturbance, Schuett-
875 Hames et al. (2011) first described a soil erosion feature as areas of exposed soil that (1) had a
876 surface area of greater than 10 square feet, and (2) was caused by harvest practice (e.g., felling,
877 bucking, or yarding). If both criteria were met, the length, width, and distance to stream were
878 recorded, and evidence of sediment delivery to the stream was noted. The number of harvest
879 related soil disturbances were grouped by 100 ft lengths of stream, as were the number of
880 features delivering sediment to the stream. Disturbances along stream bank were quantified using
881 the same methods. The surface area (mean width x length) of disturbance features were used to
882 estimate the percent coverage of soil disturbance within 50-feet of bankfull width and in the
883 equipment exclusion zone (ELZ; within 30 feet of the bankfull width). Finally, the percent of
884 harvested patches with a greater than 10% coverage of soil disturbance features in the ELZ were
885 also quantified (performance target for bank stability). These methods were used to collect data
886 for all 3 harvest treatments. These harvest treatments included 1) a 50-foot wide no cut buffer, 2)
887 clearcut, no buffer, and 3) a 56-foot radius no-cut buffer surrounding the perennial initiation
888 point (PIP). A non-parametric, two-sample Mann-Whitney U test was used to test differences in
889 mean soil and stream bank disturbance metrics between the 50-foot buffer patches and the
890 clearcut (no buffer) patches. A Fisher's exact test was used to test for differences in the relative
891 frequency of patches exceeding the performance target (more than 10% of ELZ area disturbed by
892 management related activities) between 50-foot and the clearcut buffer prescriptions.

893 Results showed that the differences between the mean values of harvest related soil and
894 streambank disturbances for clear-cut patches and the 50-ft buffers were significant for all
895 metrics (e.g., # of bank disturbance features per 100 ft, # of soil disturbance features per 100 feet,
896 # of soil disturbance features, # of soil disturbance features delivering sediment to stream, % of
897 ELZ with soil disturbance; $P \leq 0.082$). Results for soil disturbance performance targets showed
898 that all of the 50-foot buffer and PIP prescriptions met the performance targets (i.e., maintained
899 <10% harvest-related soil disturbance in the ELZ). One clearcut patch exceeded the 10%

900 coverage performance target. The difference between clearcut patches and 50-foot buffer patches
901 was significant ($p = 0.007$). The average size of harvest related soil disturbances that delivered
902 sediment to streams was 752 ft² (range: 31-9060 ft²). The average size of soil disturbance
903 features that did not deliver sediment to streams was 65 ft² (range: 13 – 214 ft²). Delivery of
904 sediment to streams was best predicted by the horizontal distance between the soil disturbance
905 and the stream channel ($P < 0.0001$). The average distance to the stream for soil disturbance
906 features that delivered sediment was 1 ft (max. = 7.7), while the average distance for non-
907 delivering soil disturbance features was 14 ft (min 3.3). Using distance-to-stream alone, 96% of
908 the observations were correctly predicted based on whether the horizontal distance to the stream
909 was greater or less than 5.4 ft ($R^2 U4 = 0.80$). The authors concluded there were more harvest-
910 related soil disturbances following harvest in the clear-cut patches than the 50-ft buffers. Further,
911 that the management practices for the 50-foot and PIP buffers were sufficient at maintaining
912 bank stability performance targets. The clearcut patches were mostly sufficient at maintaining
913 performance targets with the exception of one site.

914 Schuett-Hames et al. (2011) also collected data on soil disturbance associated with post-harvest
915 root pits created from trees being uprooted by wind or other disturbances. Four metrics were
916 used to evaluate soil disturbance associated with uprooted trees: *Root-pits per acre*. Root-
917 pits/acre was calculated by tallying the number of root-pits in each patch and dividing by the
918 patch acreage. *Root-pits per 100 ft of stream length*. Root-pits/100 ft of stream length was
919 calculated by tallying the number of root-pits in each patch (both sides of the stream), dividing
920 by the stream length, and multiplying by 100. *Root-pits with sediment delivery per acre*. Root-
921 pits/acre with evidence of sediment delivery to the channel was calculated by tallying the number
922 of root-pits where evidence of sediment delivery to the stream channel is observed in each patch
923 and dividing by the patch acreage. *Root-pits with sediment delivery per 100 ft of stream length*.
924 Root-pits with sediment delivery/100 ft of stream length were calculated by tallying the number
925 of root-pits with evidence of sediment delivery in each patch (both sides of the stream), dividing
926 by the stream length, and multiplying by 100. These metrics were measured 3 years and 5 years
927 following harvest to give an annual rate of change for each metric at 3 years, from 3-5 years, and
928 for the entire 5 years. These standardized annual rates were compared between each treatment
929 patch type and a unharvested reference patch of the same size.

930 Results showed that in the first three years after harvest, the mean annual rate of total root-pit
931 formation (all root-pits) in the 50-ft buffers was over 10 times higher than the reference rate. This
932 difference was significant ($p = 0.002$). A similar result was found in the difference between root
933 pits delivering sediment to streams ($p = 0.002$). The mean total root-pit formation rate in the
934 clear-cut patches was much lower than the reference rate (likely because there were less trees to
935 topple). This difference was significant ($P \leq 0.001$). During the second time period (years 4-5
936 after harvest) the greatest change in the root-pit formation rates was a large increase in the rate
937 for the reference patches and a decrease in rates for the 50-ft buffers. The difference in rates
938 between the reference and the 50-foot buffer were not significant for this time period. The clear-
939 cut patches continued to have the lowest rate and were still significantly lower than the reference
940 patches ($P \leq 0.001$). Over the entire first five years, the rate of total root-pit formation for the 50-
941 ft buffers was nearly double the reference rate, however, this difference was not significant. The

942 pattern was similar for root-pits with sediment delivery, however the difference between the
943 reference and buffer patches was less pronounced due to the higher percentage of root-pits
944 delivering sediment in the reference patches. The percentage of root-pits with evidence of
945 sediment delivery was much higher in the clear-cut patches than in the 50-ft buffers (20.1%) and
946 the reference (26.0%) patches but was not significantly different. Results for the PIP buffers
947 showed a similar trend as the 50-foot buffers with an increase in root pits delivering sediment to
948 the stream in the first three years, but a sharp decline after the third year. Over the course of the
949 full five years Over the entire 5 year period, the percentage of root-pits with evidence of
950 sediment delivery in the PIP buffers (17.6%) was similar to the percentage for the 50-ft buffers
951 (19.8%). These values did not differ significantly from the references.

952 The authors also investigated the factors affecting whether the post-harvest root pits delivered
953 sediment to streams for 2006 and 2008 (3 and 5 years post-harvest). In both years, sediment
954 delivery to streams was best predicted by the distance of the root-pit from the stream ($P <$
955 0.0001). Mean horizontal distance to the stream for root-pits that delivered sediment was 8.2 ft
956 compared to 28.0 ft for those that did not deliver. Using horizontal distance to stream, the
957 proportion of the total uncertainty that was attributed to the model fit was 0.39, and 80% of the
958 observations were correctly predicted based on whether the horizontal distance to stream was
959 greater or less than 12.5 ft. Width of root pits delivering soil to the stream were also larger on
960 average but its inclusion to the model did not increase fitness. The authors speculate from their
961 observations that the higher tree-fall rates in the 50-foot buffer during the first 3 years after
962 harvest was due to an increase in wind-throw. However, in the second time period the reference
963 patches showed an increase in windthrow following stronger storms during the 2006-2008
964 period. One of the two reference streams did show string evidence of mass wasting.

965 Ehinger et al. (2021; Soft Rock Study) in their investigation of sediment export following harvest
966 along Type Np streams in western Washington (same prescriptions as described above for
967 Schuett-Hames, 2011) also quantified bank erosion events to assess sediment source. To assess
968 erosion events, the researchers placed two eye screws outside of the bank full width to attach a
969 reel tape for measuring length and depth across the bank. No evidence of bank erosion events
970 were found during the pre-harvest periods (1-2 years depending on site) for any stream reach. No
971 erosion events were found at any of the treatment sites during the post-harvest period (3-4 years
972 depending on site). However, there were observations of sediment being sourced from root-pits
973 developed in 2 treatment sites during the post-harvest period, but these effects were not
974 statistically analyzed. Because of the large mass wasting event in the reference the data collected
975 does not support any strong conclusion about the effect of riparian timber harvest on bank
976 stability.

977 McIntyre et al. (2018; Hard Rock Study) also investigated post-harvest surface erosion following
978 harvest along Type Np streams (~~same prescriptions as~~[including the FP treatment as investigated](#)
979 [by](#) Schuett-Hames, 2011) on competent lithologies in western Washington. They conducted
980 visual surveys to identify recently eroded areas (source of erosion not discerned) in the treated
981 riparian areas that were 10 m² or larger. Post-harvest stream-delivering surface erosion was
982 documented at 11 of 17 sites observed. The total erosion area exceeded 110 m² at 5 of the 17

983 sites: 2 reference sites, 2 50-foot buffer sites, and 1 clearcut sites. At these five sites, post-harvest
984 surface erosion was evident adjacent to only 1.5 to 4.6% (average = 2.2%) of the total stream
985 channel length (including both mainstem and tributaries). At the remaining study sites where
986 stream-delivering erosion events occurred, the total eroded area was 60 m² or less and occurred
987 adjacent to 0.3% to 0.8% (average = 0.6%) of the stream channel length. There were no
988 statistically significant differences in stream-delivering surface erosion among treatments ($\alpha =$
989 0.05), and on average, reference and buffer treatments visually exhibited a similar amount of
990 exposed bank.

991 The researchers also investigated the frequency of uprooted trees that developed root pits during
992 the post-harvest period. The average rate of root pits developed in the 50-foot buffers was
993 approximately 3 times higher (3.6 pits/ha/yr) than in the reference sites (1.2 pits/ha/yr) for 3
994 years following harvest. However, year to year values were highly variable with reference sites
995 showing higher numbers of root pits per acre than either buffer treatment in the first year
996 following treatment (27.4 vs. 18.5 vs. 6.4 for reference, 50-foot, and clearcuts respectively).

997 The results of the above studies on bank and riparian surface erosion after harvest show some
998 evidence that bank erosion and soil disturbance is generally higher in treated areas than in
999 untreated areas. Further, that bank erosion is likely higher in clearcut treatments without buffers
1000 than in treatments with no-cut buffers. However, development of root-pits (with and without
1001 sediment delivery pathways to streams) are more likely in treatments with no-cut buffers which
1002 is likely because no trees were left in the clearcuts to be toppled. When compared to a reference,
1003 the trends of surface erosion and soil disturbance shows there is generally an increase in the
1004 treated buffers within the first few years. However, these differences appear to stabilize within
1005 five years. Finally, soil disturbance and bank erosion (especially when caused by windthrow) are
1006 highly variable and in many instances ([e.g., Ehinger et al. 2021; McIntyre et al. 2018](#)) do not
1007 exceed the natural range of variability found in reference streams ([e.g., Ehinger et al. 2021;](#)
1008 [McIntyre et al. 2018](#)).

1009 *Sediment*

1010 The function of riparian areas to regulate and filter the flow of sediments into streams is essential
1011 not only for water clarity and pool formation but also because of the ability of sediments to carry
1012 nutrients and pollutants (Cooper et al., 1987; Hoffman et al., 2009; Polyakov et al., 2005). .
1013 Sediment flux into streams can be affected by landscape factors, streamflow, vegetation
1014 composition, and disturbance including riparian and adjacent upland forest management
1015 (Crandall et al., 2021; Devotta et al., 2021; Vanderbilt et al., 2003). The movement of sediment
1016 into the active channel can, in turn, impact aquatic habitat and geomorphic processes, especially
1017 in small, forested streams (Benda et al. 2005; Gomi et al., 2005; Hassan et al., 2005).

1018 The effects of riparian area timber harvest on sediment flux into streams has been documented,
1019 investigated, and incorporated into riparian forest management plans in western North America
1020 since the 1970s with the development of the Clean Water Act of 1972 (Bilby et al., 1989;
1021 Gregory 1990; Gresswell et al., 1989; Naiman et al., 1998; Salo & Cundy, 1986; Swanson et al.,
1022 1982; Swanson & Dyrness, 1975). Prior to the Forests and Fish Report (FFR 1999), several

1023 studies from western North America investigated the effects of riparian zone timber harvest
1024 practices on sediment flux into streams.

1025 Specific to Washington, Rashin et al. (2006) evaluated the effectiveness of Washington State best
1026 management practices (BMPs) for controlling sediment related water quality impacts. Although
1027 this study was published in 2006, the data analyzed in this study were collected between 1992
1028 and 1995. In their evaluation, Rashin et al. (2006) assessed site erosion, sediment delivery,
1029 channel disturbance, and aquatic habitat condition within the first two years of harvest along
1030 fish- and ~~non-fish-bearing~~ streams across Washington state. From their results,
1031 the authors concluded that the site-specific factors influencing the effectiveness of BMPs in
1032 preventing chronic sediment delivery into streams were 1) the proximity of ground disturbance
1033 to the stream, 2) presence of a stream buffer, 3) falling and yarding practices that minimized
1034 disturbance to stream channel, and 4) timing of harvest activities for certain climate zones where
1035 frozen ground or snow cover may be exploited. The landscape factors that influenced BMP
1036 effectiveness were 1) the density (specific metric not reported) of unbuffered small streams at
1037 harvest sites, and 2) steepness of stream valley slopes. The authors conclude with a
1038 recommendation of excluding timber falling and yarding activities at least 10 m from streams
1039 and outside of steep inner gorges.

1040 Similar results were reported by Lewis (1998) in their evaluation of logging activities' effect on
1041 erosion and suspended sediment transport in the Caspar Creek Watersheds of northwestern
1042 California. From their results the authors concluded that the dominant factors influencing the
1043 difference in suspended sediment loads between watersheds was the difference in road
1044 alignment, yarding methods, and presence of stream protection zones (i.e., buffers). Because of
1045 studies like these reviewed, contemporary riparian forest management practices in the western
1046 United States include rules that limit harvesting, use of equipment, and procedures that disturb
1047 soil in areas closest to the stream or on steep and unstable slopes ([WAC 222-30-022](#); WAC 22-
1048 30-021; 2022 [ODE](#); IDAPA 20.02.01)

1049 Since 2000, many of the studies published that evaluate changes in sediment delivery or water
1050 turbidity following riparian timber harvest show similar results in that contemporary BMPs are
1051 effective in mitigating increases in sediment delivery to streams (Hatten et al., 2018; Reiter et al.,
1052 2009). For example, the studies reviewed that report a significant change in sediment delivery
1053 following harvest show evidence that these changes only persist for a short period of time (1-3
1054 years) and that the magnitude of these changes are related to the intensity of the harvest
1055 prescriptions (Karwan et al., 2007; Macdonald et al., 2003a).

1056
1057 For example, Macdonald et al. (2003a) compared changes in stream discharge rates and in-
1058 stream suspended sediment concentrations during spring snowmelt between two harvest
1059 intensities and one unharvested control, for pre- and post-harvest in first order streams of interior
1060 British Columbia. Both treated riparian areas received a harvest of 55% of the watershed; one
1061 (low-retention) removed all merchantable timber >15 cm DBH for pine and > 20 cm DBH for
1062 spruce within 20 m of the stream; the other (high-retention) removed all merchantable timber >
1063 30 cm within 20 m of the stream. The results showed an increase in spring snowmelt discharge

1064 for both treatments above predicted values for the study (5 years). However, increased in-stream
1065 total suspended sediments (TSS) only persisted for two-years post-harvest in the high-retention
1066 treatment, and for 3-years in the low-retention.

1067
1068 Karwan et al. (2007) investigated the effects of riparian timber harvest and road construction on
1069 TSS concentrations in the Mica Creek Experimental Watershed in northern Idaho. Treatments in
1070 the paired-watershed experiment consisted of 1) commercial clearcut of the watershed area by
1071 50%, and was broadcast burned and replanted, 2) partial cut in which half the canopy was
1072 removed in 50% of the watershed area 3) a no-harvest control. All harvests were done according
1073 to best management practices and the Idaho Forest Practices Act. This included equipment
1074 exclusion zones of 50- and 30-feet for fish- and non-fish-bearing streams, respectively. On all
1075 skid trails, drainage features, such as water bars, were installed for erosion control at the end of
1076 the harvest period. Results showed that road construction in both watersheds did not result in
1077 significant impacts on monthly sediment loads in either treated watershed during the immediate
1078 (1-year post-harvest) or recovery (2-4 years post-harvest) time intervals. A significant and
1079 immediate impact of harvest on monthly sediment loads in the clear-cut watershed ($p = 0.00011$),
1080 and a marginally significant impact of harvest on monthly sediment loads in the partial cut ($p =$
1081 0.081) were observed. However, after one year, the TSS loads in both treatments became
1082 statistically indistinguishable from the control.

1083
1084 Specific to Washington, McIntyre et al. (2021) evaluated the effectiveness of riparian buffers on
1085 non-fish-bearing streams underlain by competent lithologies (“Hard Rock”) in western
1086 Washington. [Riparian management zones \(RMZs\) Buffers](#) were treated with one of ~~three~~
1087 ~~prescriptions~~ [four treatments including three harvest prescriptions \(\(-1\) unharvested reference, -2\)](#)
1088 [a minimum](#) two-sided 50-ft riparian buffer along the entire ~~riparian management zone (RMZ),~~
1089 [\(-2\) a minimum](#) two-sided 50-ft riparian buffer along at least 50% of the RMZ, and ~~4(3)~~
1090 [clearcut to stream edge \(no- buffer\), and an unharvested reference](#). Results for suspended sediment
1091 export (SSE) following treatment showed episodic increases with storm events that rapidly
1092 declined. However, changes in SSE were poorly correlated with discharge and exhibited high
1093 variation between treatment sites. The authors suggest that these results show evidence that
1094 changes in SSE magnitudes were not related to harvest. Further, they conclude that the sites were
1095 likely sediment-limited considering the underlying lithology.

1096
1097 Site factors such as underlying lithology and physiography can interact with the effect of timber
1098 harvest operations on sediment delivery into streams. Bywater-Reyes et al. (2017) assessed the
1099 influence of natural controls (basin lithology and physiography) and forest management on
1100 suspended sediment yields in temperate headwater catchments in northeastern Oregon. Results
1101 from this study indicate that site lithology was the first order control over suspended sediment
1102 yield (SSY) with SSY varying by an order of magnitude across lithologies observed.
1103 Specifically, SSY was greater in catchments underlain by Siletz Volcanics ($r = 0.6$), the Trask
1104 River Formation ($r = 0.4$), and landslide deposits ($r = 0.9$) and displayed an exponential
1105 relationship when plotted against the percentage of watershed area underlain by these lithologies.
1106 In contrast, lithology had a strong negative correlation with percent area underlain by diabase (r

1107 = 0.7), with the lowest SSY associated with 100% diabase. Following timber harvest, increases
1108 in SSY occurred in all harvested catchments but returned to pre-harvest levels within 1 year
1109 except for sites that were underlain by sedimentary formations and were clearcut without
1110 protective buffers. The authors conclude that sites underlain with a friable lithology (e.g.,
1111 sedimentary formations) had, on average, SSYs an order of magnitude higher following harvest
1112 than those on more resistant lithologies (intrusive rocks).

1113 Mueller & Pitlick, (2013) found similar results in their assessment of the relative effect of
1114 lithology, basin relief, mean basin slope, and drainage density on in stream sediment supply for
1115 83 drainage basins in Idaho and Wyoming. The strongest correlation of in stream sediment
1116 supply was with lithology relative softness (based on grouping of rock types – granitic,
1117 metasedimentary, volcanic, and sedimentary). Sediment concentrations at bankfull width
1118 increased by as much as 100-fold as basin lithology became dominated by softer sedimentary
1119 and volcanic rock compared to lithologies dominated by harder granitic and metasedimentary
1120 rock. Finally, Wissmar et al. (2004), developed and field-tested erosion risk indices for
1121 watersheds in western Washington based on land cover. These erosion risk indices used the
1122 presence of unstable soils (determined by geological formation and underlying lithology), rain-
1123 on-snow events, immature forest cover (stands <35 years old where open canopies and
1124 undeveloped root systems could contribute to hillslope instability), presence and coverage of
1125 roads, and critical slope (hillslope gradients >36%, for terrain with surficial deposits of coarse-
1126 textured colluvial materials). Results of this study showed these variables could explain ~65% of
1127 the variation associated with sediment input into channels. The lowest risk areas contained the
1128 fewest of these variables (most commonly critical slope with either rain-on snow events or
1129 immature forests), while higher risk areas contained a combination of 4 or more of these factors
1130 indicating a compounding effect.

1131 Changes in sediment yield may also interact with increases in discharge rates caused by timber
1132 harvest as well as physiographic site factors. For example, Bywater-Reyes et al. (2018)
1133 quantified how sediment yields vary with catchment lithology and physiography, discharge, and
1134 disturbance history over 60 years in the H.J. Andrews experimental watershed in the western
1135 Cascade Range of Oregon. Methods for determining suspended sediment concentration involved
1136 using either vertically integrated storm-based grab samples, or discharge-proportional composite
1137 samples where composite samples were collected every three weeks at the outlet of each
1138 catchment. Data sets were taken from 10 watersheds, 7 with a history of management (mixture of
1139 selective canopy removal, patch-cut, 25-100% clearcut, broadcast burning, road building, and
1140 thinning), and 3 with no history of management that were used as a reference. A linear mixed
1141 effects model (log transformed to meet the normality assumption) was used to predict annual
1142 sediment yield. In this model, site was treated as a random effect while discharge and
1143 physiographic variables were treated as fixed variables. This allowed for the evaluation of the
1144 relationships between sediment yield and physiographic features (slope, elevation, roughness,
1145 and index of sediment connectivity) while accounting for site. To account for the effect of
1146 disturbance history a variable was added to the model when the watershed had a history of
1147 management or natural disturbances. If the models for the disturbed watersheds significantly
1148 underpredicted the sediment discharge, the timing of the sudden increases were further examined

1149 to assess whether it correlated with a disturbance event (e.g., harvesting, road building, and
1150 slash-burning.) The results of this study show that watershed physiography combined with
1151 cumulative annual discharge explains 67% of the variation in annual sediment yield across the
1152 60-year data set regardless of lithology. Relative to other physiographic variables, watershed
1153 slope was the greatest predictor of annual suspended sediment yield. However, the results
1154 showed that annual sediment yields also moderately correlated with many other physiographic
1155 variables and caution that the strong relationship with watershed slope is likely a proxy for many
1156 processes, encompassing multiple catchment characteristics.

1157 In contrast, Safeeq et al. (2020) compared instream and bedload sediment supply under multiple
1158 harvesting treatments in watersheds of western Oregon that were paired with control watersheds
1159 by size, aspect, and topography. The treatment watershed was 100% clearcut during the period
1160 from 1962-1966, broadcast burned in 1966, and re-seeded in 1968. For this study 15-minute
1161 streamflow data was recorded for both watersheds, and after large storm events. Sediment data
1162 was collected from 1952 (pre-harvest) through 1988 for suspended sediment data, and 2016 for
1163 sediment bedload. The control watershed was forested, and had no treatments (e.g., harvest)
1164 during the study period. Their results estimate that following streamside harvest, increased
1165 streamflow alone is estimated to be responsible for <10% of sediment transport into streams
1166 while the increased sediment supply caused by harvest operations is responsible for >90% of the
1167 sediment transported into streams.

1168 Puntenney-Desmond et al. (2020) found similar results in their assessment of differences in
1169 instream sediment contributions from the buffer area, harvest area, and buffer-harvest interface.
1170 Sediment concentration in the runoff was 15.8 times higher for the harvested area than in the
1171 riparian buffer, and 4.2 times greater than in the harvest-buffer interface. Total sediment yields
1172 ($\text{mg m}^{-2} \text{min}^{-1}$) from the harvested area (sediment concentration x flow rate) were approximately
1173 2 times greater than in the buffer areas, and 1.2 times greater in the harvest-buffer interface than
1174 in the buffer area.

1175 *Summary of Factors Impacting Sediment Delivery into Streams*

1176 From the studies reviewed there is evidence that sediment delivery into streams following timber
1177 harvest is influenced by not only the intensity of the harvest operation (e.g., presence of retention
1178 buffers, yarding and equipment use immediately adjacent to the stream, upland clearcut vs.
1179 thinning), but also by physiography (e.g., hillslope gradient), lithology relative softness, and
1180 climate (e.g., precipitation, frequency of large storm events). Thus, the change in magnitude of
1181 sediment delivery following harvest is context dependent and these landscape factors can interact
1182 with one another to compound these changes. However, from the studies reviewed above there is
1183 evidence that the implementation of BMPs since the 1970s in the northwestern United States
1184 lessen the impact and duration of these changes.

1186 [Shade and stream temperature](#)

1187 Canopy cover provides shade for streams that decreases the amount of incoming solar radiation
1188 and thus influences stream temperatures, although that influence can be highly variable
1189 depending on shade structure and density surrounding stream courses. Temperature regulation is
1190 vital for sensitive salmonid fish species that require cooler waters, and shade is often the primary
1191 function assessed when developing state regulations (Groom et al., 2011; Groom et al., 2018;
1192 Teply et al., 2014). The importance of shade and cooler in-stream temperatures for fish habitat
1193 has been thoroughly investigated (Bjornn & Reiser, 1991; Chapman & Bjornn, 1969; Ebersole et
1194 al., 2001; Sullivan et al., 2000). The streamside shade will likely become even more critical with
1195 the predicted increases in air temperature over the next century (Manuta et al., 2009). While
1196 stream temperature is initially reflective of moisture source (e.g., snowmelt, liquid precipitation,
1197 groundwater inputs) and watershed subsurface soil characteristics. As water flows downstream
1198 and into higher-order streams, the net rate of temperature gain or loss is the sum of incident
1199 radiation, evaporation, conduction, and advection (Brown, 1983; Bescheta et al., 1987).

1200 Bescheta et al. (1987) presented evidence that direct beam solar radiation inputs are of the
1201 highest importance to the stream's net heat exchange rate per unit area compared to other factors.
1202 Within the net heat exchange calculation, the heat released from evaporation generally cancels
1203 out the heat gained from warm air temperatures (convective and advective heat transfer). Thus,
1204 temperature fluctuations are expected to be more severe in less-shaded/more-exposed streams.
1205 This has been supported by many experimental field and simulation studies showing evidence
1206 that the reduction of effective shade can lead to considerable increases in peak summer stream
1207 temperatures primarily due to the increase of incoming solar radiation. However, while increases
1208 in solar radiation are accepted as the most important factor in stream temperature changes and
1209 fluctuations following harvest, other factors are also important and may compound these effects.

1210 Guenther et al. (2014) investigated the relationship between changes in stream temperature and
1211 changes in wind speed, vapor pressure, and evaporation following riparian thinning treatments
1212 along headwater streams in southwestern British Columbia. Treatment involved reduction of
1213 basal area by 50% (resulting in 14% reduction in canopy closure) in the upland and riparian
1214 forests. Results showed a post-harvest increase in wind speed, vapor pressure deficit, air
1215 temperature and evaporation above the stream, which coincided with increased stream
1216 temperatures and lower stability. The authors report that prior to harvest, vapor pressure
1217 gradients often favored condensation over evaporation. Further, they concluded that the
1218 relationships between the riparian and microclimate variables after harvesting became more
1219 strongly coupled to ambient climatic conditions due to increased ventilation. Contemporary
1220 riparian management practices in western North America vary by state. However, all require
1221 retention of protective buffers that preserve some percentage of shade or canopy cover to
1222 maintain or mitigate changes in stream temperatures, especially along fish-bearing streams.
1223 Many studies published in the last two decades report evidence that these practices have been
1224 effective in mitigating stream temperature changes after harvest.

1225 For example, Bladon et al. (2016), assessed the effectiveness of riparian management
1226 prescriptions developed for the Oregon Forest Practices Act (FPA). Oregon State requires a 15 m

1227 buffer on either side of small fish-bearing streams with a 6 m no-cut buffer, and a minimum
1228 retention for conifer basal area of ~3.7 m² for every 300 m (~1000 ft) length of stream. This
1229 resulted in a reduction of mean canopy closure from ~96% in the pre-harvest period to ~89% in
1230 the post-harvest period in the treatment reaches. In contrast, mean canopy closure in the
1231 reference reaches changed from ~92% to ~91% from pre- to post-treatment periods. Results
1232 showed there was a significant increase in the 7-day moving maximum temperature from pre- to
1233 post-harvest values when data was constrained to the period of July 15 – August 15 by 0.6 +/-
1234 0.2 °C. However, when analyzed by individually paired sites, and when interannual and site
1235 variability was accounted for, no significant changes in stream temperature were observed for 3
1236 years post-harvest (length of study).

1237 However, Groom et al., (2011a, b) showed evidence that the more stringent rules of the
1238 Northwest Oregon State Forest Management Plan (FMP; applied to riparian management zones
1239 on state owned land) was even more effective at maintaining stream temperatures post-harvest.
1240 The FMP requires a 52 m wide buffer for all fish-bearing streams, with an 8 m no cut buffer
1241 immediately adjacent to the stream. The results from Groom et al. (2011b) showed that FPA
1242 (Oregon Forest Practices) post-harvest shade values differed from pre-harvest values (mean
1243 change in Shade from 85% to 78%), while no difference was found for FMP shade values pre-
1244 harvest to post-harvest (mean change in Shade from 90% to 89%). Following harvest, maximum
1245 temperatures at FPA increased relative to FMP on average by 0.71 °C. Similarly, mean
1246 temperatures increased by 0.37 °C (range: 0.24 - 0.50), minimum temperatures by 0.13 °C
1247 (range: 0.03 - 0.23), and diel fluctuation increased by 0.58 °C (range: 0.41 - 0.75) relative to
1248 FMP sites.

1249 Groom et al (2011a) developed prediction models from this data to estimate the probability of
1250 riparian harvest under each regulation causing an increase in stream temperatures >0.3 °C (the
1251 Protecting Cold Water criterion developed by the Department of Environmental Quality). Results
1252 indicate that sites harvested according to FPA standards exhibited a 40.1% probability of a
1253 temperature change of > 0.3°C from pre- to post harvest. Conversely, harvest to FMP standards
1254 resulted in an 8.6% probability of exceedance that did not significantly differ from all other
1255 comparisons.

1256 In Montana, Sugden et al. (2019) investigated the effectiveness of state regulation which requires
1257 timber be retained within a minimum of 15.2 m (50 feet) of the stream. Within the riparian
1258 management zone, no more than half the trees greater than 204 mm (8 in) diameter at breast
1259 height (DBH) can be removed. In no case, however, can stocking levels of leave trees be reduced
1260 to less than 217 trees per hectare. Data for canopy cover, stream temperature, and fish population
1261 were collected for 30 harvest reaches in western Montana (northern Rocky Mountain Region),
1262 for a minimum of one-year pre- and one-year post-harvest. Shade over the stream surface was
1263 not directly measured in this study. Instead, canopy cover was used as proxy, using two
1264 independent estimates of canopy cover (1) used cruise data to populate a canopy cover model
1265 within Forest Vegetation Simulator, and (2) measured canopy cover in the harvested reach every
1266 30 m, before and after harvest. Within harvest units, mean basal area was reduced by 13%
1267 (range: 0 – 36%), and again further by a mean of 2% due to windthrow. Mean canopy cover

1268 within the riparian management area reduced from 77% (pre-treatment) to 74% (post-treatment),
1269 and mean canopy cover over the stream changed from 66% (pre-treatment) to 67% (post-
1270 treatment) based on densiometer measurements. Neither of these changes were significant.
1271 Results for stream temperature also showed no significant changes in stream temperatures or fish
1272 populations in one-year post treatment compared to pre-treatment values.

1273 Specific to Washington, Cupp & Lofgren (2014) conducted a study to test the effectiveness of
1274 riparian timber harvest rules for eastern Washington in preserving shade and stream
1275 temperatures. Regulations for fish-bearing streams in eastern Washington (in the mixed
1276 conifer/mid elevation zone) includes an “All Available Shade Rule” (ASR) for streams in the bull
1277 trout habitat zones, and a “Standard Shade Rule” (SR). Under the ASR it is required to retain all
1278 available shade within 75 feet of the stream. Under SR some harvest of shade providing trees is
1279 allowed within the 75-foot buffer depending on elevation and pre-harvest canopy cover.
1280 Unharvested reference reaches were located upstream from treatment reaches. Prior to harvest
1281 treatments, canopy closure measurements ranged from 89% to 97%, with a mean of 93%.
1282 Results showed post-harvest shade values decreased in SR sites (mean effect of -2.8%, $p =$
1283 0.002), as did the canopy closure values (mean effect of -4.5%, $p < 0.001$). Shade and canopy
1284 closure values did not significantly change after treatment in the ASR sites. Post-harvest mean
1285 daily maximum stream temperature increased 0.16 °C in the SR harvest reaches, whereas stream
1286 temperatures in both the ASR sites and in the no-harvest reference reaches increased on average
1287 by 0.02 °C. Sample period means of daily maximum temperature responses varied from -1.1 °C
1288 to 0.7 °C in the first two years post-harvest for the ASR sites, from -0.5 to 0.8 °C, in the SR
1289 sites, and -0.5 to 0.9 °C in the reference sites. While these values show a slight increase in mean
1290 temperatures and temperature ranges with treatment, the authors interpret these results as
1291 evidence that temperature effects of the SR and ASR were similar to reference conditions along
1292 sampled reaches.

1293 Riparian harvest rules along ~~non-fish-bearing~~ streams tend to allow for narrower
1294 buffer widths (sometimes with no retention buffers) or more intense thinning within the buffer
1295 than for fish-bearing streams. For example, in western Washington the Forest Practices (FP)
1296 buffer prescription requires a minimum two-sided 15 m (50 ft) wide buffer along a minimum of
1297 50% of the length of a non-fish-bearing perennial stream (i.e., up to 50% of the stream may have
1298 no buffer) with a 9.1 m (30 ft) equipment exclusion zone along the entire stream. Two recent
1299 studies (Ehinger et al., 2021 “Soft Rock”; McIntyre et al., 2021 “Hard Rock”) have compared
1300 these FP buffers, and for Hard Rock to two experimental alternative buffer treatments (~~a~~ 50 ft
1301 buffer along 100% of the stream length (100%), and no buffer (0%) treatment), and to an
1302 unharvested reference (REF) sites. Hard Rock on sites were underlain by competent lithologies
1303 and Soft Rock by (McIntyre et al., 2021; “Hard Rock”) or incompetent (friable easily eroded)
1304 lithologies. ~~(Ehinger et al. 2021; “Soft Rock).~~

1305 Results from the Hard Rock study showed that riparian canopy cover declined after harvest in all
1306 buffer treatments reaching a minimum around 4 years post-harvest (after mortality stabilized).
1307 The treatments, ranked from least to most change, were REF, 100%, FP, and 0% for all metrics
1308 ~~and~~ across all years. Effective shade results showed decreases of 11, 36, and 74 percent in the

1309 100%, FP, and 0% treatments, respectively. These changes in shade were significant for all
1310 treatments. This led to changes in mean stream temperature from pre- to post-harvest in the
1311 100% treatment by 1.1°C in the first two years following treatment, but returned to pre-
1312 harvest levels by post-harvest year 3. In contrast, the mean difference in pre- to post-harvest
1313 stream temperatures in the FP ranged from 0.5°C to 1.1°C in the first year, and changed little over
1314 the entire post-harvest period. Results for the 0% treatment showed a mean increase of 3.8°C
1315 immediately following harvest and declined over time to 0.8°C by post-harvest year 9. These
1316 results suggest that the 100% treatment was most effective at preventing increases to stream
1317 temperature followed by the FP and 0% treatments. A weak and nearly significant (P-value
1318 range: 0.008 - 0.108) negative relationship between canopy cover and stream temperature for the
1319 first 4 years after treatment was detected. These results provide evidence that the effectiveness of
1320 buffers in maintaining stream temperatures post-harvest is relative to the intensity of the
1321 treatment (e.g., presence of buffer, reduction in canopy cover). Further, post-treatment mortality
1322 within the buffer from events such as windthrow can cause fluctuations in stream temperature
1323 response during the first decade. Results from the Soft Rock Study showed similar trends in
1324 canopy cover reduction and stream temperature increases. Authors of the Soft Rock study note
1325 that stream temperature changes varied as a function of the proportion of the stream buffered and
1326 tree mortality.

1327 Outside of Washington, several studies conducted in western North America since 2000 have
1328 shown results similar to the Hard Rock and Soft Rock studies. For example, Roon et al. (2021b)
1329 compared stream temperature changes following variable riparian thinning intensities in the
1330 redwood forests of northern California. Treatments to riparian stands included reduction of
1331 canopy cover that resulted in reduction of effective shade by either (19-30%) or by (4-5%). Their
1332 results showed that local changes in stream temperature were dependent on thinning intensity,
1333 with higher levels of canopy cover reduction leading to increases in local stream temperatures. In
1334 the reaches with higher reductions in shade (19-30%) there was accumulation of 45° to 115°C
1335 additional degree days from pre- to post treatment years, while the reaches with lower reductions
1336 in shade (4-5%) only accumulated 10° to 15°C additional degree days. Further, travel distance of
1337 increased stream temperatures also appeared to be dependent on thinning intensity. The lower
1338 shade reduction reaches had an increased temperature effect downstream with travel distance of
1339 75-150 m, while the high shade reduction sites had a downstream travel distance of 300- ~1000
1340 m. Roon et al. (2021a) reported changes in in average daily maximum, maximum weekly
1341 average of the maximum (MWMT), average daily mean, or maximum weekly average of the
1342 mean (MWAT) at these same sites under the same timeline. The lower thinning intensity (4-5%
1343 effective shade reduction) showed no significant changes in any temperature metrics. However,
1344 The more intensely thinned sites (19-30% reduction in effective shade) showed an increase
1345 MWMT during spring by a mean of 1.7°C (0.9 - 2.5 °C), summer by a mean of 2.8°C (1.8, 3.8
1346 °C), and fall by a mean of 1.0°C (0.5, 1.5 °C) and increased in downstream reaches during spring
1347 by a mean of 1.0°C (0.0, 2.0 °C) and summer by a mean of 1.4°C (0.3, 2.6 °C). The authors
1348 interpret their results as evidence that that changes in shade of 5% or less caused minimal
1349 changes in temperature while reductions in shade of 20–30% resulted in much larger increases in
1350 temperature.

1351 Reiter et al. (2020) compared the changes in stream temperatures following different harvest
1352 treatments along headwater streams in the Trask River Watershed in the northwestern coast range
1353 of Oregon. Treatments included a clearcut to stream (no buffer but half of sites contained some
1354 leave trees along stream bank), upland clearcut with a 10 m no-cut buffer, upland thinning (basal
1355 area reduction to 30-50% of original stand) with a 10 m no-cut buffer, and an unharvested
1356 reference. Results showed that post-harvest stream temperature increases were only significant in
1357 the clear-cut treatments without buffers with a mean increase of 3.6°C (SE = 0.4°C) for four
1358 years after the study. They note that temperature changes were more severe in the unbuffered
1359 streams with no leave trees (4.2 and 4.4°C), however, this difference was not analyzed. No
1360 significant changes in stream temperature were detected in either treatment with a 10 m no-cut
1361 buffer. The authors speculate that 10 m wide buffers were sufficient in maintaining stream
1362 temperatures post-harvest in small, forested headwater streams.

1363 In the sub-boreal forest ecosystems of British Columbia, Canada, Macdonald et al. (2003b)
1364 compared pre- to post-harvest stream temperature changes in first-order headwater streams under
1365 3 different riparian forest treatments. These treatments included 1) low-retention – removal of all
1366 merchantable timber >15 or >20 cm DBH for pine or spruce respectively, within 20 m of the
1367 stream 2) high-retention – removal of merchantable timber >30 cm DBH within 20-30 m of the
1368 stream, and 3) patch-cut – high retention for the lower 60% of watershed approaching streams
1369 and removal of all vegetation in the upper 40% of the watershed. Results showed significant
1370 increase in stream temperatures ranging from 4 – 6 °C in the low-retention and patch cut in the
1371 first three years following harvest. However, by year five, mortality in the high-retention buffer
1372 (due to windthrow) resulted in canopy cover reduction and increases in stream temperatures that
1373 became equivalent to the other treatments. The authors conclude that while the variation in
1374 harvest intensity initially appeared to dictate stream temperature responses, site effects (e.g.,
1375 windthrow susceptibility) can impact the effectiveness of the buffer. While the studies above all
1376 show evidence that the impact of riparian forest harvest on stream temperatures are related to the
1377 severity of the harvest prescription (e.g., buffer width, thinning intensity, canopy reduction) the
1378 results are variable within treatments indicating other site factors are also important when
1379 evaluating buffer effectiveness. For example, in their review of experimental studies conducted
1380 in the Pacific Northwest of Canada and the United States, Martin et al. (2021) reported high
1381 variability in temperature response to streamside buffers. They report a substantial variability and
1382 overlap in the effect size of the mean 7-day maximum temperature metric with no-cut buffers,
1383 no-cut plus variable retention buffers, and no-cut patch buffers ≤ 20 m wide. The largest
1384 temperature response (> 3.4 °C) occurred in the clearcut buffers while treatments with buffers
1385 (i.e., no cut buffers without variable retention) had the smallest response (< 0 °C). The variable
1386 retention buffers < 20 m showed variable response (0.6 – 1.4 °C). They conclude that the
1387 variation in temperature response following riparian harvest may be associated with multiple
1388 factors such as geology, hydrology, topography, latitude, and stream azimuth.

1389 Bladon et al. (2018) investigated the changes in stream temperatures following treatments that
1390 varied from clearcuts to stream to buffers > 20 m in western Oregon. They performed a
1391 regression analysis to assess the relative relationship between catchment lithology and the
1392 percentage catchment harvested with stream temperature at all sites. Their results showed that at

1393 the upstream harvested sites there was a strong relationship between stream
1394 increases and catchment lithologies, but no statistically significant relationship between stream
1395 temperature changes and percent of catchment harvested. Sites downstream from harvested areas
1396 showed a significant relationship with the interaction of percentage of catchment harvested and
1397 the underlying lithologies ($p = 0.01$). The greatest temperature increases at downstream sites
1398 were in areas with a higher percentage of catchment harvested and were underlain by more
1399 resistant lithologies. There was no evidence for increases in stream temperatures in catchments
1400 with a high percentage of harvest that were underlain by permeable geology. The authors suggest
1401 that this relationship may be due to the buffering effect of increases in summer low flows and
1402 greater groundwater or hyporheic exchange. They conclude that the variability of rock
1403 permeability and the relative contribution of groundwater during summer months, and their
1404 effect on stream temperatures following harvest should be investigated further.

1405 There is evidence that geomorphology alone can impact stream temperature fluctuations
1406 throughout the year. Hunter & Quinn, (2009) compared seasonal fluctuations in stream
1407 temperatures between two watersheds in the Olympic Peninsula, Washington. Both watersheds
1408 were similar in all characteristics except for bed substrate. One was underlain by alluvial bed
1409 substrate while the other was underlain by bedrock. Results from this study show consistent
1410 differences in stream temperature response in alluvial versus bedrock channels. Seasonal
1411 maximum and minimum average daily temperatures varied less at the alluvial site compared to
1412 the bedrock site. This, the authors suggest, may be due to hyporheic exchange in alluvial
1413 channels helping to buffer surface water temperatures from gaining or losing heat. In addition,
1414 groundwater may also contribute to the increased stability at the alluvial site. Aside from shade
1415 reduction from timber harvest, there is evidence that light availability and canopy cover naturally
1416 changes over time as riparian stands develop. For example, Warren et al. (2013) compared
1417 canopy cover and stream light availability between old-growth-forests (>500 years old) and
1418 young harvest-aged stands (~40-60 years old) in the H.J. Andrews Experimental Forest in the
1419 Cascade mountains of Oregon. Streams were paired based on reach length and bankfull width,
1420 and north ($n = 2$), and south ($n = 2$) facing watersheds. Canopy cover was estimated using a
1421 convex spherical densiometer, and light reaching the stream bed was estimated using a
1422 fluorescent dye that degrades overtime from light exposure. Overall, three of the four paired old-
1423 growth reaches (2 south-facing, 1 north-facing) had significantly lower mean percent canopy
1424 cover ($p < 0.10$), and significantly higher mean decline in fluorescent dye concentrations ($p <$
1425 0.01). The authors interpret these results as evidence that old-growth forest canopies were more
1426 complex and had more frequent gaps allowing for more light availability and lower mean canopy
1427 cover, on average, than in adjacent young, second growth forests.

1428 Kaylor et al. (2017) presented similar results when they compared canopy cover and light
1429 availability between small mountain streams adjacent to late-successional forests (dominant
1430 canopy trees >300 years old) and second-growth forests that had been harvested to the stream
1431 50-60 years prior to data collection. Like Warren et al. (2013), canopy cover was estimated with
1432 a convex spherical densiometer; and light availability to streams was estimated with a
1433 photodegrading fluorescent dye. However, for this study, fluorescent dye degradation was
1434 converted to photosynthetically active radiation (PAR) by building a linear relationship between

1435 the dye degradation and PAR sensors. Results showed that mean PAR reaching streams was 1.7
1436 times greater, and canopy openness was 6.1% greater in >300-year-old forests than in 30–100-
1437 year-old forests. Of the 14 paired sites, differences in canopy openness and PAR were significant
1438 for 6 sites. The authors compared and combined their data with published data from 10 other
1439 similar studies. The combined datapoints for canopy openness (%) were plotted against stand age
1440 and fit it with a negative exponential curve. From the slope of the curve, the authors estimate that
1441 canopy openness reaches its minimum value in regenerating forests at ~30 years and maintains
1442 with little variability until ~100 years.

1443 *Summary of Factors Affecting Shade and Stream Temperature*

1444 From the studies reviewed above, the results show evidence that changes in canopy cover and
1445 effective shade are, not surprisingly, directly related to the intensity of harvest operation. Initial
1446 reduction in canopy cover and shade from pre- to post-harvest are influenced by the basal area
1447 removed and the width of the retention buffer. However, there is evidence that multiple site
1448 factors can interact with harvest operations (e.g., target basal areas).

1449

1450
1451

1452 Results/discussion by focal question

1453 Focal Question 1

1454 1. *What are the effects of timber harvest intensities and extent on the riparian functions, with an*
 1455 *emphasis on the five key functions listed above, in comparison to conditions before harvest?*

1456 From the perspective of an experimental design, this question inquires how the values of the
 1457 metrics used to describe the five key functions (large woody debris recruitment, sediment
 1458 filtration, stream bank stability, shade, litterfall and nutrients) differ from pre- to post-harvest
 1459 within particular riparian areas of interest. An attempt to answer this question would require data
 1460 collection before and after treatment with or without a control site. Thus, only studies that used a
 1461 BACI or BAI approach are appropriate for discussing this question. From our review, 22 papers
 1462 report pre- to post-harvest changes in the magnitude of one or more of the key functions with the
 1463 majority of these papers focusing on changes in shade. No studies published since 2000 that
 1464 apply an experimental design in western North America to quantify changes in bank stability
 1465 could be found in the literature.

Function	Count
Shade	12
Litter	3
LW	2
Sediment	4
Nutrients	3
Bank Stability	0

1466

1467 *Shade*

1468 Table 4. Treatment and responses for selected publications investigating shade relevant to Q1.

Reference	Treatment	Response
Bladon et al. (2016)	Buffer width of 15 meters (~50 feet)	3 years post-harvest (n = 6) Mean canopy closure was reduced from ~96% (pre-harvest) to ~89% (post-harvest).
Cupp & Lofgren (2014)	Buffer width of 75 feet	<u>1-2 years post-harvest</u> *ASR: Of 16 sites, 13 showed a decrease in shade ranging from 1 to 4%. 2 sites showed no change and 1 site showed an increase in shade of 4% (mean decrease of 1%). **SR: Of 14 sites, 13 showed a decrease in shade ranging from 1 to 10%, and 1 site showed an increase of 1% (mean decrease of 4%).
Gravelle & Link (2007)	Clearcut to stream; Thinning to 50% canopy cover	<u>1- and 2-years post-harvest</u> Pre-harvest shade ranged from 56% to 88% with a mean of 70% in control reaches (n = 4), 63% in clearcut reaches (n = 2), and 74% in thinned reaches (n = 2). In the clearcut reaches, post-harvest shade was reduced to a mean of 52% and 41% for years one and 2, respectively. In the thinned reaches, post-harvest shade remained near 75% for years 1 and 2.

Groom et al. (2011b)	Buffer width of 21 meters (~69 feet; Private); 52 meters (~170 feet; State)	<u>1 year post-harvest</u> For private sites (n = 18) mean post-harvest shade values decreased significantly from 85% to 78%; No statistical difference was found for state site (n = 15) mean shade values from pre- to post-harvest (90% to 89%).
McIntyre et al. (2021)	Buffer widths of ≥50 feet (100%); ≥50 feet with clearcut gaps (FP); Clearcut to stream (unbuffered)	<u>9 years post-harvest</u> Riparian cover declined after harvest in all buffer treatments reaching a <u>minimum at 4 years</u> . 100% buffers (n=4) showed a change in mean shade ranging from +1 to -10 % over nine years <u>post-harvest</u> . FP buffers (n = 43) showed a change in mean shade ranging from -12 to -32% over nine years. The unbuffered sites (n=4) showed a change in shade ranging from -27 to -87% over nine years. The 100% buffer recovered to pre-harvest values by year 9.
Reiter et al. (2020)	Clearcut, no buffer (CC_NB), clearcut with 10-m no cut buffer (CC_B), thinning with 10 m no-cut buffer (TH_B).	<u>1 year post-harvest</u> The CC_NB (n = 3) showed a reduction in shade ranging from 1.8 to 3.2% (mean = 2.4%). The CC_B treatment (n=3) showed a reduction ranging from 18.6 to 76.6% (mean = 56.6%). The TH_B (n = 1) showed an increase in shade of 2.8%.
Roon et al. (2021a)	Buffer width of 45 meters (~150 feet)	<u>1 year post-harvest</u> Of the two watersheds surveyed one showed a significant reduction in mean shade and canopy closure of 18.7 and 23.0%, respectively. The second showed a non-significant reduction of mean shade and canopy closure by 4.1 and 1.9%.
Sugden et al. (2019)	Buffer width of 15.2 meters (~50 feet)	<u>1 year post-harvest</u> <u>Mean post-harvest canopy cover increased by 1% (n = 28; range = -48 to +17%) measured with a densiometer. ***FVS Modeling based on tree metrics estimated a mean reduction in shade of 4.1% from pre- to post-harvest</u>
Swartz et al. (2020)	Buffer width of ~20-meter (65 feet) diameter gaps along streambank	<u>1-2 years post-harvest</u> Treatment reach (n = 4) mean shading declined by only 4% (SD ± 0.02%) post-harvest.

1469

1470 Specific to fish-bearing streams of eastern Washington, Cupp & Lofgren (2014) reported changes
1471 in canopy closure (quantified with handheld densiometer) and shade (quantified with fisheye lens
1472 digital camera) within reaches adjacent to riparian forests harvested under the All Available
1473 Shade Rule (ASR) and the Standard Shade Rule (SR). Both shade rules have a 30-ft no-cut
1474 buffer (core zone) immediately adjacent to the stream. The SR prescription allows thinning in the
1475 buffer zone 30-75 feet (inner zone) from the stream while the AAS prescription requires
1476 retention of all shade providing trees in this area. Results showed post-harvest shade values
1477 decreased in SR sites (mean effect of -2.8%, p = 0.002), as did the canopy closure values (mean
1478 effect of -4.5%, p < 0.001). Shade and canopy closure values did not significantly change in the
1479 treatment reaches of the ASR sites. Mean shade reduction in the SR treatment sites exceeded the

1480 mean shade reduction in the ASR sites by 3%. Canopy closure reduction was also greater in the
1481 SR sites than in the ASR sites by a mean of 4%.

1482 For ~~non-fish-bearing~~non-fish-bearing streams of western Washington, McIntyre et al. (2021)
1483 report changes in canopy closure following ~~3-three~~ different ~~harvest-riparian buffer~~ prescriptions.
1484 Riparian buffer Pprescriptions included a minimum two-sided 50-ft wide riparian buffer along
1485 the entire stream (100%), a minimum two-sided 50-ft riparian buffer along at least 50% of the
1486 stream consistent with the current Forest Practices buffer prescription (FP), and a clearcut to
1487 stream edge without a buffer (0%). The canopy cover was estimated at mid-stream with a
1488 handheld densiometer and was converted to effective shade values (for the 5 years post-harvest).
1489 Hemispherical canopy photos were also taken for 24 years pre-harvest and 53 years post-harvest
1490 and converted to Canopy and Topographic Density (percentage of the photograph obscured by
1491 vegetation or topography). Results for canopy cover showed that riparian cover declined after
1492 harvest in all buffer treatments reaching a minimum around 4 years post-harvest. The treatments,
1493 ranked from least to most change, were 100%, FP, and 0% for all metrics and across all years.
1494 Effective shade results showed decreases of 11, 36, and 74 percent in the 100%, FP, and 0%
1495 treatments, respectively, by 3 years post-harvest. However, by post-harvest year 9, canopy
1496 closure returned to pre-harvest levels in the 100% treatment but remained 15% and 27% below
1497 pre-harvest values at the FP and 0% treatments, respectively. —Another study, Janisch et al.
1498 (2012) also compared the effects of similar treatments (clearcut to stream, a full continuous
1499 buffer (10-15 m wide), and a patched buffer (~50-110 m long were retained in distinct patches
1500 along some portion of the channel) to canopy cover. Canopy cover in all streams averaged 95%
1501 (SE = 0.4) prior to harvest. Following treatment, canopy cover in the clearcut catchments
1502 averaged 53%, (SE = 7.4) canopy cover in the patch buffer treatment averaged 76%, (SE = 5.1)
1503 and canopy cover in the continuous buffer treatment averaged 86% (SE = 1.7). The changes were
1504 significant in the clearcut and patch buffers.
1505

1506 Outside of Washington, Bladon et al. (2016) assessed the effects of harvest treatments under the
1507 Oregon Forest Practices Act (FPA) on shade reduction and stream temperature. This study took
1508 place in the Siuslaw National Forest in the Oregon Coast Range in the Alsea Watershed.
1509 Treatment under the FPA includes a 15 m riparian management area with a minimum of ~3.7 m²
1510 conifer basal area retained for every 300 m length of stream and an additional 4-5 wildlife leave
1511 trees per hectare. This resulted in a mean canopy closure reduction from ~96% (pre-harvest) to
1512 ~89% (post-harvest) based on measurements from a densiometer along the stream channel for 3
1513 years pre- and 3 years post-harvest. Unfortunately, the authors did not compare these changes
1514 with statistical analysis. Groom et al. (2011b) compared changes in shade from pre- to post-
1515 harvest under the FPA and under the Northwest Oregon State Forest Management Plan (FMP).
1516 The FMP requires a 52 m wide buffer for all fish-bearing streams, with an 8 m no cut buffer
1517 immediately adjacent to the stream.

1518 Results from Groom et al. (2011b) showed that FPA site post-harvest shade values differed from
1519 pre-harvest values (mean change in Shade from 85% to 78%); While no difference was found for
1520 FMP site shade values pre-harvest to post-harvest (mean change in Shade from 90% to 89%). In
1521 the Trask Watershed of the northwestern Oregon Coast range, Reiter et al. (2020) compared three
1522 riparian zone treatments: 1) clearcut, no buffer (CC_NB; n = 4), 2) clearcut with 10-m no cut

1523 buffer (CC_B; n = 3), 3) thinning with 10 m no-cut buffer (TH_B; n = 1) in small [non-fish](#)
 1524 [bearingnon-fish-bearing](#) streams. Pre- to post-harvest values in shade were quantified with
 1525 hemispherical analysis over the stream one-year prior and one-year post-treatment. However,
 1526 post-harvest overstory buffer width varied within each treatment depending on landscape factors.
 1527 For this reason, we will present the change in percent shade with residual buffer width (Table 6).
 1528 Again, changes in shade were not statistically analyzed.

1529 In fish-bearing streams within the McKenzie River basin in the western Cascade Mountains of
 1530 Oregon Swartz et al. (2020) assessed the effects of experimental canopy gap treatments on shade
 1531 and light availability to the stream. In each treatment reach (n = 6), 20 m gaps were prescribed to
 1532 mimic gap openings that naturally occur after individual large tree mortality or small-scale
 1533 disturbance events in late successional forests. Shade was recorded in the year before and the
 1534 year after treatment with hemispherical photos. Changes in effective shade (1 year post-harvest)
 1535 were estimated in HemiView 2.1 software. Mean stream shading could not be evaluated in the
 1536 full BACI analysis because post-treatment hemispherical photographs could not be taken at all
 1537 sites due to fire impeding access in 2018. For the remaining sites, the areas beneath each gap had
 1538 notable localized declines in shade, through the entirety of the treatment reach mean shading
 1539 declined by only 4% (SD ± 0.02%).

1540 Table 5. Results for changes in shade following treatment for the Trask River Watershed Study
 1541 headwaters. Reproduced from Reiter et al (2020).

Treatment	Mean residual buffer width (2-sided)	Pre-harvest shade (%)	Post-harvest shade (%)
CC_B	33.2	85.9	82.7
CC_B	22.6	91.3	89.1
CC_B	23.9	84.7	82.9
CC_NB	0.0	83.6	7.0
CC_NB	0.0	85.5	10.9
CC_NB	16.0	84.3	65.7
CC_NB	14.1	80.6	76.6
TH_B	*	81.2	84.0

1542 CC_B = clearcut with 10 m buffer, CC_NB = clearcut no buffer, TH_B upland thinning with
 1543 buffer. *Unable to determine exact buffer width because adjacent to thinning

1544 Gravelle & Link (2007) compared changes in shade following treatment for [non-fish-bearingnon-](#)
 1545 [fish-bearing](#) streams in northern Idaho. For non-fish-bearing streams there is a 30 ft (9.1 m)
 1546 equipment exclusion zone on each side of the ordinary high-water mark (definable bank). There
 1547 are no shade requirements and no leave tree requirements, but skidding logs in or through
 1548 streams is prohibited. Harvesting treatments included (1) clearcut and (2) thinning to a 50%
 1549 shade removal. Canopy cover measurements were made using a concave spherical densiometer.
 1550 Preharvest canopy measurements ranged from 56% to 88%, with an average of 63% in the
 1551 clearcut reaches, and 74% in the partial cut reaches. In the clearcut reaches, canopy was reduced
 1552 to 52% in 2002 and 41% in 2003, immediately following broadcast burning and replanting. In

1553 2004 and 2005, overall canopy was measured at 56% and 54%, respectively. Streamside shade
1554 recovery can be attributed entirely to low-lying understory species, as evidenced by the increase
1555 in understory/deciduous cover of 26% in 2003 to 39% and 37% in 2004 and 2005, respectively.
1556 In the partial cut reaches, canopy shade remained near 75%.

1557 In fish-bearing streams of Montana, Sugden et al. (2019) assessed the effectiveness of state
1558 riparian management harvest prescriptions in maintaining canopy cover. Montana state law
1559 requires timber be retained within a minimum of 15.2 m of fish-bearing streams, with equipment
1560 exclusion zones extended on steep slopes for up to 30.5 m. Within the riparian management
1561 zone, no more than half the trees greater than 204 mm (8 in) diameter at breast height (DBH) can
1562 be removed. In no case, however, can stocking levels of leave trees be reduced to less than 217
1563 trees per hectare. Shade over the stream surface was not directly measured in this study. Rather,
1564 canopy cover was used as a general proxy, with two independent estimates of canopy cover
1565 employed. One method used the riparian cruise data to populate a canopy cover model within the
1566 Forest Vegetation Simulator (FVS), which estimated canopy cover for each study site, pre- and
1567 post-harvest. The second method measured canopy cover in the harvest reach every 30 m, both
1568 before and after timber harvest, using a concave spherical forest densiometer. Mean canopy
1569 cover in the SMZ, as modelled in FVS, decreased from 77% to 74% following timber harvest
1570 and 73% when subtracting windthrow to differentiate between direct and indirect impacts of
1571 management (Table 3). The mean canopy cover over the stream channel based on densiometer
1572 measurements was 66% pre-harvest and 67% post-harvest. Neither of these changes was
1573 statistically significant.

1574 Roon et al. (2021a) compared the effects of two experimental thinning treatments on shade in
1575 second growth redwood stands (40-60 years old) of northern California. This study took place
1576 between 2016 and 2018 with thinning treatments applied during 2017 giving 1-year pre-
1577 treatment and 1-year of post-treatment data. Two study sites prescribed treatment on one side of
1578 the stream of a 45 m buffer width with a 22.5 m inner zone with 85% canopy retention and a
1579 22.5 m outer zone that retained 70% canopy cover (Tectah watershed). At the third treatment site,
1580 thinning prescriptions included removal of up to 40% of the basal area within the riparian zone
1581 on slopes less than 20% on both sides of the channel along a ~100–150 m reach (Lost Man
1582 watershed, Redwood national park). Shade over streams was measured with hemispherical
1583 photos and effective shade was calculated in HemiView Canopy Analysis Software version 2.1.
1584 Results for the Tectah watershed showed a significant reduction in canopy closure by a mean of
1585 18.7%, (95% CI: -21.0, -16.3) and a significant reduction of effective shade by a mean of 23.0%
1586 (-25.8, -20.1) one-year post treatment. In the Lost man watershed, a non-significant reduction of
1587 mean shade by 4.1% (-8.0, -0.5), and mean canopy closure by 1.9% was observed in 2018.
1588 Results for below canopy light availability showed significant increases by a mean of 33% (27.3,
1589 38.5) in the Tectah watershed, and non-significant increases in Lost man watershed of 2.5% (-
1590 1.6, 5.6) by 2018.

1591 In general, the results from the studies reviewed above suggest changes in shade or canopy cover
1592 from pre- to post-harvest are directly impacted by the intensity of the treatment prescription.
1593 Buffer treatments vary between states and within states by stream type (e.g., fish-bearing or non-

1594 fish-bearing), For the studies that quantified pre- to post-changes in shade along fish-bearing
 1595 streams (Cupp & Lofgren, 2014; Sugden et al. 2019), results show evidence that the application
 1596 of best management practices (BMPs) cause minimal or non-significant changes in shade
 1597 following harvest. For non-fish-bearing streams harvest prescriptions are much more variable.
 1598 Further, there are many more examples of application and comparison of different experimental
 1599 buffer treatments which vary by width or thinning targets.

1600 *Litter*

1601

1602 Table 7. Treatment and responses for selected publications investigating Litter relevant to Q1.

Reference	Treatment	Response
McIntyre et al. (2018)	Buffer width of ~50 feet (100%); 50 feet with clearcut gaps (FP); Clearcut to stream (unbuffered)	<u>2-years post-harvest</u> Total litterfall input showed a significant decrease in the FP buffers (n = 2; $\Delta = -0.2711$ g) and the unbuffered (n = 2; $\Delta = -0.3823$ g) treatments. Total Leaf litterfall (deciduous and conifer leaves combined) also showed a significant decrease in the FP buffers (n = 2; $\Delta = -0.1255$ g) and the unbuffered (n = 2; $\Delta = -0.2779$ g). Conifer litterfall input significantly decreased in the FP (n = 2; $\Delta = -0.0437$) and unbuffered (n = 2; $\Delta = -0.1574$ g) treatments. Deciduous litterfall decreased significantly only in the unbuffered (n = 2; $\Delta = -0.1563$ g) treatment. Wood input (twigs and cones) decreased significantly in the FP (n = 2; $\Delta = -0.2665$ g) and unbuffered (n = 2; $\Delta = -0.2203$ g) treatments.
Kiffney & Richardson (2010)	Upland clearcuts with (1) no buffer, (2) 10 m buffer (~30 feet), (3) 30 m buffers	<u>8 years post-harvest</u> The no buffer treatment showed an ~91% reduction of litterfall in the first year with recovery to an ~11% reduction by year 8. The 10 m buffer treatment showed an initial reduction in litterfall by ~2%, but an increase of ~37% by year 8, compared to pre-harvest. The 30 m buffer treatment showed an initial increase in litterfall by ~11% in the first year which increased to ~74% by year 8 relative to pre-harvest levels.

1603

1604 Specific to western Washington, McIntyre et al. (2018) compared the change in litterfall inputs
 1605 from pre- to post-harvest under three different riparian harvest treatments. Treatments included a
 1606 [minimum](#) two-sided 50-ft riparian buffer along at least 50% of the stream (FP; with clearcut to
 1607 stream's edge outside of the buffer), a [minimum](#) two-sided 50-ft buffer along the entire stream
 1608 (100%), and a clearcut to stream without a buffer (0%). Litterfall was collected with litter traps
 1609 placed along the mainstem channel of each site. Litter was dried and sorted by type (e.g.,
 1610 deciduous, conifer, small wood) and ashed to compare weight. Results for litterfall input showed
 1611 a decrease in total litterfall input in the FP (P = 0.0034) and 0% (P = 0.0001) treatments between
 1612 pre- and post-treatment periods. Leaf litterfall (deciduous and conifer leaves combined) input
 1613 decreased in the FP (P = 0.0114) and 0% (P < 0.0001) treatments in the post-treatment period. In
 1614 addition, conifer (conifer needles and scales) litterfall input decreased in the FP (P = 0.0437) and
 1615 0% (P < 0.0001) treatments, deciduous leaves in the 0% (P < 0.0001) treatment, wood (twigs and

1616 cones) in the FP (P = 0.0044) and 0% (P = 0.0153) treatments, and misc. (e.g., moss and flowers)
 1617 in the 0% (P = 0.0422) treatment.

1618 In the Malcom Knapp Research Forests of British Columbia, Canada, Kiffney & Richardson
 1619 (2010) compared changes in litter input between riparian harvest prescriptions that included
 1620 clear-cut to stream edge, 10 m wide buffer reserve, and 30 m buffer reserves over the course of 8
 1621 years. No thinning was applied within the reserves. Upland treatment at all sites used clearcutting
 1622 methods. Vertical litter inputs were collected monthly and at approximately 6–8-week intervals
 1623 during each season for years 1,2,6,7, and 8 years after harvest. Litter was separated into
 1624 broadleaf deciduous, twig, needles, and other (seeds, cones, and moss) categories following
 1625 collection and subsequently dried and weighed using a microbalance. Results for post-harvest
 1626 changes in litterfall input by treatment per year are summarized in Table 7. Actual values of pre-
 1627 post-harvest changes in litterfall input by type, treatment, and year were not directly reported,
 1628 however, the authors report that post-harvest inputs of needles, twigs, and total particulate matter
 1629 were significantly lower for clearcuts compared to all other treatments.

1630 Table 7. Percent change in total litterfall percentage post-harvest by treatment per year from
 1631 Kiffney & Richardson (2010). Table reproduced and modified from Yeung et al. (2019)
 1632 supplementary materials Appendix C, Table C3.

Harvest type (% of watershed area harvested)	Change in litterfall (%)	Time after harvest (year)
Clearcut (33%) no buffer	~ -91	1
	~ -78	2
	~ -79	6
	~ -47	7
	~ -11	8
Clearcut (23%); with 10-m riparian buffers	~ -2	1
	~ 6	2
	~ -14	6
	~ 6	7
	~ 37	8
Clearcut (18%); with 30-m riparian buffers	~ 11	1
	~ 44	2
	~ 14	6
	~ -6	7

Formatted: Font: Not Highlight

1633

1634 *Large Wood (LW) recruitment*

1635 Table 8. Treatment and responses for selected publications investigating Large Wood relevant to
 1636 Q1.

Reference	Treatment	Response
McIntyre et al. (2021)	Buffer width of -50 feet for full stream length (100%); 50 feet with clearcut gaps for 50% of the stream length (FP); Clearcut to stream (unbuffered)	<u>8 years post-harvest</u> Large wood recruitment rates were greatest during the first two years post-harvest, then decreased. Mean LW density increased by 66, 44, and 47% in the 100% (n = 4), FP (n=3), and unbuffered treatments (n = 4), respectively, in the first 2 years. LW density continued to increase in the FP treatment by 42 and 41%, respectively, in years 5 and 8 post-harvest.
Ehinger et al. (2021)	50 feet for 50% of the stream length (FP) 1) 50 feet, 2) <50ft buffers (variable), and 3) unbuffered, harvested to the edge of the channel	<u>3 years post-harvest</u> Treatments were analyzed by buffer type: mean LW piece counts increased in the 50 feet-foot buffer (n = 8) and the unbuffered (n = 7) treatments-type by 8 and 13%, respectively, and decreased in the <50 feet-foot (n = 6) treatments-buffer type by 15%.

1637

1638 Specific to western Washington, McIntyre et al. (2021) compared the change in mean in-stream
 1639 large wood from pre- to post-harvest under three different riparian harvest treatments in non-fish-
 1640 bearing streams. Treatments included a minimum two-sided 50-ft riparian buffer along at least
 1641 50% of the stream (FP; with clearcut to stream’s edge outside of the buffer), a two-sided
 1642 minimum 50-ft buffer along the entire stream (100%), and a clearcut to stream without a buffer
 1643 (0%). Results showed a 66% (P <0.001), 44% (P = 0.05) and 47% (P = 0.01) increase in mean
 1644 large wood density in the 100%, FP and 0% treatments, respectively, in the first 2 years post-
 1645 harvest compared with the pre-harvest period and after controlling for temporal changes in the
 1646 references. Five years post-treatment the mean LW density in the FP continued to increase 42%
 1647 (P = 0.08), and again 8 years post-treatment (41%; P = 0.09).

1648 Ehinger et al. (2021) also quantified changes in in-stream LW following ~~similar~~ the FP riparian
 1649 harvest prescription. Because of unstable slopes, total buffer area was 18 to 163% greater than
 1650 the prescribed 50-foot-buffer. This resulted in 32 different buffer types 1) buffers encompassing
 1651 the full width (50 feet), 2) <50ft buffers, and 3) unbuffered, harvested to the edge of the channel.
 1652 Because of the separation into multiple treatments, sample sizes became small and unbalanced.
 1653 Thus, no statistical analyses were conducted, and only descriptive statistics were applied for
 1654 changes in stand structure and wood loading. However, given the lack of studies presenting
 1655 changes in LW recruitment from pre- to post-harvest, it is presented here for comparison. Results
 1656 showed the full buffer sites and <50 ft buffer sites received an average of 23 and 10 pieces/100 m
 1657 and 2.3 and 0.7 m³/100 m of large wood, respectively, post-harvest. The majority of recruited
 1658 large wood pieces had stems with roots attached (SWRW); 70, and 100% in the full buffer, and
 1659 <50 ft buffer types, respectively. Pre-harvest channel large wood loading ranged from 55.8 to 111
 1660 pieces/100 m and from 9.8 to 25.2 m³/100 m among buffer types. Piece counts increased in the
 1661 full buffer and unbuffered sites (8 and 13%, respectively), and decreased in the <50 ft buffers
 1662 (15%).

1663 *Sediment*

1664 Table 9. Treatment and responses for selected publications investigating Sediment relevant to
 1665 Q1.

Reference	Treatment	Response
Hatten et al. (2018)	Buffer width of 15 m (~ 50 feet), Oregon Forest practices	<u>1 year post-harvest after 2 harvest events (n = 3)</u> Mean suspended sediment concentrations (SSC) was 32 mg L ⁻¹ (~63%) lower after the first harvest and 28.3 mg L ⁻¹ (~55%) lower after the second harvest when compared to the pre-harvest concentrations.
Bywater-Reyes et al. (2017)	Unbuffered clearcuts; 50 ft buffers. Oregon Forest Practices	<u>3 years post-harvest</u> The first year following harvest suspended sediment yield (SSY) increased in the unbuffered (n = 2) and buffered (n = 1) catchments. By year 2, SSY returned to pre-harvest levels in the buffered, and one of the unbuffered catchments. In one unbuffered catchment, SSY continued to increase annual for all three years.
Karwan et al. (2007)	Buffer width of 75 foot (Idaho Forest Practices Act) with (1) upland clearcut and (2) 50% canopy removal	<u>4 years post-harvest</u> Total suspended sediment (TSS) load from the clearcut exceeded the predicted load by 152% (6,791 kg km ⁻²) in the first year following harvest. The 50% canopy removal showed a non-significant increase in TSS. Neither treatment showed a statistical difference in TSS during the recovery time 2-4 years after harvest compared to pre-harvest.

1666

1667 No studies from Washington published since 2000 provide changes in sediment concentration or
 1668 transport from pre- to post-harvest. The Hard Rock study (McIntyre et al., 2021) reported their

1669 results for water turbidity and suspended sediment export (SSE) were stochastic in nature and the
1670 relationships between SSE export and treatment effects were not strong enough to confidently
1671 draw conclusions. The lack of SSE in some high discharge events suggests that the basins are
1672 likely to be supply limited. The Soft Rock study (Ehinger et al., 2021) similarly reported that
1673 their results for changes in sediment post-harvest were highly variable. The SSE data in the Soft
1674 Rock study indicated that the marine sedimentary lithologies were more erodible than then
1675 lithologies sampled in the Hard Rock Study. However, prediction equations could not be
1676 calculated to predict the response of the treatment sites after harvest. Thus, strong conclusions
1677 about the effectiveness of the Forest Practices harvest prescription rules on discharge and SSE
1678 could not be drawn.

1679 Hatten et al. (2018) compared pre- to post-harvest suspended sediment concentrations (SSC) in a
1680 western Oregon Alsea watershed. Treatments followed contemporary harvesting practices (no
1681 buffer in non-fish-bearing streams with equipment exclusion zones, and a 15 m no-cut-buffer in
1682 fish-bearing streams) resulted in non-significant changes in SSC at all treatment sites.
1683 Surprisingly, in the fish-bearing streams there was a decrease in SSC (~63% and ~55%, after first
1684 and second harvest, respectively) compared to pre-harvest values. Bywater-Reyes et al. (2017)
1685 compared pre- to post-harvest changes in suspended sediment yield (SSY) following harvest in
1686 the Trask River Watershed of western Oregon. Harvest treatments of study sub-watersheds
1687 consisted of clearcuts (UM2 and GC3) and a clearcut with buffers (50 ft; ~15 m; PH4).
1688 Following timber harvest, (water year 2013), increases in SSY occurred in all harvested
1689 catchments. The SSY in both PH4 (clearcut with buffers) and GC3 (clearcut without buffers)
1690 declined to pre-harvest levels by water year 2014. Interestingly, the SSY in UM2 (clearcut
1691 without buffers) increased annually throughout the post-harvest period, ultimately resulting in
1692 the highest SSY of all catchments during the final two years (2015-2016) of the study after
1693 producing the lowest SSY in the pre-harvest period. Actual values for SSY and significance were
1694 not reported.

1695 Karwan et al. (2007) compared changes in total suspended solids (TSS) in streams from pre- to
1696 post-harvest in northern Idaho. Treatments in the paired-watershed experiment consisted of 1)
1697 commercial clearcut of the watershed area of 50%, and was broadcast burned and replanted by
1698 the end of May 2003, and 2) partial cut in which a target of 50% the canopy was removed in 50%
1699 of the watershed in 2001, with final 10% of log processing and hauling in early summer of 2002.
1700 All harvests were carried out according to best management practices and in accordance with the
1701 Idaho Forest Practices Act. Results showed a significant and immediate impact of harvest on
1702 monthly sediment loads in the clear-cut watershed ($p = 0.00011$), and a marginally significant
1703 impact of harvest on monthly sediment loads in the partial cut ($p = 0.081$). Total sediment load
1704 from the clearcut over the immediate harvest interval exceeded predicted load by 152% (6,791
1705 kg km⁻²); however, individual monthly loads varied around this amount. The largest increases in
1706 percentage and magnitude occurred during snowmelt months, namely April 2002 (560%, 2,958
1707 kg km⁻²) and May 2002 (171%, 3,394 kg km⁻²). Neither treatment showed a statistical
1708 difference in TSS during the recovery time 2-4 years after harvest (clearcut: $p = 0.2336$; partial
1709 cut: $p = 0.1739$) compared to the calibration loads (pre-harvest).

1710 *Nutrients*

1711 Table 9. Treatment and responses for selected publications investigating Sediment relevant to
 1712 Q1.

Reference	Treatment	Response
McIntyre et al. (2021)	Buffer width of 50 feet (100%); 50 feet with clearcut gaps (FP); Clearcut to stream (unbuffered)	Increases in total-N export of 5.73, 10.85, and 15.94 kg/ha/yr post-harvest in the 100%, FP, and 0% treatments, respectively, were detected in the first 2 years post-harvest ; and of 6.20, 5.34, and 8.49 kg/ha/yr in the extended period (7-8 years post-harvest). Results for nitrate-N export showed changes similar to but slightly less than those seen in the total-N analysis with a relative increase in nitrate-N export of 4.79, 9.63, and 14.41 kg/ha/yr post-harvest in the 100%, FP, and 0% treatments, respectively in the first 2 years. None of the changes in the extended period were significant.
Gravelle et al. (2009)	75 foot buffers (Idaho Forest Practices Act) with (1) upland clearcut and (2) 50% canopy removal	<u>4 years post-harvest</u> Significant increases in nitrogen were observed in the clearcut (n = 1) and partial cut treatments (n = 1). Increases at the clearcut site was greatest from 0.06 mg-N L ⁻¹ (pre-harvest) to 0.35 mg-N L ⁻¹ (post-harvest period, 4 years). There was also an observable seasonal effect on NO ₃ + NO ₂ concentrations with the peak concentration of 0.89 mg-N L ⁻¹ , with mean monthly concentrations of 0.43 mg-N L ⁻¹ and 0.59 mg-N L ⁻¹ in water years 3 and 4 following harvest, respectively, in the clearcut. No significant changes of in-stream concentration of any other nutrient recorded were found between time periods and treatments.
Deval et al. (2021)	Clearcut to stream with 24-47% vegetation removal (Phase I); Clearcut with 36 – 50% vegetation removal (Phase II)	<u>6 years post-harvest (Phase I), 8 years post-harvest (Phase II); (n = 7)</u> Mean annual NO ₃ + NO ₂ concentrations increased significantly at all treatment sites during both treatment Phases with the greatest increases occurring during the Phase II period (increases ranging from 1.73 kg ha ⁻¹ yr ⁻¹ – 3.95 kg ha ⁻¹ yr ⁻¹). NO ₃ + NO ₂ concentrations followed an increasing trend throughout the post-harvest period with evidence of recovery in year 8 indicated by the flattening of the cumulative load curve.

1713
 1714 The “Hard Rock” study (McIntyre et al., 2021) results showed an increase in total-N export of
 1715 5.73 (P = 0.121), 10.85 (P = 0.006), and 15.94 (P = 0.000) kg/ha/yr post-harvest in the 100%, FP,
 1716 and 0% treatments, respectively, in the first 2 years; and of 6.20 (P = 0.095), 5.34 (P = 0.147),
 1717 and 8.49 (P = 0.026) kg/ha/yr in the extended period (7-8 years post-harvest). Results for nitrate-
 1718 N export showed changes similar to but slightly less than those seen in the total-N analysis with
 1719 a relative increase in nitrate-N export of 4.79 (P = 0.123), 9.63 (P = 0.004), and 14.41 (P <0.001)
 1720 kg/ha/yr post-harvest in the 100%, FP, and 0% treatments, respectively in the first 2 years. None
 1721 of the changes in the extended period were significant. However, the authors note that there was
 1722 high variability in the data for the extended period and nitrate-N export only returned to pre-
 1723 harvest levels in one watershed. Total phosphorus export increased post-harvest by a similar
 1724 magnitude in all treatments: 0.10 (P = 0.006), 0.13 (P = 0.001), and 0.09 (P = 0.010) kg/ha/yr in

1725 the 100%, FP, and 0% treatments, respectively in the first 2 years post-harvest. Changes in
1726 phosphorus were not reported in the extended period.

1727 Gravelle et al. (2009) compared pre- to post changes in NO^3 and NO^2 concentrations in
1728 headwater streams following a clearcut and a partial cut (50% removal of canopy cover) in
1729 northern Idaho. Riparian buffers and leave trees are not required for [non-fish-bearing](#)
1730 [non-fish-bearing](#) headwater streams in Idaho. Results showed statistically significant increases in NO^3 and
1731 NO^2 concentrations following clearcut and partial harvest cuts in headwater streams ($p < 0.001$).
1732 Increases at the clearcut treatment site were greatest, where mean monthly concentrations
1733 increased from 0.06 mg-N L⁻¹ during the calibration period to 0.35 mg-N L⁻¹ in the post-
1734 harvest period. Mean monthly concentrations in the partial cut increased from 0.04 mg-N L⁻¹ in the
1735 pre-harvest period to 0.05 mg-N L⁻¹ in the post-harvest period. No significant changes of
1736 in-stream concentration of any other nutrient recorded (total Kjeldahl nitrogen (TKN), TP, total
1737 ammonia nitrogen (TAN) consisting of unionized (NH_3) and ionized (NH_4^+) ammonia, and
1738 unfiltered orthophosphate (OP)) were found between time periods and treatments.

1739 Deval et al. (2021) compared changes in the same nutrient concentrations in the same area of
1740 northern Idaho but with an additional harvest prescription several years later. For this analysis,
1741 time periods were broken into four distinct phases: 1) pre-disturbance (1992–1997), 2) post-road
1742 (1997–2001), 3) experimental-harvest Phase I (PH-I) (2001–2007), and 4) operational sequential
1743 harvest Phase II (PH-II) when the extent and frequency of harvests increased (2007–2016). PH-I
1744 represents an experimental treatment phase during which harvest activities were experimentally
1745 controlled (only upstream headwater watersheds were harvested and mature vegetation (size or
1746 age threshold for “mature” not reported) removal ranged between 24% and 47%) followed by
1747 site management operations including broadcast burning and replanting. PH-II represents the
1748 post-experimental phase where the study area transitioned to operational treatments that
1749 consisted of additional road construction and timber harvest, with site management operations
1750 including pile burning and competition release herbicide application. During this operational
1751 phase, the mature vegetation (size or age threshold for “mature” not reported) removal in the
1752 upstream watersheds ranged between 36% and 50%. The response in $\text{NO}^3 + \text{NO}^2$ concentrations
1753 was negligible at all treatment sites following the road construction activities. However, $\text{NO}^3 +$
1754 NO^2 concentrations during the PH-I period increased significantly ($p < 0.001$) at all treatment
1755 sites. Similar to the PH-I period, all watersheds experienced significant increases in $\text{NO}^3 + \text{NO}^2$
1756 concentration during the PH-II treatment period ($p < 0.001$). Similar to Gravelle et al. (2009),
1757 significant increases in all other nutrients recorded were not detected.

1758

1759 [Focal Question 1a](#)

1760 *1a. What are the effects of thinning (intensity, extent) on the riparian functions, over the short*
1761 *and long-term compared to untreated stands?*

1762 Based on the wording of this question, papers deemed appropriate are those that compare
1763 changes in measurable data indicative of the riparian functions between harvested and
1764 unharvested stands. Further, studies chosen for this question should compare the response of

1765 these functions based on different thinning intensities. Thus, the design of the studies reviewed
 1766 for this review should be a BACI or ACI design with results reported for differences between
 1767 treatment and reference reaches. Also included are a few simulation modeling experiments that
 1768 follow these designs.

1769

Function	Count
Shade	2
Litter	0
LW	2
Sediment	1
Nutrients	1
Bank Stability	0

1770

1771 *Shade*

1772 Table 10. Treatment and responses for selected publications investigating Shade relevant to Q1a.

Reference	Treatment	Response
Anderson et. al. (2007)	69 m buffers (B1); variable width buffer averaging 22 m (VB); streamside retention buffer averaging 9 m (SR-T)	<u>2-5 years post-harvest</u> Adjacent upland to each buffer treatment was thinned to a range of 98 – 297 trees per hectare. Visible sky at stream center increased with decreasing buffer width. Untreated stands maintained ~4.2% visible sky at stream center. VB and B1 sites showed an increase of visible sky to ~9.3% and the SR-T sites showed an increase to ~9.6%.
Roon et. al. (2021a)	45 m buffer width with 70-85% canopy retention (CC); Up to 40% basal area removal along stream (BA)	<u>1-year post-harvest</u> The CC sites showed a mean canopy cover reduction of 18.7%, (-21.0, -16.3) and a significant reduction of effective shade by a mean of 23.0% (-25.8, -20.1). The BA sites showed a reduction of mean shade by 4.1% (-8.0, -0.5), and mean canopy closure by 1.9%.

1773

1774 Anderson et. al. (2007) compared changes in canopy cover at stream centers between sites
 1775 adjacent to different riparian zone treatments and an untreated control. This study was conducted
 1776 in young headwater forests of western Oregon. Treatments included three buffer widths: 1) one
 1777 site-potential tree averaging 69 m (B1), 2) variable width buffer averaging 22 m (VB), or 3)
 1778 streamside retention buffer averaging 9 m (SR-T). Adjacent upland to each buffer treatment was
 1779 thinned to ~198 trees per hectare. Results showed that visible sky at stream center only differed
 1780 significantly between SR-T (9.6%) and the untreated (4.2%) sites post-harvest. These results
 1781 were reported for the period 2-5 years post-harvest.

1782 Roon et. al. (2021a) used a BACI analysis to evaluate significant changes in canopy cover
 1783 relative to untreated reaches following 2 different thinning intensities in second growth redwood

1784 forests of northern California. One study site prescribed treatment on one side of the stream of a
 1785 45 m buffer width with a 22.5 m inner zone with a target 85% canopy retention and a 22.5 m
 1786 outer zone that retained 70% canopy cover (Green Diamond Resource Company, Tectah
 1787 watershed). The treatment site, thinning prescriptions included removal of up to 40% of the basal
 1788 area within the riparian zone on slopes less than 20% on both sides of the channel along a ~100–
 1789 150 m reach (Lost Man watershed, Redwood national park). Control reaches were located
 1790 upstream from treatment reaches. Data analysis was conducted separately for each experimental
 1791 watershed (i.e., 1 Lost man site, 2 Tectah sites). Results for the Tectah watershed showed a
 1792 significant reduction in canopy closure by a mean of 18.7%, (95% CI: -21.0, -16.3) and a
 1793 significant reduction of effective shade by a mean of 23.0% (-25.8, -20.1) one-year post
 1794 treatment. In the Lost Man watershed, a non-significant reduction of mean shade by 4.1% (-8.0, -
 1795 0.5), and mean canopy closure by 1.9% was observed. Results for below canopy light availability
 1796 showed significant increases by a mean of 33% (27.3, 38.5) in the Tectah watershed, and non-
 1797 significant increases in Lost Man watershed of 2.5% (-1.6, 5.6). Data for canopy closure and
 1798 effective shade were recorded for 1-year pre- and 1-year post-harvest.

1799 *LW*

1800 Table 10. Treatment and responses for selected publications investigating Large Wood relevant to
 1801 Q1a.

Reference	Treatment	Response
Benda et al. (2016)	simulation modeling of single entry thinning with and without a 10 m width no-cut buffers; and a double entry thinning occurring 25 years after first with and without 10 m no-cut buffers	<u>Simulated 100-year post harvest results</u> The model output for single entry thinning treatments predicts a 33% or 66% reduction of in-stream wood over a century relative to the unharvested reference for harvest on one side or both sides of the stream, respectively. Double entry thinning treatments without a buffer predicted further reduction in wood recruitment over a century of simulation with 42 and 84% reduction of in stream wood relative to the reference stream when one side and both sides of the channel were harvested.
Schuett Hames and Stewart (2019a)	30-ft no-cut buffer width, and thinning 30-75 ft from the stream (SR); retention of all shade providing trees in this area (AAS)	<u>5-years post-harvest</u> The SR and AAS LW input rates by volume were nearly 300% and 50% higher than the reference stream rates, respectively. Wood recruitment in the SR sites was significantly greater than in the AAS and reference sites. Conversely, differences in wood recruitment did not differ significantly between the AAS and reference sites.

1802
 1803 Benda et al. (2016) used simulation modeling to estimate the changes in in-stream LW volume
 1804 over time between sites with thinning treatments and unharvested reference sites. They used
 1805 ORGANON growth models to simulate forest growth and LW recruitment over a 100-year
 1806 period. The model simulated treatments of single entry thinning from below (thinning from
 1807 below removes the smallest trees to simulate suppression mortality) with and without a 10 m
 1808 width no-cut buffers; and a double entry thinning from below with the second thinning occurring

1809 25 years after the first with and without 10 m no-cut buffers (results with 10 m buffer presented
1810 in question 1b). Each thinning treatment was also combined with some mechanical introduction
1811 of thinned trees into the stream encompassing a range between 5 and 20 % of the thinned trees.
1812 The single-entry thin reduces stand density to 225 tph in 2015 (-67 %) and declines further to
1813 160 tph by 2110 (-77 %). The double entry thinning resulted in 123 tph after the second thinning
1814 in 2040 (-82%) and maintained that density until 2110. Both thinning treatments resulted in a
1815 substantial reduction of dead trees that could contribute to in-stream wood loads. The model
1816 output for single entry thinning treatments predicts a 33% or 66% reduction of in-stream wood
1817 over a century relative to the unharvested reference for harvest on one side or both sides of the
1818 stream, respectively. Including mechanical tipping of 5,10,15, and 20% of cut stems without a
1819 buffer in the single-entry thinning treatment changes the relative in-stream percentages of wood
1820 relative to the reference stream to -15, -6, +1, and +6%, respectively. Double entry thinning
1821 treatments without a buffer predicted further reduction in wood recruitment over a century of
1822 simulation with 42 and 84% reduction of in stream wood relative to the reference stream when
1823 one side and both sides of the channel were harvested. To offset the predicted changes of in
1824 stream wood volume following double entry harvest would require tipping of 10% of cut stems.
1825 The authors conclude that thinning without some mitigation efforts resulted in large losses of in
1826 stream wood over a century.

1827 Schuett Hames and Stewart (2019a) compared recruitment rates of LW and volume of in-stream
1828 LW between different riparian buffer thinning treatments and unharvested reference sites.
1829 Treatments evaluated included prescriptions for standard shade rule (a 30-ft no-cut buffer width,
1830 and thinning 30-75 ft from the stream), and all available shade rule (requires retention of all
1831 shade providing trees in this area) for eastern Washington. Results showed cumulative wood
1832 recruitment from tree fall over the five-year post-harvest interval was highest in the standard
1833 shade rule (SR) group, lower in the all-available-shade rule (AAS) group and lowest in the
1834 reference (REF) group. The SR and AAS rates by volume were nearly 300% and 50% higher
1835 than the REF rates, respectively. Wood recruitment in the SR sites was significantly greater than
1836 in the AAS and reference sites ($P < 0.05$). Conversely, differences in wood recruitment did not
1837 differ significantly between the AAS and reference sites.

1838 *Sediment*

1839 Karwan et al. (2007) used BACI analysis to compare changes in total suspended solid (TSS)
1840 yields between thinned sites and unharvested reference sites. This study was conducted in the
1841 Mica Creek Experimental watershed of northern Idaho and focused on ~~non-fish bearing~~
1842 ~~non-fish~~ [bearing](#) headwater streams. The thinning treatment included a target 50% canopy removal
1843 without no-cut buffers. Results showed a marginally significant ($P = 0.081$) increase in TSS
1844 relative to the reference streams in the first year following treatment. However, differences in
1845 TSS between the treatment streams and reference streams were not significant ($p = 0.174$) in the
1846 period 2-4 years post-harvest.

1847 *Nutrients*

1848 Yang et al. (2021) compared changes in stream chemistry between streams along thinned stands
1849 and unharvested reference stands in young mixed conifer headwater basins of the Sierra National

1850 Forest. Thinning treatment included mastication of shrub cover to < 10% and harvesting of trees
 1851 to a target basal area of 27–55 m² ha⁻¹. Data for dissolved organic carbon (DOC) and dissolved
 1852 organic nitrogen (DON) were recorded for 2 years prior to and 3 years after treatment. For
 1853 stream water, volume-weighted concentrations of DOC were 66- 94% higher in thinned
 1854 watersheds than in control watersheds for all three consecutive drought years following thinning
 1855 (p = 0.06, 0.01, and 0.05 for years 1,2, and 3 post-harvest, respectively). No differences in DOC
 1856 concentrations were found between thinned and control watersheds before thinning (p = 0.50,
 1857 and 0.74 for pre-harvest years 1 and 2, respectively). Volume-weighted concentrations of DIN
 1858 were 24% higher in thinned than in control watersheds only in the third year following thinning
 1859 (p = 0.04). No differences in DIN were detected between treatment and reference streams in the
 1860 2 pre-harvest years (P ≥ 0.44). Note: Drought occurred at both sites during the three post-harvest
 1861 years which may have compounded these effects. This is discussed in more detail in question 3.

1862

1863 **Focal Question 1b**

1864 *Ib. How do buffer widths and adjacent upland timber harvest prescriptions influence impacts of*
 1865 *riparian thinning treatments?*

1866 An experimental design that could provide information useful in answering this question would
 1867 involve a comparison of sites with different buffer widths, all with upland harvest, and data
 1868 would need to be recorded before and after thinning, with or without a control site (BAI, BACI),
 1869 or differences after thinning between treatment and control sites (ACI). Three papers include an
 1870 experimental design that investigate different buffer widths or different upland treatments along
 1871 with riparian thinning treatments.

1872 *Shade*

1873 Anderson et al. (2007) compared changes in canopy cover at stream centers between sites
 1874 adjacent to different riparian zone treatments and an untreated control. This study was conducted
 1875 in young headwater forests of western Oregon. Treatments included three buffer widths (1) one
 1876 site-potential tree averaging 69 m (B1), (2) variable width buffer averaging 22 m (VB), or (3)
 1877 streamside retention buffer averaging 9 m (SR-T); the adjacent upland to each buffer was thinned
 1878 to ~198 trees per hectare. Results showed that visible sky at stream center only differed
 1879 significantly between SR-T (9.6%) and the untreated (4.2%) sites post-harvest. These results
 1880 were reported for the period 2-5 years post-harvest.

1881 *LW*

1882 Table 11. Treatment and responses for selected publications investigating Large Wood relevant to
 1883 Q1b.

Reference	Treatment	Response
-----------	-----------	----------

Burton et al. (2016)	Buffer widths were 6, 15, or 70 meters and upland thinning was to 200 trees per ha (tph); unthinned reference stand of ~400 tph.	<u>5 years post-harvest</u> slightly higher volumes of wood were found in sites with a narrow 6-m buffer (not significant), as compared with the 15-m and 70-m buffer sites 5 years after harvest.
Benda et al. (2016)	simulation modeling of thinning from below with and without a 10 m width no-cut buffers;	<u>Simulated 100-year post harvest results</u> Adding a 10 m buffer reduced total reduction of in stream wood to 11 and 22% for thinning on one and both sides of the channel, respectively, from the predicted 42 and 84% reduction without the 10 m buffer.

1884

1885 Burton et al. (2016) examined the relationship between annual in-stream wood loading and
 1886 riparian buffer widths adjacent to upland thinning operations. Buffer widths were 6, 15, or 70
 1887 meters and upland thinning was to 200 trees per ha (tph), with a second thinning (~10 years later)
 1888 to ~85 tph, alongside an unthinned reference stand of ~400 tph. Their results showed that slightly
 1889 higher volumes of wood were found in sites with a narrow 6-m buffer (not significant), as
 1890 compared with the 15-m and 70-m buffer sites in the first 5 years after the first harvest and
 1891 maintained through year 1 of the second harvest (end of study). The authors attributed this
 1892 difference to a higher likelihood of logging debris and/or windthrow, but these factors were not
 1893 analyzed.

1894 Benda et al. (2016) used simulation modeling to estimate the changes in in-stream LW volume
 1895 over time between sites with thinning treatments and unharvested reference sites. They used
 1896 ORGANON growth models to simulate forest growth and LW recruitment over a 100-year
 1897 period. The model simulated treatments of single entry thinning from below (thinning from
 1898 below removes the smallest trees to simulate suppression mortality) with and without a 10 m
 1899 width no-cut buffers; and a double entry thinning from below with the second thinning occurring
 1900 25 years after the first with and without 10 m no-cut buffers. Each thinning treatment was also
 1901 combined with some mechanical introduction of thinned trees into the stream encompassing a
 1902 range between 5 and 20 % of the thinned trees. The single-entry thin reduces stand density to 225
 1903 tph in 2015 (-67 %) and declines further to 160 tph by 2110 (-77 %). The double entry thinning
 1904 resulted in 123 tph after the second thinning in 2040 (-82%) and maintained that density until
 1905 2110. Both thinning treatments resulted in a substantial reduction of dead trees that could
 1906 contribute to in-stream. The model output for single entry thinning treatments predicts a 33% or
 1907 66% reduction of in-stream wood over a century relative to the unharvested reference for harvest
 1908 on one side or both sides of the stream, respectively. Adding the 10-m no cut buffer reduced total
 1909 loss to 7 and 14%. Including mechanical tipping of 5,10,15, and 20% of cut stems without a
 1910 buffer in the single-entry, thinning treatment changed the relative in-stream percentages of wood
 1911 relative to the reference stream to -15, -6, +1, and +6%, respectively. To completely offset the
 1912 loss of in stream wood due to single entry thinning, mechanical tipping of 14 and 12% were

1913 required without and with buffers. Double entry thinning treatments without a buffer predicted
1914 further reduction in wood recruitment over a century of simulation with 42 and 84% reduction of
1915 in stream wood relative to the reference stream when one side and both sides of the channel were
1916 harvested. Adding a 10 m buffer reduced total reduction of in stream wood to 11 and 22% for
1917 thinning on one and both sides of the channel. To offset the predicted changes of in stream wood
1918 volume following double entry harvest would require tipping of 10 and 7% of cut stems without
1919 and with the 10-m buffer. The authors conclude that thinning without some mitigation efforts
1920 resulted in large losses of in stream wood over a century.

1921

1922 Focal Question 1c

1923 *1c. What are the effects of clearcut gaps in riparian stands (intensity, extent) on the riparian*
1924 *functions, over the short and long-term, compared to untreated stands?*

1925 This question uses the general term “clearcut gaps” as a treatment within the riparian area but
1926 does not define a minimum or maximum threshold for gap size. Thus, studies reviewed that used
1927 a “patch” treatment were included as having information useful in answering this question. The
1928 question also identifies a comparison with untreated stands. Therefore, any design with a control
1929 site (BACI, ACI) is appropriate.

1930 There appears to be a paucity of studies in the literature that investigate the effects of gaps or
1931 patch harvesting treatments on riparian function within riparian stands. Only 4 papers discussed
1932 the effects of prescribed gaps or patches in the riparian area on riparian function.

1933 The “Hard Rock” study from McIntyre et al. (2021) and the “Soft Rock” study from Ehinger et
1934 al. (2021) present the most relevant results useful for answering this question. Riparian buffer
1935 prescriptions for ~~non-fish-bearing~~ streams in western Washington use a gap
1936 design. In this design, a 50-foot buffer is required along at least 50% of the treated stream length.
1937 The remaining 50% or less of the treated riparian management zone can be clear cut to the
1938 stream edge. The Hard Rock study compared differences in shade, in-stream sediment and
1939 nutrient concentrations, and large wood recruitment between treated and unharvested ~~reaches~~
1940 ~~basins~~ for 8-9 years post-harvest. ~~The first iteration of the Hard Rock study (McIntyre et al.~~
1941 ~~2021) also compared differences in litter inputs following treatment for 2 years post-harvest~~
1942 ~~between treatment and reference reaches.~~

1943 The Soft Rock study compared differences in the same functions between treated and
1944 unharvested ~~reaches~~basins, with 3-6 years of post-harvest sampling depending on the function
1945 under investigation. ~~The first phase of the Hard Rock study (McIntyre et al. 2018) also compared~~
1946 ~~differences in litter inputs following treatment for 2 years post-harvest between treatment and~~
1947 ~~reference reaches. However, because of unstable slopes in some of the sites in the Soft Rock~~
1948 ~~study, many of the buffers were required to be wider than 50 feet (ranging from 18–160% wider~~
1949 ~~than 50 feet). Conversely, some of the sites treated ended up with buffers narrower than 50 feet.~~
1950 ~~Further, there was limited availability of sites that fit the criteria (marine sediment lithology,~~
1951 ~~timing of treatment). Because of these limitations, statistical analysis, and comparison of~~
1952 ~~response between treatments and references for stream temperature and shade could not be~~

1953 performed. However, descriptive statistics were provided that contain useful information. Results
 1954 from formal statistical analyses are provided for all other functions.

1955 *Shade*

1956 Table 12. Treatment and responses for selected publications investigating Shade relevant to Q1c.

1957

Reference	Treatment	Response
McIntyre et al. (2021)	Buffer width of 50 feet (100%); 50 feet with clearcut gaps (FP); Clearcut to stream (unbuffered)	9-years post-harvest Effective shade results showed decreases of 11, 36, and 74 percent in the 100%, FP, and 0% treatments, respectively. Results for canopy cover showed that riparian cover declined after harvest in all buffer treatments reaching a minimum around 4 years post-harvest (after mortality stabilized).
Janisch et al. (2012)	Patched buffer: clearcut to stream with ~50-110 m patches retained; continuous buffer 10-15 m	<u>1-year post-harvest</u> After treatment, canopy cover in the clearcut catchments averaged 53%, canopy cover in the patch buffer treatment averaged 76%, and canopy cover in the continuous buffer treatment averaged 86%. The canopy cover of the clearcut and patch buffer treatments were significantly lower than in the reference streams.
Swartz et al. (2020)	20 m diameter clearcut gaps over stream at 30 m intervals.	<u>1-year post-harvest (n = 6)</u> Post-harvest significant increase in mean reach light to a mean of 3.91 (SD ± 1.63) moles of photons m ⁻² day ⁻¹ , overall resulting in a mean change in light of 2.93 (SD ± 1.50) moles of photons m ⁻² day ⁻¹ . The areas beneath each gap had notable localized declines in shade, though the entirety of the treatment reach (100 m) mean shading declined by only 4% (SD ± 0.02%).

1958

1959 The Hard Rock study reported significant decreases in canopy cover (measured at 1 meter above
 1960 the stream surface with a spherical densiometer) for all treated sites immediately following
 1961 harvest compared to the reference sites ($p < 0.05$). The mean canopy cover decreased from 96%
 1962 (pre-harvest) to 72% in the first-year post-harvest and continued to decline for four years
 1963 reaching a minimum of 54%. After year four, mean canopy cover began to recover increasing
 1964 annually until year 9 to 74%. In contrast, mean canopy cover in the reference sites was 95%
 1965 before harvest and never fell below 85% for 9 years. In the Soft Rock study, mean canopy
 1966 closure decreased in the treatment sites from 97% in the pre-harvest period to 75%, 68%, and
 1967 69% in the first, second, and third post-harvest years, respectively; and was further related to the
 1968 proportion of stream buffered and to post-harvest windthrow within the buffer. Canopy closure
 1969 remained stable in the reference sites throughout the course of the study, ranging from 95 to
 1970 99%.

1971 Janisch et al. (2012) compared canopy cover before and after application of a “patched buffer”
 1972 treatment with unharvested control reaches in headwater streams of western Washington. The
 1973 “patched buffer” treatment included retention of portions of the riparian forests ~50-110 m long
 1974 in distinct patches along the channel with the remaining riparian area clearcut. There was no
 1975 standard width for patched buffers, with buffers spanning the full width of the floodplain area

1976 and/or extending some undefined distance away from the stream. Canopy density was measured
1977 once in the summer prior to logging and once in the summer following logging. The percentage
1978 of visible sky was determined from digital photos taken with a fish-eye lens using Hemiview
1979 Canopy Analysis software. Canopy cover in all streams averaged 95% prior to harvest and did
1980 not differ between treatment and reference streams. Following treatment, canopy cover in the
1981 patch buffer treatment averaged 76% and differed significantly from reference reaches.

1982 Swartz et al. (2020) tested the effects of adding canopy gaps within young (40 – 60 years old),
1983 regenerating forests of western Oregon on stream light availability and stream temperatures.
1984 While light availability and stream temperature are not functions described in the FPHCP, they
1985 are directly related to shade (an FPHCP function) Also, they directly affect water quality and
1986 aquatic habitat productivity which are functional objectives within the FPHCP. Further,
1987 considering the paucity of studies available that investigate the effects of clearcut gaps, the
1988 results are presented here. The addition of gaps in the young regenerating forests were used to
1989 theoretically mimic the natural disturbance regimes and the higher canopy complexity of late-
1990 successional forests. The researchers used a BACI design on six replicated streams within the
1991 McKenzie River Basin. In each treatment reach, gaps were designed to create openings in the
1992 canopy that were approximately 20 m in diameter. Gaps were centered on a tree next to the
1993 stream and spaced approximately 30 meters apart along each reach. The BACI analysis showed
1994 strong evidence for significant increase in mean reach light ($p < 0.01$) up to 3.91 (SD \pm 1.63)
1995 moles of photons $m^{-2} day^{-1}$ and an overall mean change in light of 2.93 (SD \pm 1.50) moles of
1996 photons $m^{-2} day^{-1}$. Mean stream shading could not be evaluated in the full BACI analysis
1997 because post-treatment hemispherical photographs could not be taken at all sites due to fire
1998 impeding access. For the remaining sites, the areas beneath each gap had notable localized
1999 declines in shade, though the entirety of the treatment reach mean shading declined by only 4%
2000 (SD \pm 0.02%).

2001 *Litter*

2002 The Hard Rock study only quantified changes in litter input for 2 years after treatment (McIntyre
2003 et al., 2018). While significant decreases in litter input were observed from pre- to post-harvest
2004 in the treatment sites (described in focal question 1) these values were not significant when
2005 compared to the changes in the reference sites. Litter input was not ~~quantified~~ collected in the
2006 Soft Rock study.

2007 *LW*

2008 For the Hard Rock study, large wood recruitment and loading were only compared between the
2009 reference reaches and the buffered portion of the treatment reaches. The authors report large
2010 wood recruitment into the channel was 3 times greater on average in the treatment buffer than in
2011 the reference over the 8-year post-treatment period. However, while considerable, these
2012 differences were not significant for any analyzed post-harvest interval (e.g., 1-2 years post, 1-5
2013 years post, or 1-8 years post). The lack of significance was attributed to the large variability in
2014 recruitment values among treatment sites. The greatest increase in LW recruitment in the
2015 treatment sites relative to the reference sites occurred in the first 2 years post-harvest. Large

2016 wood loading (pieces/m of channel length) increased significantly ($\alpha = 0.10$) in the treatment
2017 reaches, relative to the reference sites in the first 2 years (47%; $p = 0.05$), 5 years (42%; $p =$
2018 0.08), and 8 years (41%; $p = 0.09$) post-harvest. For the Soft Rock study there was little post-
2019 harvest large wood input in reference sites: an average of 4.3 pieces and 0.34 m³ of combined in-
2020 and over-channel volume per 100 m of channel. In contrast, the full buffer sites and <50 ft buffer
2021 sites received an average of 23 and 10 pieces/100 m and 2.3 and 0.7 m³/100 m of large wood,
2022 respectively.

2023 *Sediment*

2024 For the Hard Rock study, results for water turbidity and suspended sediment export (SSE) were
2025 stochastic in nature and the relationships between SSE and treatment effects were not strong
2026 enough to confidently draw conclusions. Water turbidity and SSE increased with stream
2027 discharge during large storm events but rapidly declined. The Soft Rock study reported similar
2028 issues with the data for SSE in that it appeared to be driven by site and event specific factors and
2029 strong conclusions could not be drawn. The authors report that the softer lithologies sampled as
2030 part of this study were more erodible than the competent lithologies sampled in the companion
2031 Hard Rock Study.

2032 *Nutrients*

2033 The Hard Rock study analyzed changes in total nitrogen and nitrate export in the gap buffers
2034 relative to untreated reference streams. Results showed an increase in total nitrogen export in the
2035 treatment sites of 10.85 kg/ha/yr ($p = 0.006$) in the first two years post-harvest relative to the
2036 reference sites. In the extended periods, total nitrogen export increased by 5.34 ($p = 0.147$)
2037 kg/ha/yr relative to the reference streams. Results for NO³ export showed similar but slightly
2038 lower increases than total nitrogen with a relative increase in NO³ export of 9.63 ($p = 0.004$)
2039 kg/ha/yr for the first two years post-harvest relative to the reference. None of the changes in
2040 nitrate exports in the extended period were significant. The Soft Rock study reported significant
2041 increases in concentrations of total nitrogen ($p < 0.05$) and NO³ ($p < 0.05$) post-harvest in the
2042 treatment sites relative to the reference sites. The change in export appeared related to the
2043 proportion of stream buffered.

2044

2045 [Focal Question 1d](#)

2046 *1d. How do buffer widths and upland timber harvest influence impacts of clearcut gaps*
2047 *treatments?*

2048 The wording of this question implies that the effects of clearcut gaps (discussed in focal question
2049 1c) on riparian function could be impacted when paired with different buffer widths and upland
2050 harvest prescriptions. Similar to the results of the search in literature for focal question 1c, there
2051 was a paucity of riparian function studies that implemented a clearcut gap or patch cutting
2052 method within the riparian area. The added layer of complexity in this question specifying
2053 differences in buffer widths and upland harvests only further refined the selection of appropriate
2054 papers. Of the studies reviewed above, none included the evaluation of different buffer widths or

2055 different upland harvests in their experimental design. The Hard Rock study compared the
2056 clearcut gap buffers to full retention buffer and unbuffered sites (discussed in the literature
2057 review section), but different widths were not compared in the gap buffer treatments.

2058

2059 [Focal Question 1e](#)

2060 *1e. What are the effects of any combinations of the above treatments?*

2061 No studies found in our search compared the effects of combined treatments on one or more of
2062 the five functions, likely because combining multiple treatments into one design has the potential
2063 to confound results and are difficult to implement with sufficient sample sizes. The majority of
2064 the studies listed in our review investigate the effects of buffer width, thinning treatments, and
2065 upland treatments separately.

2066 The only papers with some extractable evidence of the compounding/ameliorating effects of
2067 combined treatments were focused on shade. One study, Reiter et al. (2020), compared the
2068 effects of thinned and unthinned buffers, and clearcut on changes in percent shade over adjacent
2069 streams (discussed in focal question 1). However, changes in shade were not statistically
2070 analyzed and the implementation of the upland thinning treatment only occurred at one site
2071 (Table 6).

2072

2073 [Focal Question 2](#)

2074 *2. How and to what degree do specific site conditions (e.g., topography, channel width and
2075 orientation, riparian stand age and composition) influence the response of the riparian
2076 functions?*

2077 Multiple studies have investigated the influences of site conditions on riparian function. Few
2078 studies reviewed (4) investigated the interaction between specific site conditions (e.g., slope,
2079 lithology, elevation) and harvest on the response of riparian function. However, if these specific
2080 site conditions influence the magnitude of riparian function in the absence of harvest, it is
2081 possible they can compound the effects of harvest on their response. Thus, studies that assess the
2082 relationship between site factors and riparian function may provide some useful insight for
2083 management and are presented below. Further, we also included studies that investigated the
2084 relationships between road development and sediment transport because road development is
2085 directly related to changes in local topography.

2086 *Litter*

2087 Table 13. Treatment and responses for selected publications investigating Litter relevant to Q2.

Reference	Treatment	Response
-----------	-----------	----------

Hart et al. (2013)	Remove plants in a 5 x 8 m section adjacent to stream < 10 cm DBH and >12 cm height every 2 months. 5 m fence extending underground and parallel to the stream	<u>1-2 years post-treatment (n = 5)</u> . Deciduous-site vertical litter input (504 g m ⁻¹ y ⁻¹) exceeded that from coniferous sites (394 g m ⁻¹ y ⁻¹ , 336.4–451.7) by 110 g/m ² (28.6–191.6) over the full year. Annual lateral inputs at deciduous sites (109 g m ⁻¹ y ⁻¹) were 46 g/m more than at coniferous sites (63 g m ⁻¹ y ⁻¹). Lateral inputs calculated for a 3-m-wide stream accounted for 9.6% (5.4–12.5) of total annual inputs at coniferous sites and 12.7% (10.2–14.5) of total inputs at deciduous sites. The strongest deciduous inputs to streams occurred in November. Annual lateral litter input increased with slope at deciduous sites (R ² = 0.4073) but showed no strong relationship at coniferous sites (R ² = 0.1863).
Bilby & Heffner (2016)	Simulation modeling and field sampling	<u>1-year of litterfall data</u> the majority of the litter recruited into the stream originated from within 10 m of the stream regardless of litter or stand type. No difference was found in delivery distance and litter type (needles or broadleaf) at young sites (ages not specified; canopy height mean = 32.4 m). However, needles released at mature (canopy height mean = 47 m) sites had a higher proportion of cumulative input from greater distances than needles or alder leaves released at younger sites. Litter travel distance was linearly related to wind speed (p < 0.0001). Doubling wind speed at one site led to a 67-87% expansion of the riparian contribution zone in the study area.

2088

2089 Hart et al. (2013) compared litter delivery into streams between riparian zones dominated by
 2090 deciduous (red alder) and coniferous (Douglas-fir) tree species in western Oregon. Results from
 2091 this study show that deciduous forests dominated by red alder delivered significantly greater
 2092 vertical and lateral inputs (g m⁻² y⁻¹) to adjacent streams than did coniferous forests dominated by
 2093 Douglas-fir. Deciduous-site vertical litter input (mean = 504 g m⁻² y⁻¹) exceeded that from
 2094 coniferous sites (394 g m⁻² y⁻¹) by 110 g/m² over the full year. Annual lateral inputs at
 2095 deciduous sites (109 g m⁻² y⁻¹) were 46 g m⁻² y⁻¹ more than at coniferous sites (63 g m⁻² y⁻¹).
 2096 The timing of the inputs also differed, with the greatest differences occurring in November
 2097 during autumn peak inputs for the deciduous forests. Further, annual lateral litter input increased
 2098 with slope at deciduous sites (R² = 0.4073, p = 0.0771), but showed no strong relationship at
 2099 coniferous sites (R² = 0.1863, p = 0.2855). These results were partially consistent with Bilby &
 2100 Heffner (2016) in that they suggest litter type, and topography (slope) can affect the litter input
 2101 rates.

2102 Bilby & Heffner (2016) used a combination of field experiments, literature review, and modeling
 2103 to estimate the relative importance of factors affecting litter delivery from riparian areas into
 2104 streams of western Washington in the Cascade mountains at high and low elevations. Their
 2105 results for conifer needles released at mature sites had a higher proportion of cumulative input
 2106 from greater distances than needles or leaves released at younger sites. The authors suggest from
 2107 their interpretation of the model that the width of the litter contributing area was ~35% greater at
 2108 mature sites than at young sites. The mean age of “mature” and “young” sites was not specified
 2109 but the mean tree heights were 47.0 m and 32.4 m for the mature and young sites, respectively.
 2110 Thus, tree height is related to the width of the litter contributing area for conifer needles. Litter

2111 travel distance was also linearly related to wind speed ($p < 0.0001$). Doubling wind speed at one
 2112 site led to a 67-87% expansion of the riparian litter contribution zone in the study area.
 2113 Interpretation of the regression curves revealed a trend that suggests hillslope gradient affects the
 2114 width of the litter contributing area as well. However, the authors did not apply statistical
 2115 analysis to these values and only speculated that increasing the slope from 0-45% would increase
 2116 the width of the litter contributing area by up to 70%.

2117 *LW*

2118

2119

2120 Table 14. Treatment and responses for selected publications investigating Large Wood relevant to
 2121 Q2.

Reference	Treatment	Response
Wing & Skaugset (2002)	Relationships between channel and habitat characteristics with LW piece count and volume	<u>Observation data from in 3793 stream reaches in western Oregon State.</u> LW volume: reaches with < 2.3% gradient averaged 5.8 m ³ while higher gradient streams averaged 17.9 m ³ per reach for all land types (ownership, forested and non-forested). Reaches with gradients less than 4.7% averaged LW volume of 11.5 m ³ , while mean volume at higher gradient reaches was 25.2 m ³ . LW pieces: Streams <12.2 m bank full width averaged 11.1 LW pieces per reach while larger channels averaged 4.9 pieces per reach. For key LW pieces (logs at least 0.60 m in diameter and 10 m long), stream gradient was again most important. gradient < 4.9% averaged 0.5 key LW pieces per reach while streams with higher gradients averaged 0.9 key LW pieces per reach.
Sobota et al. (2006)	patterns of riparian tree fall directions	<u>Data was collected from 21 field sites</u> Projections of LW recruitment estimated that sites with uniform steep side slopes (>40%) produced between 1.5 to 2.4 times more in stream LW by number of tree boles than sites with uniform moderate side slopes (< 40%).

2122

2123 Wing & Skaugset (2002) investigated the relationships between channel and habitat
 2124 characteristics with LW piece count and volume in stream reaches in western Oregon. This study
 2125 analyzed an extensive spatial database of aquatic habitat conditions created for western Oregon
 2126 using stream habitat classification techniques and a geographic information system (GIS).
 2127 Regression tree analysis (an exploratory regression analysis that allows for the inclusion of
 2128 multiple explanatory variables) was used to compare the relative strength of each variable in
 2129 predicting LW volume. Explanatory variables used in this analysis included morphology of
 2130 active channel (hillslope, terrace, terrace hillslope, unconstrained), and lithology (e.g., alluvium,

2131 basalt, etc.). Results for channel characteristics showed that stream gradient was the most
2132 important explanatory variable for LW volume. The split for stream gradient occurred for reaches
2133 with < 2.3% gradient (mean LW volume: 5.8 m³ per reach) while higher gradient streams showed
2134 a mean LW volume of 17.9 m³ per reach.

2135
2136 For LW pieces in forested stream reaches bankfull channel width was the most important
2137 explanatory variable with the split occurring for streams channels less than 12.2 m wide. LW
2138 pieces for streams <12.2 m wide averaged 11.1 LW pieces per reach while larger channels
2139 averaged 4.9 pieces per reach; in this model the BFW split explained 7% of the variation in LW
2140 pieces found in forested streams. For key LW pieces (logs at least 0.60 m in diameter and 10 m
2141 long) in forested reaches, stream gradient was again the most important explanatory variable
2142 with the split occurring at a slope of 4.9%. The streams with a gradient < 4.9% averaged 0.5 key
2143 LW pieces per reach while streams with higher gradients averaged 0.9 key LW pieces per reach;
2144 in this model stream gradient explained 8% of the variation in key LW pieces found in streams.

2145
2146 Lithology caused second, third or fourth level splits after stream gradient or BFW. Specifically,
2147 Mesozoic sedimentary and metamorphic geologies, located in southern Oregon stream reaches,
2148 were grouped and split from basalt, Cascade, and marine sedimentary geologies. In stream
2149 reaches with Mesozoic sedimentary and metamorphic geologies, the quantity of [LWD-large](#)
2150 [wood](#) was roughly half the amount found in other geologies. The only exception to this grouping
2151 was for LW volume in larger stream reaches, where basalt and marine sedimentary geologies
2152 contained more LW volume when grouped separately from all other geologies in a fourth-level
2153 split. The authors conclude that the geomorphic characteristic of stream reaches, in particular
2154 stream gradient and bankfull width, correlated best with LW presence.

2155
2156 Sobota et al. (2006), evaluated patterns of riparian tree fall directions in diverse environmental
2157 conditions and evaluate correlations with tree characteristics, forest structural variables, and
2158 topographic features. Specifically, the authors were interested in correlations between fall
2159 directionality and tree species type, tree size, riparian forest structure, and valley topography
2160 (side slope). Data was collected from 21 field sites located west of the Cascade Mountains crest
2161 (11 sites: Coast Range and west slopes of the Cascades), and in the interior Columbia Basin (10
2162 sites: east slopes of the Cascades, Blue Mountains, and Northern Rockies) of Oregon,
2163 Washington, Idaho, and Montana, USA. Streams were second- to fourth-order channels and had
2164 riparian forests that were approximately 40 to >200 years old. Model projections of LW
2165 recruitment estimated that sites with uniform steep side slopes (>40%) produced between 1.5 to
2166 2.4 times more in stream LW by number of tree boles than sites with uniform moderate side
2167 slopes (< 40%). The authors warn that while side slope categories (>40%, <40%) was the
2168 strongest predictor of tree fall direction in this study, they believe the differences in tree fall
2169 direction between these categories mainly characterized differences between fluvial (88% of
2170 moderate slope sites) and hillslope landforms (71% of steep slope sites). They suggest that the
2171 implications from this study are most applicable to small- to medium-size streams (second to
2172 fourth order) in mountainous regions where sustained large wood recruitment from riparian
2173 forest mortality is the significant management concern.

2174 *Sediment*

2175 Table 15. Treatment and responses for selected publications investigating Sediment relevant to
 2176 Q2.

Reference	Treatment	Response
Bywater-Reyes et al. (2017)	basin lithology and physiography effects on sediment delivery	<u>6 years of data from the Trask River Watershed</u> Site lithology was the first order control over suspended sediment yield (SSY). SSY was greater in catchments underlain by Siletz Volcanics ($r = 0.6$), the Trask River Formation ($r = 0.4$), and landslide deposits ($r = 0.9$). There was a strong negative correlation of SSY with percent area underlain by diabase ($r = -0.7$), with the lowest SSY associated with 100% diabase
Bywater-Reyes et al. (2018)	catchment lithography, physiography, discharge, and disturbance history effects on sediment delivery	<u>60 years of data in the H.J. Andrews experimental watershed (n = 10)</u> Watershed slope variability combined with cumulative annual discharge explained 67% of the variation in annual sediment yield. When considering disturbance, the largest magnitude changes in sediment movement, were after floods with a ≥ 30-year return interval .
Mueller & Pitlick (2013)	correlation analysis to assess the relative impact of lithology, basin relief, mean basin slope, and drainage density on in stream sediment supply.	<u>Data sets ranging 1-96 years for 83 basins</u> the strongest correlation of bankfull sediment concentration was with basin lithology , and showed little correlation strength with slope, relief and drainage density. As lithologies become dominated by softer parent materials (volcanic and sedimentary rocks), bankfull sediment concentrations increased by as much as 100-fold.
Litschert & MacDonald (2009)	Post-harvest stream sediment delivery pathway development frequency and characteristics.	<u>1-year post-harvest data (n = 200 harvest units)</u> 19 harvest units developed sediment delivery pathways. Pathway length and probability of connecting to stream was significantly correlated with mean annual precipitation, cosine of the aspect, elevation, and hillslope gradient .

2177
 2178

2179 Bywater-Reyes et al. (2017) assessed the influence of natural controls (basin lithology and
 2180 physiography) and forest management on suspended sediment yields in temperate headwater
 2181 catchments. This study analyzed 6 years of data from the Trask River Watershed in northeastern
 2182 Oregon and included data from harvested and unharvested sub-catchments underlain by
 2183 heterogenous lithologies. Results from this study indicate that site lithology was the first order
 2184 control over suspended sediment yield (SSY) with SSY varying by an order of magnitude across
 2185 lithologies observed. Specifically, SSY was greater in catchments underlain by Siletz Volcanics
 2186 ($r = 0.6$), the Trask River Formation ($r = 0.4$), and landslide deposits ($r = 0.9$) and displayed an
 2187 exponential relationship when plotted against the percentage of watershed area underlain by
 2188 these lithologies. In contrast, site lithology had a strong negative correlation with percent area
 2189 underlain by diabase ($r = 0.7$), with the lowest SSY associated with 100% diabase. Following
 2190 timber harvest, increases in SSY occurred in all harvested catchments but returned to pre-harvest
 2191 levels within 1 year except for sites that were underlain by sedimentary formations and were

2192 clearcut without protective buffers. The authors conclude that sites underlain with a friable
2193 lithology (e.g., sedimentary formations) had on average, SSYs an order of magnitude higher
2194 following harvest than those on more resistant lithologies (intrusive rocks).

2195 Bywater-Reyes et al. (2018) quantified how sediment yields vary with catchment lithography and
2196 physiography, discharge, and disturbance history (management or natural disturbances) over 60
2197 years in the H.J. Andrews experimental watershed in the western Cascade Range of Oregon. A
2198 linear mixed effects model (log transformed to meet the normality assumption) was used to
2199 predict annual sediment yield. In this model, site was treated as a random effect while discharge
2200 and physiographic variables were treated as fixed variables. This allowed for the evaluation of
2201 the relationships between sediment yield and physiographic features (slope, elevation, roughness,
2202 and index of sediment connectivity) while accounting for site. To account for the effect of
2203 disturbance history a variable was added to the model when the watershed had a history of
2204 management or natural disturbances. If the models for the disturbed watersheds significantly
2205 underpredicted the sediment discharge, the timing of the sudden increases were further examined
2206 to assess whether it correlated with a disturbance event. The results showed that watershed
2207 physiography combined with cumulative annual discharge explained 67% of the variation in
2208 annual sediment yield across the 60-year data set. Relative to other physiographic variables,
2209 watershed slope was the greatest predictor of annual suspended sediment yield. However, the
2210 results showed that annual sediment yields also moderately correlated with many other
2211 physiographic variables and caution that the strong relationship with watershed slope is likely a
2212 proxy for many processes, encompassing multiple catchment characteristics.

2213 Mueller & Pitlick (2013) used correlation analysis to assess the relative impact of lithology,
2214 basin relief, mean basin slope, and drainage density on in stream sediment supply defined by the
2215 bankfull sediment concentration (bedload and suspended load). The study used sediment
2216 concentration data from 83 drainage basins in Idaho and Wyoming. Lithologies of the study area
2217 were divided into four categories ranging from hardest to softest- granitic, metasedimentary,
2218 volcanic, and sedimentary. The results showed the strongest correlation of bankfull sediment
2219 concentration was with basin lithology, and showed little correlation strength with slope, relief
2220 and drainage density. As lithologies become dominated by softer parent materials (volcanic and
2221 sedimentary rocks), bankfull sediment concentrations increased by as much as 100-fold. The
2222 authors interpret these results as evidence that lithology can be more important in estimating
2223 sediment supply than topography.

2224 Rachels et al. (2020) used sediment source fingerprinting techniques to quantify the proportional
2225 relationship of sediment sources (hillslope, roads, streambanks) in harvested and un-harvested
2226 watersheds of the Oregon Coast Range. The study included one catchment (Enos Creek) that was
2227 partially clearcut harvested in the summer of 2016 and an unharvested reference catchment
2228 (Scheele Creek) located ~3.5 km northwest of Enos Creek. The paired watersheds had similar
2229 road networks, drainage areas, lithologies and topographies. The treatment watershed was
2230 harvested with a skyline buffer technique in the summer of 2016 under the Oregon Forest
2231 practices Act policy that requires a minimum 15 m no-cut buffer. The proportion of suspended
2232 sediment sources were similar in the harvested ($90.3 \pm 3.4\%$ from stream bank; $7.1 \pm 3.1\%$ from
2233 hillslope) and unharvest ($93.1 \pm 1.8\%$ from streambank; $6.9 \pm 1.8\%$ from hillslope) watersheds.

2234 However, the harvested watershed contained a small portion of sediment from roads ($3.6 \pm$
2235 3.6%), while the unharvested reference watershed suspended sediment contained no sediment
2236 sourced from roads. In the harvested watersheds the sediment mass eroded from the general
2237 harvest areas (96.5 ± 57.0 g) was approximately 10 times greater than the amount trapped in the
2238 riparian buffer (9.1 ± 1.9 g), and 4.6 times greater than the amount of sediment collected from
2239 the unharvested hillslope (21.0 ± 3.3 g). These results suggest that the riparian buffer was
2240 efficient in reducing sediment erosion relative to the harvested area. The caveat of this study was
2241 the limited sample size (1 treatment, 1 paired reference watershed) and does not incorporate the
2242 effects of different watershed physiography on sediment erosion. However, it is presented here as
2243 evidence that the formation of roads within a riparian area may interact with timber harvest to
2244 increase the potential flow of sediments from roads.

2245 Litschert & MacDonald, (2009) investigated the frequency of sediment delivery pathways in
2246 riparian management areas and their physical characteristics and connectivity following harvest.
2247 In this study the authors describe sediment delivery pathways (“features”) as rills, gullies, and
2248 sediment plumes that form when excess sediment relative to overland flows transports sediment
2249 from the hillslope to the stream. The authors surveyed 200 riparian management areas (RMA) in
2250 four different National Forests of the Sierra Nevada and Cascade Mountains of California. USFS
2251 policy requires 90-m wide RMA along each side of perennial streams and 45-m wide RMA along
2252 each side of all ephemeral and intermittent streams. When features were found within an RMA,
2253 data for years since harvest, soil depth, soil erodibility (K), feature length, feature gradient,
2254 aspect, elevation, hillslope gradient, hillslope curvature, surface roughness, and connectivity
2255 were recorded for analysis. Association between these variables were analyzed with a
2256 Spearman’s rank correlation. The variables most strongly associated with feature length were
2257 used to develop a multiple linear regression model to predict feature length. Only 19 of the 200
2258 harvest units had sediment development pathways. Feature pathways ranged in age (time since
2259 harvest) from 2 to 18 years, and in length from 10 m to 220 m. Of the 19 feature pathways, only
2260 six were connected to streams, and five of those originated from skid trails. Feature pathway
2261 length was significantly related to mean annual precipitation, cosine of the aspect, elevation, and
2262 hillslope gradient ($R^2 = 64\%$, $p = 0.004$). These results suggest that within treated riparian areas
2263 topographic characteristics such as aspect, elevation and hillslope gradient can affect delivery of
2264 sediment into streams.

2265 Rashin et al. (2006) evaluated the effectiveness of Washington State best management practices
2266 (BMPs) for controlling sediment related water quality impacts. Although this study was
2267 published in 2006, the data analyzed in this study were collected between 1992 and 1995. In their
2268 evaluation, Rashin et al. (2006) assessed site erosion, sediment delivery, channel disturbance,
2269 and aquatic habitat condition within the first two years of harvest along fish- and ~~non-fish~~
2270 ~~bearing~~non-fish-bearing streams across Washington state. From their results, the authors
2271 concluded that the site-specific factors influencing the effectiveness of BMPs in preventing
2272 chronic sediment delivery into streams were 1) the proximity of ground disturbance to the
2273 stream, 2) presence of a stream buffer, 3) falling and yarding practices that minimized
2274 disturbance to stream channel, and 4) timing of harvest activities for certain climate zones where
2275 frozen ground or snow cover may be exploited. The landscape factors that influenced BMP

2276 effectiveness were 1) the density (specific metric not reported) of unbuffered small streams at
2277 harvest sites, and 2) steepness of stream valley slopes. The authors conclude with a
2278 recommendation of excluding timber falling and yarding activities at least 10 m from streams
2279 and outside of steep inner gorges.

2280 From the studies reviewed there is evidence that sediment delivery into streams following timber
2281 harvest is influenced by not only the intensity of the harvest operation (e.g., presence of retention
2282 buffers, yarding and equipment use immediately adjacent to the stream, upland clearcut vs.
2283 thinning), but also by physiography (especially hillslope gradient), lithology relative softness,
2284 and the presence of roads. Thus, the change in magnitude of sediment delivery following harvest
2285 is context dependent and these landscape factors can interact with one another to compound
2286 these changes. However, from the studies reviewed in the sediment section of the literature
2287 review, there is evidence that the implementation of BMPs since the 1970s in the northwestern
2288 United States has lessened the impact and duration of these changes.

2289 *Nutrient*

2290 None of the studies published since 2000 and conducted in western North America provide
2291 experimental evidence of the effects of site factors on nutrient flux into streams. However, Zhang
2292 et al. (2010) conducted a global review and meta-analysis of the effectiveness of buffers in
2293 reducing nonpoint source pollution. They reported slope (hillslope gradient) as having a linear
2294 relationship with buffer pollutant removal efficacy that switched from positive to negative when
2295 slope increased beyond 10% (i.e., hillslope gradients of ~10% were optimal for buffer efficacy in
2296 removing pollutants).

2297

2298 *Focal Question 3*

2299 *3. What is the frequency of weather-related effects (e.g., windthrow, ice storms, excessive heat,*
2300 *flood and drought events) on riparian areas? What are the weather-related effects (positive and*
2301 *negative) on the riparian functions, and how are they distinguished from harvest effects? How do*
2302 *these effects differ between treated and untreated riparian forests?*

2303 The first part of this question “What is the frequency of weather-related effects (e.g., windthrow,
2304 ice storms, excessive heat, flood and drought events) on riparian areas?” is a generally worded
2305 question asking how often weather events in riparian areas occur. The second part of this
2306 question “What are the weather-related effects (positive and negative) on the riparian functions,
2307 and how are they distinguished from harvest effects?” contains within it 2 parts 1) what the
2308 effects on the riparian functions are, and 2) how they are distinguished from timber harvest
2309 effect. Any study reviewed that answers one or more parts of this question have been included.

2310 *Shade*

2311 McIntyre et al. (2021), the “Hard Rock” study, compared changes in shade from pre- to post-
2312 harvest between three riparian harvest treatments and a reference. Treatments included a two-
2313 sided 50-ft riparian buffer along at least 50% of the stream (FP; with clearcut to stream’s edge
2314 outside of the buffer), a two-sided 50-ft buffer along the entire stream (100%), and a clearcut to

2315 stream without a buffer (0%). The canopy cover was measured 1 meter above the stream surface
 2316 with a spherical densiometer. The changes in canopy cover were distinguished from harvest
 2317 effects and compared to unharvested reference sites by using a BACI design. For the FP
 2318 treatment, mean canopy cover declined from 96% to 72% in the first-year post-harvest but
 2319 continued to decline for 4 years to a minimum of 54%. In the 100% treatment mean canopy
 2320 cover was more stable, decreasing from 94% to 88% in the first year and reaching a minimum of
 2321 82% also by year 4. Canopy cover began to increase after year 4 through year 9 in both
 2322 treatments. In contrast, the reference sites experienced much smaller reductions in canopy cover
 2323 from 95% to 89% in the first four years. The cause of mortality in the treatment sites was
 2324 primarily attributed to windthrow. However, while post-harvest mortality in the treatment sites
 2325 were higher on average than in the reference sites there was a high amount of variability between
 2326 sites in both the treated and reference sites. For example, in the first 2 years following harvest
 2327 mortality ranged from 1.8 to 34.6% (loss of basal area) between sites in the FP treatment. In
 2328 contrast, mortality in the reference sites ranged from 1.1 to 20.4% (loss of basal area) during the
 2329 same period.

2330 *Litter*

2331 Bilby & Heffner (2016) showed evidence that wind speed has a strong effect on the width of
 2332 litter delivery areas within riparian areas. They used a combination of field experiments and
 2333 simulation modeling to estimate the influence of different site factors (physiography, stand age,
 2334 species composition, wind speed) on litter delivery into streams. Their results showed that litter
 2335 travel distance was also linearly related to wind speed ($p < 0.0001$). Doubling wind speed at one
 2336 site led to a 67-87% expansion of the riparian litter contribution zone in the study area. However,
 2337 this study does not compare the differences in the influence of wind speed on the width of the
 2338 litter contributing area between harvested and unharvested sites.

2339 *LW*

2340 Table 16. Treatment and responses for selected publications investigating Large Wood relevant to
 2341 Q3.

Reference	Treatment	Response
McIntyre et al. (2021)	50 feet (100%); 50 feet with clearcut gaps (FP); Clearcut to stream (unbuffered)	<u>8-years post-harvest data 100% (n = 4), FP (n = 3), and unbuffered treatments (n = 4)</u> The FP-Reference contrast in mortality was not significant 2 years post-harvest, but it was at 5- and 8-years post-harvest as mortality in FP increased relative to the Reference over time. Wind/physical damage was the primary cause of mortality for all treatments, including the Reference. In the 100% treatment it accounted for 78% and 90% of the loss of basal area and density (stem/ha), respectively; in the FP it accounted for 78% and 65%, in the reference it accounted for 52% and 43%.
Liquori (2006)	Buffer widths ranging from 25-100 feet	<u>3 years post-harvest (n = 20)</u> within no-cut buffers, windthrow caused mortality was up to 3 times greater than competition induced mortality for 3 years following treatment with tree fall probability highest in the outer areas (closest to upland clearcuts) of the buffers. highest at the outside edges of buffers (50+ feet), ~ 60% of total treefall, ~18% in the 0 -25-foot zone, and ~22% in the 25-50-foot zone.

Martin & Grotenfendt (2007)	Buffer widths 20 m or greater	Differences in mortality for the treatment sites were similar to the reference sites for the first 0-10 m from the stream (22% increase). However, mortality in the outer half of the buffers (10-20 m) from the stream in the treatment sites was more than double (120% increase) what was observed in the reference sites. The authors estimate that windthrow mortality was twofold and fivefold greater in the inner and outer halves of the treatment buffers than in the reference buffers, respectively.
Bahuguna et al. (2010)	Buffer widths 10 m, and 30 m	<u>7-years post-harvest (n = 3)</u> In the first 2 years , 11% of the timber was blown down in the 10 m buffer, compared to 4% in the 30 m buffer, and 1% in the unharvested controls. Following 8 years post-harvest , a significant amount of annual windthrow caused mortality occurred in the unharvested control at 30%, compared to 15% in both 30 m and 10 m buffers.
Schuettt-Hames & Stewart (2011, 2019b)	Buffer widths 50 feet	<u>10-years post-harvest</u> 3 years after treatment annual tree fall rates (live and dead) were over 8 times (by % of standing trees) and 5 times (by trees/acre/yr) higher in the 50-foot buffers than in the reference. 4-5 years after treatment mortality was still higher in the treated sites (27.3%) than in the reference (13.6%), but the difference was not significant. 10 years after treatment stand mortality in the 50-ft buffer treatment stabilized.

2342

2343 Chapter 3 of the Hard Rock study compared changes in stand mortality and LW input from pre-
 2344 to post-harvest and between treated and untreated reference sites. Results showed that by year 8,
 2345 post-harvest mortality as a percentage of pre-harvest basal area was lower in the reference
 2346 (16.1%) than in the 100% (24.3%) and FP (50.8%) treatments. The FP–Reference contrast in
 2347 mortality was not significant 2 years post-harvest, but it was at 5- and 8-years post-harvest as
 2348 mortality in FP increased relative to the Reference over time. The contrast in mortality between
 2349 the 100% and Reference were not significant for any time interval 8 years post-harvest.
 2350 Wind/physical damage was the primary cause of mortality for all treatments, including the
 2351 Reference. In the 100% treatment it accounted for 78% and 90% of the loss of basal area and
 2352 density (stem/ha), respectively; in FP it accounted for 78% and 65% of the loss. Wind accounted
 2353 for a smaller proportion of mortality in the reference (52% and 43%, respectively).
 2354

2355 LW recruitment to the channel was greater in the 100% and FP treatment than in the reference for
 2356 each pre- to post-harvest time interval. Eight years post-harvest mean recruitment of large wood
 2357 volume was two to nearly three times greater in 100% and FPB RMZs than in the references.
 2358 Annual LW recruitment rates were greatest during the first two years, then decreased. However,
 2359 there was a great deal of variability in recruitment rates within treatment sites and the differences
 2360 between treatments were not significant. Mean LW loading into the channel (pieces/m of channel
 2361 length) differed significantly between treatments in the magnitude of change over time. There
 2362 was a 66%, 44% and 47% increase in mean large wood density in the 100%, FP and 0%
 2363 treatments, respectively, in the first 2 years post-harvest compared with the pre-harvest period
 2364 and after controlling for temporal changes in the references. By year 8, only the FP treatment
 2365 showed a significantly higher proportional increase (41%) in wood loading when compared to

2366 the reference. In the time interval 2-8 years post-harvest wood loading in the 100% treatment
2367 stabilized.

2368
2369 Liquori (2006) investigated treefall characteristics within riparian buffer sites in a managed tree
2370 farm in the Cascade Mountains of western Washington. Buffer widths ranged between 25-100
2371 feet along ~~non-fish-bearing~~non-fish-bearing and ~~fish-bearing~~fish-bearing streams. Results
2372 showed that within no-cut buffers, windthrow caused mortality was up to 3 times greater than
2373 competition induced mortality for 3 years following treatment with tree fall probability highest in
2374 the outer areas (closest to upland clearcuts) of the buffers. Their results showed that treefall was
2375 generally highest at the outside edges of buffers (50+ feet), representing about 60% of the total
2376 observed treefall, while the 0–25-foot zone represented ~18%, and the 25–50-foot zone
2377 represented ~22%. The researchers interpret these results as evidence that windthrow
2378 susceptibility within riparian buffers increases with increasing distance from the stream.

2379
2380 Martin & Grotenfendt (2007) compared riparian stand mortality and in-stream LW recruitment
2381 characteristics between riparian buffer strips with upland timber harvest and riparian stands of
2382 unharvested watersheds using aerial photography in the northern and southern portions of
2383 Southeast Alaska. All buffer strips in this study were a minimum of 20 m wide and included
2384 selective harvest within the 20 m zone (thinning intensity not specified or included in the
2385 analyses as an effect). The results from this study showed significantly higher mortality (based
2386 on cumulative stand mortality: downed tree counts divided by standing tree counts + downed tree
2387 counts by number/ha), significantly lower stand density (269 trees/ha in buffer units and 328
2388 trees/ha in reference units), and a significantly higher proportion of LW recruitment from the
2389 buffer zones of the treatment sites than in the reference sites. Also, results showed that mortality
2390 varied with distance to the stream. Differences in mortality for the treatment sites were similar to
2391 the reference sites for the first 0-10 m from the stream (only a 22% increase in the treated sites).
2392 However, mortality in the outer half of the stream buffers (10-20 m) across treatment sites was
2393 more than double (120% increase) that observed within the reference sites. The authors estimate
2394 that windthrow mortality was twofold and fivefold greater in the inner and outer halves of the
2395 treatment buffers than in the reference buffers, respectively.

2396
2397 Bahuguna et al. (2010) evaluated the difference in windthrow caused mortality between 10 m, 30
2398 m buffer widths (neither had thinning within the buffer and both had upland clear-cuts) and
2399 unharvested controls in the Coast Mountains, British Columbia. Following harvest, 11% of
2400 initially standing timber was blown down in the first and second years in the 10 m buffer,
2401 compared to 4% in the 30 m buffer, and 1% in the unharvested controls. However, after 8 years
2402 post-harvest, a significant amount of annual mortality occurred when winter storms brought
2403 down multiple trees in the unharvested control at 30%, compared to 15% in both 30 m and 10 m
2404 buffers. These results show evidence that timber harvest can increase windthrow caused
2405 mortality within protective buffers in the short term but can stabilize within a decade. Further,
2406 this study shows evidence that windthrow caused mortality is stochastic and large storm events
2407 can cause significant mortality within untreated riparian forests.

2408

2409 Schuett-Hames and Stewart (2019a) compared changes in stand mortality and LW recruitment
2410 between treated and untreated riparian areas along fish-bearing streams in eastern Washington.
2411 Treatments were prescribed under the Standard Shade Rule (SR), under the All-Available Shade
2412 rule (AAS), and unharvested reference sites. Both shade rules have a 30-ft no-cut buffer (core
2413 zone) immediately adjacent to the stream. The SR prescription allows thinning in the buffer zone
2414 30-75 feet (inner zone) from the stream while the AAS prescription requires retention of all
2415 shade providing trees in this area. Thinning non-shade providing trees within the inner zone is
2416 allowed under the AAS rule. Results from a mixed model comparison showed that the frequency
2417 of wood input from fallen trees was significantly greater in SR group compared to both the
2418 reference and AAS groups ($p < 0.001$), while the difference between reference and AAS groups
2419 was not significant. Over 60% of pieces recruited from AAS and SR fallen trees consisted of
2420 stems with attached rootwads (SWAR), double the proportion in the reference sites. The
2421 reference-AAS and reference-SR differences in recruitment of SWAR pieces were significant (p
2422 < 0.001). The authors comment that the higher mortality and recruitment of LW in the SR sites
2423 was primarily due to windthrow.

2424 Schuett-Hames et al, (2011) compared tree mortality and LW recruitment between treated and
2425 untreated riparian stands along ~~non-fish-bearing~~non-fish-bearing streams in western Washington.
2426 Treated sites were prescribed a 50-foot-wide no-cut buffer. Annual fall rates of live and dead
2427 standing stems combined were over 8 times (by % of standing trees) and 5 times (by
2428 trees/acre/yr) higher in the 50-foot buffers than in the reference buffers 3 years after treatment.
2429 These differences were significant for both metrics ($p < 0.001$). Over the entire five-year period,
2430 the percentages of standing trees that were uprooted and broken (as well as the combined total)
2431 were significantly greater in the 50-foot buffer. Wind was the dominant tree fall process,
2432 accounting for nearly 75% of combined fallen trees, 11% fell from other trees falling against
2433 them and 1.8% of fallen trees fell from bank erosion. Differences in mortality followed a similar
2434 pattern to tree fall rates. In the 50-foot buffer sites mortality rates were significantly higher (3.5
2435 times higher) than in the reference sites for the first three years following harvest. However, in
2436 years 4-5 mortality rates increased in the reference buffers after high-intensity storms resulting in
2437 non-significant differences in mortality during this period. The cumulative percentage of live
2438 trees that died over the entire five-year period was 27.3% in the 50-ft buffers compared to 13.6%
2439 in the reference reaches, but the difference was not statistically significant. The authors suggest
2440 that the lack of significance was likely due to the high variability in mortality between sites in
2441 the 50-foot buffers.

2442 In the follow-up study, Schuett-Hames & Stewart (2019b) reported that over a 10-year period,
2443 stand mortality in the 50-ft buffer treatment stabilized and showed a cumulative 14.1% reduction
2444 in live basal, while the reference stands showed a 2.7% increase in live basal area. The
2445 differences in these values were not significant. Cumulative LW recruited into stream channel
2446 over the 10-period was double in the 50-ft buffer treatment streams than in the reference streams.

2447 In general, the studies reviewed above show evidence that upland timber harvest with riparian
2448 retention buffers initially increases stand mortality within the buffers and increases LW
2449 recruitment relative to unharvested reference stands in the short-term. Hence, treated riparian
2450 forests appear to have a higher susceptibility to windthrow caused mortality, at least in the short

2451 term, compared to untreated stands. Depending on the streams in question, an increase in LW
 2452 could be considered a positive or negative impact This increase in mortality and LW recruitment
 2453 is attributed to an increase in the susceptibility to windthrow within the riparian buffers relative
 2454 to the unharvested controls. Further, multiple studies (Liquori, 2006; Martin & Grotefendt, 2007,
 2455 Schuett-Hames & Stewart 2019a) showed evidence that the increase in windthrow caused
 2456 mortality is highest in the outer area of the riparian buffers (area closest to upland treatments).
 2457 There is some evidence that thinning within the buffer can also affect mortality rates, but these
 2458 studies are few. In the three studies that collected post-harvest data for 8 or more years
 2459 (Bahuguna et al., 2010; McIntyre et al., 2021; Schuett-Hames & Stewart 2019b), there is
 2460 indication that mortality in the riparian buffers and annual LW recruitment into adjacent streams
 2461 stabilizes within 5-10 years. However, in the subsequent decades following treatments with
 2462 upland clearcuts there is evidence that LW recruitment rates can continue to decrease and in
 2463 stream wood loads may become depleted before recruitment rates can recover (Nowakowski &
 2464 Wohl, 2008; Reid & Hassan, 2020) depending on applied management practices (e.g., buffer
 2465 widths, road construction, etc.). For example, Teply et al. (2007) used simulation modeling to
 2466 estimate the effectiveness of Idaho Forest Practices for riparian buffers and found no significant
 2467 difference between predicted LW loads for harvested and unharvested sites 30-, 60-, or 100-years
 2468 post-harvest.

2469 *Nutrient*

2470 Table 17. Treatment and responses for selected publications investigating Nutrients relevant to
 2471 Q3.

Reference	Treatment	Response
Vanderbilt et al. (2003)	long-term datasets from six watersheds in the H.J. Andrews Experimental Watershed	<u>20-30 years of historical data</u> Total annual discharge was a positive predictor of annual dissolved organic nitrogen (DON) export in all watersheds with R2 values ranging between 0.42 to 0.79. No other nutrients nitrate (NO3-N), ammonium (NH4-N), and particulate organic nitrogen (PON) showed consistent patterns or relationships to any predictor variables. The increase in concentration began in July or August with the earliest rain events , and peak DON concentrations occurred in October through December before the peak in the hydrograph. DON concentrations then declined during the winter months .
Yang et al. (2021)	Mastication of riparian area shrubs to < 10% cover. Treatment effects compared with drought effects.	2 years pre-drought, 3 years following drought and treatment Drought alone altered the concentration of dissolved organic carbon (DOC) in stream water. Dissolved organic carbon (DOC) was 62% lower and the ratio of DOC to dissolved inorganic nitrogen (DIN) was 82% lower during drought years. Drought combined with thinning showed 66- 94% higher DOC than in unthinned watersheds.

2472

2473 Vanderbilt et al. (2003) analyzed long-term datasets (ranging 20-30 years for each watershed)
 2474 from six watersheds in the H.J. Andrews Experimental Watershed in the west-central Cascade

2475 Mountains of Oregon to investigate patterns in dissolved organic nitrogen (DON) and dissolved
2476 inorganic nitrogen (DIN) export with watershed hydrology. The researchers used regression
2477 analysis of annual N inputs and outputs with annual precipitation and stream discharge to
2478 analyze patterns. Their results showed that total annual discharge was a positive predictor of
2479 annual DON export in all watersheds with R^2 values ranging between 0.42 to 0.79. In contrast,
2480 relationships between total annual discharge and annual export of nitrate ($\text{NO}_3\text{-N}$), ammonium
2481 ($\text{NH}_4\text{-N}$), and particulate organic nitrogen (PON) were variable and inconsistent across
2482 watersheds. The authors speculate that different factors may control organic vs. inorganic N
2483 export. The authors emphasize the importance of analyzing data from multiple watersheds in a
2484 single climactic zone to make inferences about stream chemistry.

2485 Yang et al. (2021) investigated the effects of drought and forest thinning operations
2486 (independently and combined) on stream water chemistry in the Mediterranean climate
2487 headwater basins of the Sierra National Forest. The effects of drought alone were examined by
2488 comparing water samples collected from control watersheds for 2 years before and 3 years after
2489 drought. The effects of drought and thinning combined were examined by comparing water
2490 samples collected from treated sites to reference sites for three years post-harvest (all drought
2491 years). Drought alone altered the concentration of dissolved organic carbon (DOC) in stream
2492 water. Volume-weighted concentration of DOC was 62% lower ($p < 0.01$) and the ratio of
2493 dissolved organic carbon to dissolved inorganic nitrogen (DOC:DON) was 82% lower ($p =$
2494 0.004) in stream water in years during drought (WY 2013–2015) than in years prior to drought
2495 (WY 2009 and 2010). Drought combined with thinning altered DOC and DIN concentrations in
2496 stream. For stream water, volume-weighted concentrations of DOC were 66–94% higher in
2497 thinned watersheds than in control watersheds for all three consecutive drought years following
2498 thinning. No differences in DOC concentrations were found between thinned and control
2499 watersheds before thinning. The authors conclude that their results showed evidence that the
2500 influences of drought and thinning are more pronounced for DOC than for DIN in streams.

2501 *Drought Frequency*

2502 Wise (2010) used reconstructed newly collected tree-ring data augmented with existing
2503 chronologies from sites at three headwater streams in the Snake River Basin to estimate
2504 streamflow patterns for the 1600–2005 time-period. Streamflow patterns derived from
2505 instrumental data and from reconstructed chronologies were compared with other streamflow
2506 previously reconstructions of three other western rivers (the upper Colorado, the Sacramento,
2507 and the Verde Rivers) in similar climates to examine synchronicity among the rivers and gain
2508 insight into possible climatic controls on drought episodes. The reconstruction model developed
2509 for the analysis explained 62% of the variance in the instrumental record after adjustment for
2510 degrees of freedom. Results showed evidence that droughts of the recent past are not yet as
2511 severe, in terms of overall magnitude, as a 30-year extended period of drought discovered in the
2512 mid-1600s. However, in terms of number of individual years of $< 60\%$ mean-flow (i.e., low-flow
2513 years), the period from 1977–2001 were the most severe. Considering the frequency of
2514 consecutive drought years, the longest (7-year-droughts), occurred in the early 17th and 18th
2515 centuries. However, the 5-year drought period from 2000–2004 was the second driest period over

2516 the 415-year period examined. The correlative analysis of the chronologies developed for the
2517 upper Snake River with other rivers of the West showed mixed results with periods of positive
2518 and negative correlations. The author interprets these results as evidence that drought frequency,
2519 in general, in this area appears to be increasing in severity and that mean annual flow appears to
2520 be reducing in the latter half of the 20th and the beginning of the 21st century. The exceptions
2521 being the 1930's dustbowl, and an unusually long dry period in the early 1600s.

2522 *Fire Frequency*

2523 Dwire & Kauffman (2003) in their reviewed and summarized the available conducted on fire
2524 regimes in forested riparian areas relative to uplands in the western United States. They
2525 summarized the distinctive features of riparian areas that can influence the properties of fire as
2526 (1) higher fuel loads because of higher net primary productivity, (2) higher fuel moisture content
2527 due to proximity to water, shallow water tables, and dense shade, (3) active channels gravel bars
2528 and wet meadows may act as fuel breaks, (4) topographic position (canyon bottoms, low point on
2529 landscape) leads to higher relative humidity, fewer lightning strikes, but more human-caused
2530 ignitions, (5) microclimate may lead to cooler temperatures and higher humidity that can lessen
2531 fire intensity and spread. They highlight a need for more extensive research on the history and
2532 ecological role of fire in the riparian areas of the western United States.

2533 There is a logical assumption that fire in riparian zones would be less frequent than in adjacent
2534 uplands because of its proximity to water. However, several studies have been conducted which
2535 reconstruct historical fire regimes in riparian areas relative to adjacent uplands and have
2536 provided varying results. Everett et al. (2003) used fire-scar and stand-cohort records to estimate
2537 the frequency and seasonality of fire in Douglas-fir dominated riparian areas and adjacent
2538 uplands. They sampled sites along 49 stream segments on 24 different streams in the Wenatchee
2539 (33 segments) and Okanogan (16 segments) National Forests. The data collected allowed for
2540 reconstruction of fire occurrence back to 1896. Their results showed that the mean count of fire
2541 scars was significantly fewer in riparian areas than in adjacent uplands regardless of valley type,
2542 aspect, or plant association group. However, the difference between riparian and upland fire scars
2543 was greatest for western aspects and least for northern aspects. Also, the differences were
2544 greatest for the 'warm mesic shrub/herb' plant association group (e.g., common snowberry), and
2545 least in the cool dry grass plant association group (e.g., pinegrass, or elk sedge).

2546 Prichard et al. (2020) evaluated drivers of fire severity and fuel treatment effectiveness at the
2547 2014 Carlton Complex in north-central Washington State. While this study's objective does not
2548 specifically evaluate differences in fire severity between riparian and upland forests, it did
2549 evaluate differences in fire severity based on variations in topographic and vegetation type
2550 variables. One vegetation variable was classified broadly as "riparian vegetation" from the
2551 publicly available data set LANDFIRE. The authors used a combination of simultaneous
2552 autoregression and random forests approaches to model drivers of fire severity. In the study
2553 area's southern section (1 of 2 designated study areas), the results showed cover type was a
2554 significant predictor with negative correlations with fire severity in non-forest types and riparian
2555 forests.

2556 Conversely, Olson & Agee (2005) provide evidence that fire return intervals in the riparian areas
2557 of the Umpqua National Forests, Oregon, may not have differed significantly from adjacent
2558 upland forests. They reconstructed historical fire return intervals from fire scar cross sections
2559 taken from 15 stream reaches and 13 paired upland forests. Sites were primarily dominated by
2560 Douglas-fir, western red cedar, and western hemlock. The number of fires per plot, maximum
2561 and minimum fire return intervals, and the Weibull median fire return interval (WMPIs) were
2562 compared between riparian and upland stands using the Wilcoxon signed rank test, the Mann-
2563 Whitney U-test for unmatched samples, and the Kruskal-Wallis one-way analysis of variance.
2564 The results showed that between 1650 and 1900, 43 fire years occurred on 80 occasions. Of these
2565 80 occasions, 33 were recorded in the riparian and adjacent upslope forest, 23 were recorded in
2566 only the riparian area, and 24 were recorded only in the upland forests. The riparian WMPIs
2567 were somewhat longer (ranging from 35-39 years, with fire return intervals ranging from 4-167
2568 years) than upslope WMPIs (ranging from 27-36 years, with fire return intervals ranging from 2-
2569 110 years), but these differences were not significant. The authors, Olson & Agee (2005),
2570 interpret these results as evidence that fires in this area were likely patchy and smaller in scale
2571 with a high incidence of fires occurring only in the riparian area or only in the upland forests,
2572 and less commonly in both. The authors also suggest that fire is a natural occurrence in the
2573 riparian areas of this area and should be restored to protect riparian forest health.

2574 Another study from the Klamath Mountains in northern California showed evidence that fires in
2575 riparian forests may have been more frequent than in adjacent upland forests (Skinner, 2003).
2576 Skinner (2003) used dendrochronological methods to construct fire return intervals for 5 riparian
2577 and adjacent upland forests sites, each between 1-2 hectares. Because of the small sample size,
2578 statistical analysis was not conducted, and their results are only descriptive. The ranges of fire
2579 return intervals (FRIs) were similar between riparian and upland forests. However, the median
2580 FRI for the riparian forests was nearly double that in adjacent uplands. The authors conclude that
2581 these limited data suggest fire in the riparian areas may be more variable than in the uplands in
2582 frequency and intensity.

2583 Yet, another study from Harley et al. (2020) showed evidence that the differential fire occurrence
2584 riparian and adjacent uplands may have been dependent on weather (i.e. drought). Harley et al.
2585 (2020) reconstructed low-severity fire histories from tree rings in 38 1-ha plots. This data was
2586 supplemented with existing fire histories from 104 adjacent upland plots. 2633 fire scars were
2587 sampled from 454 (127 riparian; 329 upland) trees from two sites in the Blue Mountains in
2588 north-eastern Oregon: One in the Wallowa-Whitman (WWNF) and one in the Malheur (MNF)
2589 National Forests. Fire-scar dates were used to construct plot composite fire chronologies,
2590 excluding fire dates recorded from only one tree. These were used to compute median fire
2591 intervals for riparian and upland forests for each site and for both sites combined. A mixed linear
2592 model with fire interval as a response and plot type (riparian vs. Upland) as a predictor was used
2593 to check for statistical difference in fire frequency. The influence of climate on fire occurrence
2594 was inferred by assessing whether the summer Palmer Drought Severity Index (PDSI) differed
2595 significantly during the fire year or preceding or following years (-3 to +1 years) using
2596 superimposed epoch analysis. Results showed that Fires burned synchronously in riparian and
2597 upland plots during more than half of the fire years at both WWNF and MNF (55% and 57%,

2598 respectively). At WWNF, fires burned during 65 years of the analysis period (1650–1900); 36
2599 burned in both riparian and upland plots, 7 burned only in riparian plots and 22 burned only in
2600 upland plots. At MNF, fires burned during 74 years of the analysis period; 42 burned in both
2601 riparian and upland plots, 3 burned only in riparian plots and 29 burned only in upland plots. At
2602 both sites, average PDSI was significantly warm–dry during synchronous fire years. However,
2603 climate was not significantly cool–wet during non-synchronous fire years at either site. The
2604 authors interpret these results as evidence that historical synchronized fire occurrence was more
2605 likely during excessively dry or drought years.

2606 There is also evidence that riparian forest fire regimes have been altered in many areas from pre-
2607 Euro-American settlement due to fire suppression. Messier et al. (2012), used dendro-ecological
2608 methods to reconstruct pre-Euro-American settlement riparian forest structure and fire frequency
2609 for comparison of changes post-settlement in the Rouge River of southwestern Oregon. Fire
2610 events were dated from increment cores and fire-scar cross-sections back to the year 1600,
2611 approximately. Changes in annual radial growth rates were used to infer changes in stand density
2612 over time. Results showed the age distribution prior to 1850 followed a pulse pattern of
2613 recruitment with recruitment peaks occurring around 1850, 1800, and between 1740-1770
2614 (though this pulse was difficult to discern because the sample size of trees established prior to
2615 1740 were relatively few). After 1900, many mixed conifer sites showed a dramatic increase in
2616 the recruitment of more shade tolerant white fir (*Abies concolor*) compared to Douglas-fir
2617 (*Pseudotsuga menziesii*). White fir comprised 51% of the live trees recruited after 1900, but only
2618 18% of the live trees before 1900. Results from the 26 cross-dated fire scars spanned from 1748
2619 – 1919 with the highest number of detected fires occurring in the early-settlement period (1850-
2620 1900). The authors interpret these results as evidence that fire suppression over the last century
2621 has changed the successional pathway and stand structure of riparian forests in this area.

2622 Van de Water & North (2011) found similar results from their study in the northern Sierra
2623 Nevada. They compared current field data with reconstructed data to estimate changes in stand
2624 structure, fuel loads, and potential fire behavior over time. Additionally, they estimated how
2625 these conditions for riparian forests compared to adjacent upland forests during the reconstructed
2626 and current periods. Data for current forest structure, species composition, and fuel loads were
2627 collected from 36 adjacent riparian and upland sites (72 sites total). The reconstruction period
2628 was set at the year of the last fire (ranging from 1848 – 1990), determined from fire-scar records.
2629 Potential fire behavior, effects, and canopy bulk density were estimated for current and
2630 reconstructed stand conditions for riparian and upland sites using Forest Vegetation Simulator
2631 (FVS). Stand structure (BA, stand density, snag volume, QMD, average canopy base height),
2632 species composition, fuel load, potential fire behavior, canopy bulk density, and mortality were
2633 compared between current and reconstructed periods for riparian and upland sites, and between
2634 sampling areas (riparian vs. Upland) with an analysis of variance (ANOVA). Results showed that
2635 under current conditions, riparian forests were significantly more fire prone than upland forests,
2636 with greater stand density (635 vs. 401 stems/ha), probability of torching (0.45 vs. 0.22),
2637 predicted mortality (31% vs. 16% BA), and lower quadratic mean diameter (46 vs. 55 cm),
2638 canopy base height (6.7 vs. 9.4 m), and frequency of fire tolerant species (13% vs. 36% BA).
2639 However, the reconstructed periods showed no significant difference between riparian and

2640 upland forests for fuels and structure. The authors suggest that these results provide evidence that
 2641 the historic fire return intervals may not have differed significantly between riparian and upland
 2642 forests in this area.

2643 *Fire Effects on Function*

2644 *Litter and Nutrients*

2645 Table 17. Treatment and responses for selected publications investigating the relationship
 2646 between wildfire Litter and Nutrients relevant to Q3.

Reference	Treatment	Response
Musetta-Lambert et al. (2017)	Buffer widths 30 m; wildfire	<u>Sampling began 7-17 years after harvest or 12 years after wildfire (n = 5 harvest, 7 fire, 6 reference)</u> Total leaf litter input was significantly higher at fire sites than at harvested or reference sites. Fire sites had significantly greater leaf litter inputs by species of willow spp., Atlantic ninebark, and speckled alder than in both reference and harvested sites
Rhoades et al. (2011)	Wildfire	<u>1- year pre- and 5-years following the 2002 Hayman Fire in Colorado</u> Cation concentrations and acid neutralizing capacity (ANC) increased immediately and significantly following fire that peaked at 4 months . Ca 2+ concentrations, ANC, and conductivity remained elevated in the burned streams for 2 years compared to pre-fire conditions, and unburned streams. Stream water nitrate and turbidity increased linearly with the proportion of a basin burned or burned at high severity.
Son et al. (2015)	Wildfire	<u>2-years pre- and immediately following wildfire</u> Results for turbidity showed no significant differences between pre- and post-fire ranges immediately following fire . After first rainfall event mean turbidity increased from 11.3 NTU to 641.62 NTU. Post-fire aqueous total phosphorus (TP) and nitrogen (TN) was significantly higher than pre-fire values (390 and 6 times higher than pre-fire values for TP and TN, respectively).

2647
 2648 Musetta-Lambert et al. (2017) compared changes in leaf-litter inputs into streams following
 2649 adjacent riparian forest harvesting or wildfire to reference sites. This study took place in the
 2650 boreal forest of the White River Forest management Area in Ontario, Canada, ~75 km inland
 2651 from the northern shore of Lake Superior. This study is outside of western North America (the
 2652 focal area for this review), but it is the only study found that provides experimental evidence of
 2653 wildfire's effects on litter inputs. The study sites consisted of ~50 m reaches in 25 catchments, 10
 2654 that were harvested, 7 that experienced wildfire, and 8 references. Of these reaches a subset was
 2655 used to riparian forest structure, leaf litter inputs, and water chemistry (5 harvest, 7 fire, 6
 2656 reference). The harvested catchments were harvested 7-17 years prior to the study (minimum 30
 2657 m riparian buffers; specific harvest rules/methods not described). The wildfire catchments had
 2658 burned 12 years prior to the study and had no dead material removed. The reference catchments
 2659 had no fire or harvesting for a minimum of 40 years. Water grab samples were collected in
 2660 September, October and November 2010, and May, June and September of 2011 from the study
 2661 reaches.

2662 Water samples were analyzed to obtain measurements for pH, conductivity, dissolved organic
2663 carbon (DOC) and dissolved inorganic carbon (DIC) concentrations, soluble reactive
2664 phosphorous (SRP), along with a suite of other major elements and nutrient measurements (total
2665 N, NH₄, total P, Ca, K, Mg, etc.). Vertical leaf litter traps consisting of plastic bins were placed at
2666 10 locations along the bankfull width of each site. Lateral leaf fall was not collected or analyzed.
2667 Leaf litter inputs were focused on leaves from deciduous trees and shrubs. Leaves were separated
2668 to the lowest possible taxonomic level, dried and weighed for analysis.

2669 Univariate one-way ANOVA models were used to determine differences in water chemistry,
2670 riparian forest characteristics of juvenile tree and shrub communities (richness, Shannon's
2671 diversity index, relative occurrence of individual taxa), mature tree communities (total basal
2672 area, stem density), and litter subsidies (richness, mass input). Results for water chemistry
2673 showed that Conductivity, pH, and dissolved inorganic carbon were significantly higher at fire
2674 sites than at reference sites ($p = 0.02$, $p = 0.04$, $p = 0.03$, respectively) but did not differ between
2675 harvested and fire sites or harvested and reference sites.

2676 Results for stand structure showed there was significantly higher taxa richness in fire sites than
2677 in reference sites or harvested sites ($p = 0.04$). Taxa richness did not differ significantly between
2678 reference and harvested sites. Reference sites had significantly higher total mean densities (# ha
2679 ⁻¹) of mature riparian trees (>10 cm DBH) than fire ($p < 0.001$) and harvested sites ($p = 0.036$).
2680 Total mature tree densities in reference sites were 1.7x and 4x higher than in harvested and fire
2681 sites, respectively. Taxa richness in leaf litter subsidies did not significantly differ among
2682 disturbances ($p = 0.477$). Total leaf litter input (g m^{-1}) significantly higher at fire sites than at
2683 harvest ($p = 0.02$) or reference sites ($p = 0.02$). Fire sites had significantly greater leaf litter
2684 inputs of willow spp. ($p = 0.0002$, 0.006 , respectively), Atlantic ninebark ($p = 0.002$, 0.003 ,
2685 respectively) and speckled alder ($p = 0.02$, 0.04 , respectively) than in both reference and
2686 harvested sites. The authors interpret these results as evidence that natural fire disturbance in
2687 low-order boreal forest streams had higher leaf litter inputs, and different stand structures and
2688 composition than harvested or untreated riparian stands. They suggest that while harvested
2689 stands were more structurally similar to fire affected stands than reference stands, the future
2690 implementation of these treatments should intend to emulate the patchy nature of wildfire
2691 disturbance. This would enhance the diversity of riparian forest structure and increase litter
2692 subsidies into streams.

2693 *Nutrients*

2694 Rhoades et al. (2011) monitored stream chemistry and sediment 1-year before and for 5-years
2695 after the 2002 Hayman Fire in Colorado. Monthly water samples were collected from streams in
2696 three burned and three unburned watersheds. Pre-fire and post-fire water nitrate, cation
2697 concentration (Ca^{2+} , Mg^{2+} , K^{+}), acid neutralizing capacity (ANC) and turbidity were compared
2698 graphically and statistically between the three burned and unburned basins. Results for cation
2699 concentrations and ANC showed an immediate and significant increase that peaked during the 4-
2700 month period following the fire. The Ca^{2+} concentrations, ANC, and conductivity remained
2701 elevated in the burned streams for 2 years compared to pre-fire conditions, and unburned
2702 streams. Stream water nitrate and turbidity increased linearly with the proportion of a basin

2703 burned or burned at high severity. No other chemical analyte showed a significant response to
2704 fire severity or extent. Streams draining basins affected by extensive stand-replacement fires
2705 showed a 3.3-fold higher ($p=0.000$) nitrate concentration than basins that burned less. Also,
2706 turbidity was 2.4-fold ($p=0.000$) higher average turbidity compared to streams in basins burned
2707 less severely or extensively. In the extensively burned basins, stream water nitrate concentrations
2708 did not decline over the five years of the study and the mean concentrations of nitrate in the fifth
2709 year did not differ from the fourth year. The authors conclude that wildfire can have immediate
2710 and mid-term (up to 5 years) impacts on water chemistry and turbidity. Further, the magnitude
2711 and temporal increases of nitrate and turbidity, specifically, have a positive relationship with burn
2712 severity and extent.

2713 Son et al. (2015) compared stream water samples before and after an intense wildfire in the
2714 Cache la Poudre River basin in Colorado. Stream water samples for total phosphorus (TP) and
2715 total nitrogen (TN) were collected over 2 years (2010 – May 2012) before the fire in June 2012.
2716 Two post-fire water samples were taken: 1) immediately following containment of the fire (July
2717 4, 2012) and 2) twelve days after the fire was contained (July 16, 2012). For each pre- and post-
2718 fire sampling date water samples were collected at three randomly selected points at two sites.
2719 Riverbed sediments were also collected at each site and sieved through a 2 mm sieve to capture
2720 the geochemically reactive portion of the riverbed. The pre- and post-fire sediment and stream
2721 water quality were compared with t-test. Correlations of sediment and stream water quality with
2722 other factors (e.g., stream temperature, precipitation, streamflow) were evaluated with a
2723 Pearson's correlation at 0.05 and 0.1 significance levels. Results for turbidity showed no
2724 significant differences between pre- and post-fire ranges immediately following fire. However,
2725 after the first post-fire rainfall (2.5 mm) nephelometric turbidity ranged from 113.6 - 2099.4
2726 NTU (mean = 641.62 NTU), a considerable increase from pre-fire data (mean 11.3 NTU), and
2727 post-fire data before rainfall (47.3 NTU). Post-fire aqueous TP and TN loads ranged from 30.5 -
2728 56,086 and 45.4 - 1203 kg/day, respectively, and were significantly higher than pre-fire values
2729 (390 and 6 times higher than pre-fire values for TP and TN, respectively). The authors note that
2730 this is likely due to the transport and input of ash into the stream. After the first rainfall, all forms
2731 of P were significantly higher than pre-fire concentrations, such as soluble reactive phosphorus
2732 (SRP; $p=0.000$), dissolved organic phosphorus (DOP; $p=0.009$), and particulate phosphorus
2733 (PP; $p=0.02$). Riverbed sediment equilibrium P concentrations increased significantly ($p=$
2734 0.007) from pre- to post-fire in all sites. The authors conclude that this study shows evidence that
2735 stream TP and TN, and riverbed sediment TP all increased significantly after the first rainfall,
2736 post-fire. They further suggest that the effects of wildfire on riverbed sorption mechanisms are
2737 very complex but further research would be valuable because fire impacted sediments highly
2738 concentrated P can become a long-term source of P.

2739 *LW*

2740 Bendix & Cowell (2010) investigated the effects of fire and flooding on LW input in two
2741 tributaries of Sespe Creek (Potrero John Creek and Piedra Blanca Creek) in the Los Padres
2742 national Forest in southern California. Both sites were located within the perimeter of the Wolf
2743 Fire that burned in June of 2002. Extensive flooding in the area occurred during January and

2744 February of 2005. The study area is characterized by chaparral dominated communities and a
2745 Mediterranean-type climate. While there is a scarcity of trees in the uplands, the riparian areas
2746 contained substantial growth of *Alnus rhombifolia* (white alder), *Populus fremontii* (Fremont
2747 cottonwood), *Quercus agrifolia* (coast live oak), *Quercus dumosa* (scrub oak) and *Salix* sp.
2748 (willows) on the valley floors. Thus, any change in in-stream or riparian area LW was sourced
2749 exclusively from the riparian area. Data for LW and standing live and dead stems in the riparian
2750 area were collected in July, of 2003 (1-year pre-fire) and again in July of 2005 (3-years post-fire,
2751 5-6 months after flood events). This data was used to answer 4 questions: 1) How many of the
2752 burned snags fell during this time, and what was the species composition?, 2) Did snags differ by
2753 species or size in the rate at which they fell?, 3) How did flooding after the fire affect the rate at
2754 which snags fell?, 4) How did flooding affect the mobilization of fallen snags? Questions 1 was
2755 analyzed by comparing descriptive data (i.e., no statistical analysis). A t-test was used to compare
2756 mean diameter of standing and fallen stems (question 2). T-tests were also used to analyze
2757 differences in mean flow depth for standing vs. fallen snags and for fallen snags still present vs.
2758 snags that had been transported after flooding (questions 4 and 5). Results showed high post-fire
2759 mortality (94%) with 339 of 362 stems killed. By 2005, 57 of the 339 snags had fallen (16.8%).
2760 The majority of fallen stems were either *Alnus* or *Salix* species. Standing snags varied in size
2761 from 3 cm to 69.2 cm, whereas those that had fallen ranged from 3 cm to 33 cm. Among the
2762 fallen snags, those <10 cm were not proportionate to the overall numbers, whereas snags between
2763 10 cm and 30 cm were disproportionately likely to fall. While fewer snags in the larger size
2764 classes the mean diameter of fallen snags was larger than the mean diameter of standing snags
2765 (11.4±10.9 cm vs. 11.0±8.0 cm) and did not differ significantly. The mean flood depth for fallen
2766 snags (1.05±0.68 m) was significantly greater than those still standing (0.40±0.56 m; $p < 0.0001$,
2767 $n=339$). The three species experiencing no snagfall at all (*Abies glauca*, *Rhamnus californica* and
2768 *Quercus agrifolia*) occurred only in higher quadrats, which had experienced virtually no
2769 flooding. Of the 57 snags that had fallen by July 2005, 43 (75%) were gone from the quadrats in
2770 which they had been recorded in 2003. The snags that had been mobilized were from quadrats
2771 that had experienced deeper flood depths (1.14±0.69 m) than those that had remained. (0.80±0.62
2772 m), but the difference is insignificant. The authors interpret these findings as an indication that
2773 short-term rates of snagfall following wildfire are influenced by the species composition of
2774 burned stems and by post-fire flood depth. Thus, although wildfire resulted in many burned snags
2775 across the valley floor, the rate at which these stems are recruited into the fluvial system as
2776 woody debris varies by the ecological characteristics and the geomorphic setting.

2777

2778 Focal Question 4

2779 4. How do various treatments within riparian buffers relate to forest health and resilience to fire,
2780 disease, and other forest disturbances?

2781 While there are several studies that discuss the frequency, dynamics, or potential for
2782 disturbances, especially fire, in riparian areas of the western United States (Dwire & Kauffman,
2783 2003; Everett et al., 2003; Merschel et al., 2014) there is a dearth of studies that investigate how
2784 treatments within the riparian area or in riparian buffers relate to the riparian area's resilience to

2785 disturbance. No studies found in our literature search and review were suitable for providing
 2786 direct experimental evidence of the effects of riparian buffer treatments on riparian health and
 2787 resilience to disturbance except for several studies that provide evidence that riparian harvest
 2788 treatments have the potential to increase susceptibility to windthrow caused mortality. Post-
 2789 harvest changes in windthrow susceptibility are discussed in focal question 3. One study used
 2790 simulation modeling to estimate changes in health and susceptibility to disturbance with and
 2791 without treatment.

2792 Ceder et al. (2018) used Forest Vegetation Simulator (FVS) to predict how treatment along fish-
 2793 bearing streams of eastern Washington affects riparian stand health and susceptibility to insects,
 2794 disease, and crown fire. The projected changes in susceptibility were produced for the low- and
 2795 mid-elevation regulatory zones for timber harvest. Models were run for 50 years with and
 2796 without application of prescribed treatments. Prescriptions for these zones include a buffer width
 2797 of 75-130 ft depending on stream width category. For all treatments, no harvest is allowed within
 2798 the first 30 feet from the bankfull channel. Timber harvest is allowed in the remaining width of
 2799 the buffer but must meet a minimum basal area based on the regulatory zone. The authors report
 2800 high variability in the data and the outputs of each modeling scenario. However, they report that
 2801 overall, as riparian zone growth was simulated with and without management, tree size and stand
 2802 density increased, along with some increases in insect and disease susceptibility and potential
 2803 fire severity without management and decreases with management.

2804 **Focal Question 5**

2805 *5. How do the functions provided by riparian stands change over time (e.g., large woody debris*
 2806 *recruitment from farther away from the stream)?*

2807 This question addresses a temporal and spatial component to changes in function. The question
 2808 specifies “change over time” but provides an example with a spatial component. While harvest is
 2809 not specified as a factor, studies that quantify changes to riparian function in harvested reaches
 2810 have been included. Studies that compare differences in one or more functions between
 2811 comparable sites in different successional stages (i.e., different mean age) are also included.
 2812 Papers that investigate the changes in ~~LW~~**large wood** source distance following harvest have
 2813 been included because of the given example (*large woody debris recruitment from farther away*
 2814 *from the stream*).

2815 *Shade*

2816 Table 18. Treatment and responses for selected publications investigating Shade relevant to Q5.

Reference	Treatment	Response
Kaylor et al. (2017)	old-growth (> 300 years old) and mid-successional (50-60 years old) Douglas-	the authors estimate that canopy openness reaches its minimum value in regenerating forests at ~30 years and maintains with little variability until ~100 years . Mean canopy openness in stands 30-100 years old was 8.7% with a range from 1.2 to 32.0% (SD = 5.7). Canopy openness over streams in old-growth forests averaged 18.0% but was highly variable and ranged from 3.4 to 34.0% (SD= 5 7.9)

	fir dominated forests	
Warren et al. (2013)	old-growth-forests (>500 years old) and young second-growth stands (~40-60 years old)	Three of the four paired old-growth reaches had significantly lower mean percent canopy cover

2817

2818 Kaylor et al. (2017) compared canopy cover throughout stream networks adjacent to old-growth
 2819 (> 300 years old) and mid-successional (50-60 years old) Douglas-fir dominated forests in the
 2820 H.J. Andrews Experimental Forest in the Cascade Mountains of Oregon. Canopy openness was
 2821 quantified with a handheld spherical densiometer. Data was supplemented with a review of
 2822 literature studies conducted in the Pacific Northwest that reported stand age and canopy cover
 2823 over the stream. The combined datapoints for canopy openness (%) were plotted against stand
 2824 age and fit with a negative exponential curve. From the slope of the curve, the authors estimate
 2825 that canopy openness reaches its minimum value in regenerating forests at ~30 years and
 2826 maintains with little variability until ~100 years. Mean canopy openness in stands 30-100 years
 2827 old was 8.7% with a range from 1.2 to 32.0% (standard deviation = 5.7). Canopy openness over
 2828 streams in old-growth forests averaged 18.0% but was highly variable and ranged from 3.4 to
 2829 34.0% (standard deviation = 5 7.9).

2830 Warren et al. (2013) compared canopy cover between old-growth-forests (>500 years old) and
 2831 young second-growth stands (~40-60 years old) in the H.J. Andrews Experimental Forest in the
 2832 Cascade Mountains of Oregon. Canopy cover was estimated using a convex spherical
 2833 densiometer. Streams were paired based on reach length, bankfull width, and north (n=2), vs.
 2834 south (n=2) facing watersheds. Results showed significant differences in percent forest cover
 2835 between old-growth and second-growth reaches in both south-facing watersheds in mid-summer
 2836 ($p < 0.10$). For the north-facing watersheds, differences in canopy cover and light availability (p
 2837 < 0.10) were only significant at 1 of the two reaches. Overall, three of the four paired old-growth
 2838 reaches had significantly lower mean percent canopy cover. The authors interpret these results as
 2839 evidence that old-growth forest canopies were more complex and had more frequent gaps.

2840 *Litter*

2841 Table 19. Treatment and responses for selected publications investigating Litter relevant to Q5.

Reference	Treatment	Response
-----------	-----------	----------

Kiffney & Richardson (2010)	Upland clearcuts with (1) no buffer, (2) 10 m buffer (~30 feet), (3) 30 m buffers	<u>8 years post-harvest data</u> Differences in litter flux relative to riparian treatment persisted through year 7, while a positive trend between buffer width and litter flux remained through year 8 . The linear relationship between reserve width and litter inputs was strongest in the first year after treatment, explaining ~57% of the variation, but the relationship could only explain ~17% of the variation in litter input by buffer width by year 8.
Bilby & Heffner (2016)	Litter samples released from canopy height at one old-growth site and one young forest site.	<u>1-year of data actual age of stands not quantified, estimated by mean height (47.0 and 32.4 m)</u> Needles released at mature sites had a higher proportion of cumulative input from greater distances than needles or alder leaves released at younger sites. The model estimated that the width of the contributing area for needles was ~35% greater at older sites than at younger sites.

2842

2843 Kiffney & Richardson (2010) compared changes in litter input between riparian harvest
2844 prescriptions that included clear-cut to stream edge, 10 m wide buffer reserve, 30 m buffer
2845 reserves, and an uncut control over the course of 8 years. No thinning was applied within the
2846 reserves. Upland treatment at all sites applied clearcut. Results showed differences in litter flux
2847 relative to riparian treatment persisted through year 7, while a positive trend between reserve
2848 width and litter flux remained through year 8. Needle inputs remained 6x higher in the buffer and
2849 control sites through year 7, and 3-6x higher in year 8 than in the clearcut sites. Twig inputs into
2850 the control and buffered sites were ~25x higher than in the clearcut sites in the first year after
2851 treatment. The linear relationship between reserve width and litter inputs was strongest in the
2852 first year after treatment, explaining ~57% of the variation, but the relationship could only
2853 explain ~17% of the variation in litter input by buffer width by year 8 (i.e., the relationship
2854 degraded over time). The authors interpret these results as evidence that litter flux from riparian
2855 plants to streams, was affected by riparian reserve width and time since logging.

2856 Bilby & Heffner (2016) used linear mixed effects models developed for young and old-growth
2857 forests of western Washington to estimate controls on litter delivery. Litter samples were released
2858 from canopy height at one old-growth forest site and one young forest site. The mean age of
2859 “mature” and “young” sites was not specified but the mean tree heights were 47.0 m and 32.4 m
2860 for the mature and young sites, respectively. Results showed that needles released at mature sites
2861 had a higher proportion of cumulative input from greater distances than needles or alder leaves
2862 released at younger sites. The model estimated that the width of the contributing area for needles
2863 was ~35% greater at older sites than at younger sites.

2864 *Source distance curves for LW*

2865 Table 20. Treatment and responses for selected publications investigating LW source distance
2866 curves relevant to Q5.

Reference	Treatment	Response
Schuett-Hames & Stewart (2019a)	30-ft no-cut buffer width, and thinning 30-75 ft from the stream (SR); retention of all shade providing trees in this area (AAS)	<u>5-years post-harvest</u> Most recruited fallen trees originated in the core zone (0-30 feet; 76%, 72%, and 64% for the reference, AAS and SR groups, respectively), while the proportion from the inner zone (30–75 feet from the stream) was ~10% greater for the SR group compared to the AAS and REF groups.
Burton et al. (2016)	Buffer widths of 6, 15, or 70 meters	<u>6 years post-harvest</u> 82-85% of the wood with discernable sources came from within 15 m of the stream , and the relative contribution of wood to streams declined rapidly with increasing distance .
Martin & Grotenfendt (2007)	Minimum buffer width of 20 m	Recruitment from within 0-20 m of stream was only 17% greater in the treated sites than in the reference sites. However, recruitment from the outer 10 – 20 m was more than double in the buffered units than in the reference units. Estimate that future supply of LW is diminished by ~10% in the treated sites compared to the reference sites.

2867

2868 Schuett-Hames & Stewart (2019a) compared differences in LW recruitment between riparian
2869 management zones harvested under the current standard Shade Rules (SR), the All-Available
2870 Shade Rule (AAS), and unharvested references for fish-bearing streams in the mixed conifer
2871 habitat type (2500 - 5000 feet elevation) for eastern Washington. Both shade rules have a 30-ft
2872 no-cut buffer (core zone) immediately adjacent to the stream. The SR prescription allows
2873 thinning in the buffer zone 30-75 feet (inner zone) from the stream while the AAS prescription
2874 requires retention of all shade providing trees in this area. Results showed that cumulative wood
2875 recruitment from tree fall after the five-year post-harvest interval was highest in the SR group,
2876 lower in the AAS group and lowest in the REF group. The SR and AAS LW recruitment rates by
2877 volume were nearly 300% and 50% higher than the REF rates, respectively. Wood recruitment in
2878 the SR sites was significantly greater than in the AAS and reference sites. Conversely,
2879 differences in wood recruitment did not differ significantly between the AAS and reference sites.
2880 Considering the source distance of post-harvest recruited LW, most recruited fallen trees
2881 originated in the core zone (76%, 72%, and 64% for the REF, AAS and SR groups, respectively),
2882 while the proportion from the inner zone (30–75 feet from the stream) was ~10% greater for the
2883 SR group compared to the AAS and REF groups. These results provide evidence that the
2884 thinning treatments applied in the inner zone of the SR treatment changed the spatial pattern
2885 (source distance) of wood recruitment from fallen trees within 5 years post-harvest.

2886 Burton et al. (2016) examined the relationship between annual in-stream wood loading and
2887 riparian buffer widths adjacent to upland thinning operations. Buffer widths were 6, 15, or 70
2888 meters and upland thinning was to 200 trees per ha (tph), with a second thinning (~10 years later)
2889 to ~85 tph, alongside an unthinned reference stand ~400 tph. Data for LW in streams were
2890 collected for 6 years (5 years after the first harvest and 1 additional year after the second
2891 harvest). The results showed that between 82-85% of the wood with discernable sources (90%

2892 for wood in early stages of decay; 45% of wood in late stages of decay) came from within 15 m
2893 of the stream, and the relative contribution of wood to streams declined rapidly with increasing
2894 distance.

2895 Martin & Grotenfendt (2007) compared riparian stand mortality and in-stream LW recruitment
2896 characteristics between riparian buffer strips with upland timber harvest and riparian stands of
2897 unharvested watersheds using aerial photography. All buffer strips in this study were a minimum
2898 of 20 m wide and included selective harvest within the 20 m zone (thinning intensity not
2899 specified or included in the analyses as an effect). The results showed significantly higher
2900 mortality (based on cumulative stand mortality: downed tree counts divided by standing tree
2901 counts + downed tree counts), significantly lower stand density (269 trees/ha in buffer units and
2902 328 trees/ha in reference units), and a significantly higher proportion of LW recruitment from the
2903 buffer zones of the treatment sites than in the reference sites. LW recruitment based on the
2904 proportion of stand recruited (PSR) was significantly higher in the buffered units compared to
2905 the reference units. However, PSR from the inner 0-20 m was only 17% greater in the buffer
2906 units than in the reference units; while PSR of the outer unit (10 – 20 m) was more than double
2907 in the buffered units than in the reference units. From their analysis they also estimate that future
2908 potential supply of LW is diminished by ~10% in the buffered sites compared to the reference
2909 sites.

2910 *LW and stand age*

2911 Jackson and Wohl (2015) compared in-stream wood loads between old-growth (> 200 years) and
2912 young forests (age not reported). This study took place within the Arapaho and Roosevelt
2913 National Forests in Colorado. In-stream wood loads (m³/ha) were recorded for reaches in 10 old-
2914 growth forests and 23 young forests. Paired t- test or Kruskal-Wallis tests were used to check for
2915 significant differences in wood load. Results indicated that channel wood load (OG = 304.4 +
2916 161.1; Y = 197.8 + 245.5 m³ /ha), floodplain wood load (OG = 109.4 + 80; Y = 47.1 + 52.8 m³
2917 /ha), and total wood load (OG = 154.7 + 64.1; Y = 87.8 + 100.6 m³ /ha) per 100 m length of
2918 stream and were significantly higher in streams of old-growth forests than in young forests.
2919 Streams in old-growth forests also had significantly more wood in jams, and more total wood
2920 jams per unit length of channel than in younger forests (jam wood volume: OG = 7.10 +/- 6.9
2921 m³; Y = 1.71 +/- 2.81 m³)

2922 *Nutrient dynamics over time*

2923 Vanderbilt et al. (2003) investigated long-term datasets (ranging from 20-30 years) from six
2924 watersheds in the H.J. Andrews Experimental Watershed (HJA) in the west-central Cascade
2925 Mountains of Oregon. Their objective was to characterize long-term patterns of N dynamics in
2926 precipitation and stream water at the HJA. Patterns between nitrogen with precipitation and
2927 discharge were analyzed with logistic regression. Results showed that dissolved organic nitrogen
2928 (DON) concentrations increased in the fall in every watershed. The increase in concentration
2929 began in July or August with the earliest rain events, and peak DON concentrations occurred in
2930 October through December before the peak in the hydrograph. DON concentrations then
2931 declined during the winter months. However, other forms of N showed inconsistent patterns

2932 across all other watersheds. The authors conclude that total annual stream discharge was a
2933 positive predictor of DON output suggesting a relationship to precipitation. Also, DON had a
2934 consistent seasonal concentration pattern. All other forms of N observed showed variability and
2935 inconsistencies with annual and seasonal stream discharge. The authors speculate that different
2936 factors may control organic vs. inorganic N export. Specifically, DIN may be strongly influenced
2937 by terrestrial or in-stream biotic controls, while DON is more strongly influenced by climate.
2938 Last, the authors suggest that DON in streams may be recalcitrant, and largely unavailable to
2939 stream organisms.

2940

2941 Focal Question 6

2942 *6. Are there feedback mechanisms (e.g., microclimate changes within the riparian buffer) related*
2943 *to forest management that affect the recovery rates of riparian functions?*

2944 The studies considered appropriate for answering this question are those that quantify how forest
2945 management practices impact one or more factors that can in-turn impact the rate of recovery of
2946 riparian function. The regeneration, growth and development of vegetation within the riparian
2947 area following treatment can impact the rate of recovery of litter inputs, shade, sediment and
2948 nutrient filtration. Reduction in shade may affect the amount of light reaching the forest
2949 understory that then could impact productivity in the riparian area. Also, disturbance of soil and
2950 removal of vegetation during riparian management operations can impact streamflow and
2951 sediment supply, which in turn impacts sediment flux into streams. The studies summarized
2952 below provide experimental evidence in how these factors (e.g., vegetation productivity,
2953 streamflow discharge, sediment disturbance) are impacted by management.

2954 However, considering the second part of this question on how these feedback mechanisms affect
2955 the recovery rates of riparian function can only be inferred. To properly answer the full question
2956 a study would require an experimental design which 1) tracks the changes in site conditions (e.g.,
2957 microclimate, light availability to groundcover, exposed soil...) after treatment relative to
2958 untreated stands, 2) evaluates how these changes in site conditions lead to changes in stand
2959 development that can then impact function (e.g., vegetation), and finally 3) how these changes in
2960 development affect the recovery rates of function. This third step would require separating out
2961 the effect of these “feedback mechanism” so that the differences in recovery rates in treated
2962 stands with and without these effects (e.g., blocking newly available light to the understory) can
2963 be compared quantitatively. No studies that specifically, and entirely address these 3 objectives
2964 collectively could be found in the literature. Thus, the following reviewed studies provide
2965 evidence of how feedback mechanisms can affect function (e.g., increased light = increased
2966 primary productivity), but how these mechanisms affect the recovery rates of any particular
2967 function (e.g., timing of recovery with and without the feedback mechanism) can only be
2968 assumed.

2969 *Litter*

2970 Yeung et al. (2019) simulated post-harvest responses to leaf-litter derived coarse particulate
2971 organic matter (CPOM) quantity in a coastal rainforest stream in British Columbia. This study

2972 used a CPOM model that was calibrated using data from multiple published studies from,
 2973 primarily the Pacific Northwest region, and several other North American regions. Calibration
 2974 data included stream flow and temperature, and CPOM following different timber harvest
 2975 intensities within 4 years of harvest. The model used estimated litterfall decreases of (-10%, -
 2976 30%, -50%, -90%) for low, moderate, high, and very high basal area removal ; peak streamflow
 2977 increases of +20%, +40%, +100%, +300%); and stream temperature increases of +1°C, +2°C,
 2978 +4°C, and +6 °C. Treatment intensities in litterfall, peak flow, and stream temperature were
 2979 modeled and analyzed individually and cumulatively to estimate their relative and combined
 2980 effects on in-stream CPOM standing stocks. Results of the model showed that, in general, the
 2981 standing stocks of CPOM decreased under the independent effects of reduced litterfall and
 2982 elevated peak flows and increased with higher stream temperatures.

2983 Along the gradient of increasing timber removal, litterfall reductions on depleting CPOM
 2984 standing stocks were at least an order of magnitude greater than those of elevated peak flows.
 2985 The magnitude of CPOM changes induced by litterfall reductions was consistently greater than
 2986 stream temperature increases, but their differences in magnitude became smaller at higher levels
 2987 of disturbance severity. Only the effects of litterfall-temperature interactions on CPOM standing
 2988 stocks were significant ($p < 0.001$). The authors interpret these results as evidence that litterfall
 2989 reduction from timber harvest was the strongest control on in-stream CPOM quantity for 4 years
 2990 post-harvest. However, the authors propose that the decreased activity of CPOM consumers
 2991 caused by increasing stream temperatures may be enough to offset the loss of litterfall inputs on
 2992 standing CPOM stocks. The caveat of this study is that it did not include LW dynamics in
 2993 preserving CPOM post-harvest. There is evidence that in-stream LW can act as a catchment for
 2994 CPOM (May & Gresswell, 2003; Richardson et al. 2007).

2995 *Sediment*

2996 Table 21. Treatment and responses for selected publications investigating Sediment relevant to
 2997 Q5.

Reference	Treatment	Response
Safeeq et al. (2020)	Long-term dataset with mixture of management, storm events, and	estimate that following harvest, changes on streamflow alone was estimated in being responsible for < 10% of the resulting suspended sediment transported into streams, while the increase in sediment supply due to harvest disturbance was responsible for >90% .
Litschert & MacDonald (2009)	Post-harvest stream sediment delivery pathway development frequency.	<u>1-year post-harvest data (n = 200 harvest units)</u> The authors conclude that in general, USFS riparian forest harvest practices are effective in reducing the development of sediment delivery pathways. They also interpret these results as evidence that skid trails should be directed away from streams , maintain surface roughness, and promptly decommissioned.

2998

2999 Safeeq et al. (2020) analyzed a long-term data set to changes in streamflow, and suspended
 3000 sediment load and sediment bedload in streams between two watersheds; one with a history of

3001 timber management and one with no history of timber management. The two watersheds were
3002 located in the H.J. Andrews Experimental Forest and were paired by size, aspect, and
3003 topography. The treatment watershed was 100% clearcut during the period from 1962-1966,
3004 broadcast burned in 1966 and re-seeded in 1968. Streamflow and sediment data were taken
3005 intermittently; suspended sediment data after large storm events between 1952 (pre-harvest) and
3006 1988; and sediment bedload in 2016. The researchers used a reverse regression technique to
3007 evaluate the relative and absolute importance of changes in streamflow versus changes in
3008 sediment supply from timber harvest on sediment transport. There were no significant changes in
3009 precipitation patterns before or after harvest. The results for post-treatment sediment yields
3010 showed suspended load declined to pre-treatment levels in the first two decades following
3011 treatment and bedload remained elevated, causing the bedload proportion of the total load to
3012 increase through time. Changes in streamflow alone account for 477 Mg/km² (10%) of the
3013 suspended load and 113 Mg/km² (5%) of the bedload over the post-treatment period. Increase in
3014 suspended sediment yield due to increase in sediment supply from timber harvest activities was
3015 84% of the measured post-treatment total suspended sediment yield. The authors estimate that
3016 following harvest, changes on streamflow alone was estimated in being responsible for < 10% of
3017 the resulting suspended sediment transported into streams, while the increase in sediment supply
3018 due to harvest disturbance was responsible for >90%. Thus, while timber harvest-induced
3019 increases in streamflow does increase sediment transport, it is negligible compared to the
3020 increase in sediment source created from management practices.

3021 Litschert & MacDonald (2009) investigated the frequency of sediment delivery pathways in
3022 riparian management areas and their physical characteristics and connectivity following harvest.
3023 In this study the authors describe sediment delivery pathways (“features”) as rills, gullies, and
3024 sediment plumes that form when excess sediment relative to overland flows transports sediment
3025 from the hillslope to the stream. The authors surveyed 200 riparian management areas (RMA) in
3026 four different National Forests of the Sierra Nevada and Cascade Mountains of California. USFS
3027 policy requires 90-m wide RMA along each side of perennial streams and 45-m wide RMA along
3028 each side of all ephemeral and intermittent streams. When features were found within an RMA,
3029 data for years since harvest, soil depth, soil erodibility (K), feature length, feature gradient,
3030 aspect, elevation, hillslope gradient, hillslope curvature, surface roughness, and connectivity
3031 were recorded for analysis. Association between these variables were analyzed with a
3032 Spearman’s rank correlation. The variables most strongly associated with feature length were
3033 used to develop a multiple linear regression model to predict feature length. Only 19 of the 200
3034 harvest units had sediment development pathways. Feature pathways ranged in age (time since
3035 harvest) from 2 to 18 years, and in length from 10 m to 220 m. Of the 19 feature pathways, only
3036 six were connected to streams, and five of those originated from skid trails. Feature pathway
3037 length was significantly related to mean annual precipitation, cosine of the aspect, elevation, and
3038 hillslope gradient ($R^2 = 64\%$, $p = 0.004$). The authors conclude that in general, USFS riparian
3039 forest harvest practices are effective in reducing the development of sediment delivery pathways.
3040 They also interpret these results as evidence that skid trails should be directed away from
3041 streams, maintain surface roughness, and promptly decommissioned.
3042

3043 *Impacts on Microclimate*

3044 Anderson et al. (2007) compared changes in understory microclimate above the stream, within
3045 the channel, and within the riparian area between thinned and unthinned riparian stands. The
3046 focus of this study was on second-growth (30- to 80-year-old) riparian Douglas-fir forests along
3047 headwater streams in the western Oregon Coast and Cascade Range. Stands were either thinned
3048 to approximately 198 trees per acre (TPA) or were left unthinned and ranged from 500-865 TPA.
3049 Streams within treated stands were surrounded by buffers of either 1) one site-potential tree
3050 averaging 69 m (B1, B1-T thinned and unthinned respectively), 2) variable width buffer
3051 averaging 22 m (VB, and VB-T), or 3) streamside retention buffer averaging 9 m (SR, and SR-
3052 T). Further, directly adjacent randomly selected B1-T and VB-T buffers patch openings (0.4 ha)
3053 were created (B1-P, VB-P). Microsite and microclimate responses were repeat sampled for each
3054 treatment and compared with untreated stands (UT). Within the riparian buffer zones, daily
3055 maximum temperatures were higher in all treated stands when compared to UT stands. The
3056 differences in daily maximum temperatures between treated and untreated stands ranged from
3057 1.1°C (B1) to 4.0°C (SR-T), but the difference was only significant in one SR-T stand. Daily
3058 maximum air temperature within buffer zones adjacent to patch openings were 3.5°C higher than
3059 in UT stands. Within patch openings daily maximum temperatures were on average 6 to 9°C
3060 higher than in UT stands. Soil temperature changes were only evident within patch openings
3061 ranging from 3.6 - 8.8°C higher than in UT stands. VB-T buffers that were 15 m wide or wider
3062 exhibited changes in daily maximum air temperature above stream centers <1°C and daily
3063 minimum relative humidity <5% lower than in untreated stands. The authors conclude that in
3064 general, thinned stands are warmer and drier than unthinned stands. However, the results for
3065 differences in microclimate were only significant in narrow (9 m) thinned buffers and patch
3066 openings.

3067 Anderson & Meleason (2009) conducted a companion study to Anderson et al. (2007) and
3068 compared changes in small (5-29 cm diameter) and large (≥ 30 cm diameter) downed wood
3069 abundance and understory vegetation between treated and untreated stands 5 years after harvest.
3070 Treatments compared were the same as those described in Anderson et al. (2007) discussed
3071 above. The results for small and large downed wood were highly variable between pre- and post-
3072 harvest periods and between treatments but the authors speculate from trends in the data that
3073 both wood and vegetation responses within buffers ≥ 15 m wide were insensitive to treatments.
3074 The strongest contrast in rate of change in herb cover was between the SR-T and VB-T buffers
3075 with higher herbaceous cover in the SR-T buffers and highest in SR-T buffers adjacent to patch
3076 openings. The authors conclude that in general these thinning treatments only led to subtle
3077 changes in understory vegetation cover and composition. Because of the high variability in
3078 responses among and between treatments significance could not be confirmed. The authors
3079 further conclude that a better functional understanding of the changes in ecological processes
3080 associated with changes in habitat characteristics following changes in understory wood and
3081 vegetation cover is needed to help discern ecological significance.

3082

3083 Focal Question 7

3084 *7. What major data gaps and uncertainties exist relative to effects of timber harvest (both*
3085 *riparian and adjacent upland) on the riparian functions?*

3086 No studies that provide experimental evidence that quantifies how specific treatments within the
3087 riparian area affect bank stability were found based on our search criteria (published after 2000,
3088 conducted in western North America). However, this may be because bank erosion relates
3089 directly to sediment transport and thus bank stability is inferred by the magnitude of change in
3090 sediment export. Furthermore, the importance of vegetation retention and equipment exclusion in
3091 areas closest to the stream for maintaining bank stability appears to be well understood
3092 considering its prevalence in riparian forest management plans ([WAC 222-30-022](#); WAC 22-30-
3093 021; 2022 [ODE](#); IDAPA 20.02.01).

3094 Our search of the literature focused on how treatments within or adjacent to forested riparian
3095 areas impact one or more of the riparian functions. Most of the studies found in our search focus
3096 on the impacts of riparian treatment on LW and shade (commonly coupled with stream
3097 temperature). There is also a significant body of research that considers the impact of harvest on
3098 nutrient and sediment flux into streams. Fewer studies could be found that quantify changes in
3099 litter input following riparian management.

3100 While few studies could be found that provide direct experimental evidence of how bank
3101 stability is affected by timber harvest, two studies were found that compared the relative
3102 influence of different factors on bank stability. Both of which showed evidence that bank
3103 stability is influenced by the type of vegetation dominating the riparian area. Rood et al. (2015)
3104 compared the relative erosion resistance of riverbanks occupied by forests versus grassland along
3105 the Elk River in British Columbia, Canada. This study used a combination of field sampling and
3106 aerial photo analysis from 1995 to 2013 to estimate the differences in channel migration between
3107 forest and grass dominated riparian areas. Relative tree cover was binned into 5 categories
3108 ranging from (1) no trees to (5) completely treed. Relative channel change was binned into 2
3109 categories as ‘moderate change’ for channels that migrated between 45 and 75 m, and as ‘major
3110 change’ for channels that migrated more than 75 m. Chi square analysis was used to assess the
3111 distributions of vegetation of channels with moderate and major changes. Results of the chi
3112 square analysis showed that the distribution of the observed vegetation types differed
3113 significantly ($p < 0.05$) by channel change categories. Of the 15 sites assessed with moderate or
3114 major erosion (changes), 7 were along banks dominated by grasslands without trees (‘1’), four
3115 were assessed as a ‘2’, with some trees, and three were in a ‘3’ with a mixed zone of similar
3116 proportions of trees and clearing. Only one site with a ‘4’ showed a moderate amount of change.
3117 The authors interpret these results as evidence that trees are better than grass at stabilizing banks,
3118 and that stability increases with tree cover.

3119 Outside of the U.S., Krzeminska et al. (2019), investigated the effect of different types of
3120 riparian vegetation on stream bank stability in a small agricultural catchment in South-Eastern
3121 Norway. The dominating soil type within the catchment is coarse moraine in the forested areas
3122 and marine deposits with silt loam and silty clay loam texture in agriculture areas. The
3123 researchers used a combination of field collected data with stream bank stability modeling using

3124 Bank-Stability and Toe-Erosion Modeling (BSTEM). Three experimental plots were established,
3125 one for each dominant vegetation type, grass dominated, shrub dominated, and tree dominated.
3126 Investigations of in-situ undrained shear strength of the root-reinforced soil were done with a
3127 Field Inspection Vane Tester. Additionally, potential changes in the bank profile were monitored
3128 with a series of erosion pins, 6 pins per each plot. Changes in root cohesion and % cover over
3129 time for each vegetation type were estimated using the RipRoots sub-model in BSTEM. Their
3130 results showed a difference in bank stability based on vegetation type, that varied seasonally with
3131 groundwater level and stream water level. The grass dominated and tree dominated plots,
3132 specifically, showed the lowest estimated stability during spring (March to April) and early
3133 autumn (September to November), and the highest estimated stability during the summer months
3134 (May-June). This seasonal trend was also observed for the shrub plots but not as strongly.
3135 Steeper slopes in the grass and shrub dominated plots showed a trend of reduced stability for
3136 plots 54° slopes showing potential for failure. The tree dominated plots showed a trend of lower
3137 stability for steeper slopes, however, it wasn't as strong of a trend and the model did not predict
3138 potential for failure or 'instability'. Regardless of season, groundwater levels, or slope steepness
3139 the tree plots showed the highest estimated bank stability overall.

3140 These two studies that investigate bank stability use methods which could be applied to an
3141 experimental design that also considers differences in stability between treated (harvested) and
3142 untreated stands. The combination of field observation and simulation modeling used by
3143 Krzeminska et al. (2019), especially, could be used to estimate how timber harvest affects bank
3144 stability (or erosion) while also accounting for geomorphic and hydrological differences.

3145 Considering the topics included in the focal questions, studies that investigate the effects of
3146 clearcut gaps, and studies that quantify how treatment within the riparian zone relates to
3147 resilience to fire had the fewest studies providing experimental evidence. Other than the Hard
3148 Rock and Soft Rock studies, only 2 other studies (Janisch et al., 2012, Swartz et al., 2020) were
3149 found that investigate the effects of similar buffer treatment designs (patched buffers and riparian
3150 canopy gaps). For how treatments within the riparian zone relate to resilience to fire, there were
3151 no studies that provide experimental evidence on this topic based on the search criteria. Some
3152 studies were found to quantify the probability of fire or fire severity within riparian zones in
3153 general (Reeves et al. 2006; Van de Water & North, 2011). However, none compares the
3154 resilience of riparian stands between treated and untreated stands after fire. One study, Ceder et
3155 al. (2018) used simulation modeling to compare fire susceptibility between managed and
3156 unmanaged stands and has been included in focal question 4.

3157 Indeed, Stone et al. (2010) surveyed fire management officers from 55 national forests across 11
3158 western states and found that fewer than half (43%) of them indicated that they were conducting
3159 fuel reduction treatments in riparian areas. The primary objective for most of these treatments
3160 involved some form of fuel reduction (83%), while others focused on multiple objectives such as
3161 ecological restoration and habitat improvement. Most of these treatments (93%) were of small
3162 extent (< 300 acres) and occurred in the wildland urban interface (73%). The authors conclude
3163 that these results are promising, but that well-designed monitoring programs are needed to
3164 estimate the consequences of these treatments on fire risk and other ecological effects.

3165 The study from Prichard et al. (2020), discussed in question 3, used a combination of
3166 simultaneous autoregression (SAR) and random forest (RF) modeling approaches to model the
3167 drivers of fire severity and the effectiveness of fuel treatments in mitigating fire severity in the
3168 2014 Carlton Complex. Results from this study provided evidence on how vegetation (based on
3169 broad LANDFIRE classifications), topography, and different fuel treatments (e.g., thinning only,
3170 thin and pile burn, thin and broadcast burn, etc.) related to fire severity and fire spread. This
3171 approach has potential to be used in riparian areas burned by wildfires. In terms of the topic of
3172 how various treatments relate to riparian forest resistance and resilience to fire would require
3173 using a dataset of riparian forest stand characteristics that includes information on fuel
3174 treatments, time since last fire, and basin characteristics. This information could be used along
3175 with spatial information of burn severity immediately following a fire.

3176 References

- 3177 Anderson, P. D., & Meleason, M. A. (2009). Discerning responses of down wood and understory
3178 vegetation abundance to riparian buffer width and thinning treatments: an equivalence–inequivalence
3179 approach. *Canadian Journal of Forest Research*, 39(12), 2470-2485.
- 3180 Anderson, P. D., Larson, D. J., & Chan, S. S. (2007). Riparian buffer and density management influences on
3181 microclimate of young headwater forests of western Oregon. *Forest Science*, 53(2), 254-269.
- 3182 Arkle, R. S., & Pilliod, D. S. (2010). Prescribed fires as ecological surrogates for wildfires: a stream and
3183 riparian perspective. *Forest Ecology and Management*, 259(5), 893-903.
- 3184 Bahuguna, D., Mitchell, S. J., & Miquelajauregui, Y. (2010). Windthrow and recruitment of large woody
3185 debris in riparian stands. *Forest Ecology and Management*, 259(10), 2048-2055.
- 3186 Benda, L. E., Litschert, S. E., Reeves, G., & Pabst, R. (2016). Thinning and in-stream wood recruitment in
3187 riparian second growth forests in coastal Oregon and the use of buffers and tree tipping as
3188 mitigation. *Journal of forestry research*, 27(4), 821-836.
- 3189 Benda, L., Miller, D. A. N. I. E. L., Sias, J. O. A. N., Martin, D. O. U. G. L. A. S., Bilby, R., Veldhuisen, C., &
3190 Dunne, T. (2003, January). Wood recruitment processes and wood budgeting. In *American Fisheries*
3191 *Society Symposium* (pp. 49-74). American Fisheries Society.
- 3192 Beschta, R. L., Bilby, R.E.Brown, G.W., Holtby, L.B., and Hofstra., T.D., 1987. Stream Temperature and
3193 Aquatic Habitat: Fisheries and Forestry Interactions. In: *Streamside Management: Forestry and Fisheries*
3194 *Interactions*. E. O. Salo and T. W. Cundy (Editors). Contribution No. 57, University of Washington, Institute
3195 of Forest Resources, 471 pp.
- 3196 Bilby, R. E., & Heffner, J. T. (2016). Factors influencing litter delivery to streams. *Forest Ecology and*
3197 *Management*, 369, 29–37. <https://doi.org/10.1016/j.foreco.2016.03.031>
- 3198 Bilby, R. E., Sullivan, K., & Duncan, S. H. (1989). The generation and fate of road-surface sediment in
3199 forested watersheds in southwestern Washington. *Forest Science*, 35(2), 453-468.
- 3200 Bjornn, T. C., & Reiser, D. W. (1991). Habitat requirements of salmonids in streams. *American Fisheries*
3201 *Society Special Publication*, 19(837), 138.

- 3202 Bladon, K. D., Cook, N. A., Light, J. T., & Segura, C. (2016). A catchment-scale assessment of stream
 3203 temperature response to contemporary forest harvesting in the Oregon Coast Range. *Forest Ecology and*
 3204 *Management*, 379, 153-164.
- 3205 Bladon, K. D., Segura, C., Cook, N. A., Bywater-Reyes, S., & Reiter, M. (2018). A multicatchment analysis of
 3206 headwater and downstream temperature effects from contemporary forest harvesting. *Hydrological*
 3207 *Processes*, 32(2), 293-304.
- 3208 Brown, G. W. (1983). *Forestry and water quality. Forestry and water quality., (Ed. 2).*
- 3209 Brown, G. W., Gahler, A. R., & Marston, R. B. (1973). Nutrient losses after clear-cut logging and slash
 3210 burning in the Oregon Coast Range. *Water Resources Research*, 9(5), 1450-1453.
- 3211 Burton, J. I., Olson, D. H., & Puettmann, K. J. (2016). Effects of riparian buffer width on wood loading in
 3212 headwater streams after repeated forest thinning. *Forest Ecology and Management*, 372, 247–257.
 3213 <https://doi.org/10.1016/j.foreco.2016.03.053>
- 3214 Bywater-Reyes, S., Bladon, K. D., & Segura, C. (2018). Relative influence of landscape variables and
 3215 discharge on suspended sediment yields in temperate mountain catchments. *Water Resources*
 3216 *Research*, 54(7), 5126-5142.
- 3217 Bywater-Reyes, S., Segura, C., & Bladon, K. D. (2017). Geology and geomorphology control suspended
 3218 sediment yield and modulate increases following timber harvest in temperate headwater
 3219 streams. *Journal of Hydrology*, 548, 754-769.
- 3220 Camp, A., C. Oliver, P. Hessburg, and R. Everett. (1997). Predicting late-successional fire refugia pre-dating
 3221 European settlement in the Wenatchee Mountains. *Forest Ecology and Management* 95 63-77
- 3222 Chan, S., P. Anderson, J. Cissel, L. Lateen. and C. Thompson. 2004. Variable density management in
 3223 Riparian Reserves: lessons learned from an operational study in managed forests of western Oregon,
 3224 USA. *For. Snow Landsc. Res.* 78,1/2:151-172.
- 3225 Chapman, D. W., & Bjornn, T. C. (1969). Distribution of salmonids in streams. In *Symp. Salmon Trout*
 3226 *Streams*. Institute of Fisheries, University of British Columbia, Vancouver (pp. 153-176).
- 3227 Chen, X., Wei, X., & Scherer, R. (2005). Influence of wildfire and harvest on biomass, carbon pool, and
 3228 decomposition of large woody debris in forested streams of southern interior British Columbia. *Forest*
 3229 *Ecology and Management*, 208(1-3), 101-114.
- 3230 Chen, X., Wei, X., Scherer, R., Luider, C., & Darlington, W. (2006). A watershed scale assessment of in-
 3231 stream large woody debris patterns in the southern interior of British Columbia. *Forest Ecology and*
 3232 *Management*, 229(1-3), 50-62.
- 3233 Chesney, C. (2000). Functions of wood in small, steep streams in eastern Washington: Summary of
 3234 results for project activity in the Ahtanum, Cowiche, and Tieton basins. *Timber, Fish, Wildlife*.
- 3235 CMER 03-308 (2004) Review of the Available Literature Related to Wood Loading Dynamics in and
 3236 around Stream in eastern Washington Forests
- 3237 Cole, E., & Newton, M. (2013). Influence of streamside buffers on stream temperature response
 3238 following clear-cut harvesting in western Oregon. *Canadian journal of forest research*, 43(11), 993-1005.

- 3239 Cooper, J. R., Gilliam, J. W., Daniels, R. B., & Robarge, W. P. (1987). Riparian areas as filters for agricultural
3240 sediment. *Soil science society of America journal*, 51(2), 416-420.
- 3241 Crandall, T., Jones, E., Greenhalgh, M., Frei, R. J., Griffin, N., Severe, E., ... & Abbott, B. W. (2021).
3242 Megafire affects stream sediment flux and dissolved organic matter reactivity, but land use dominates
3243 nutrient dynamics in semiarid watersheds. *PloS one*, 16(9), e0257733.
- 3244 Cupp, C.E. and T.J. Lofgren. 2014. Effectiveness of riparian management zone prescriptions in protecting
3245 and maintaining shade and water temperature in forested streams of Eastern Washington. Cooperative
3246 Monitoring Evaluation and Research Report CMER 02-212. Washington State Forest Practices Adaptive
3247 Management Program. Washington Department of Natural Resources, Olympia, WA.
- 3248 Deval, C., Brooks, E. S., Gravelle, J. A., Link, T. E., Dobre, M., & Elliot, W. J. (2021). Long-term response in
3249 nutrient load from commercial forest management operations in a mountainous watershed. *Forest
3250 Ecology and Management*, 494, 119312.
- 3251 Devotta, D. A., Fraterrigo, J. M., Walsh, P. B., Lowe, S., Sewell, D. K., Schindler, D. E., & Hu, F. S. (2021).
3252 Watershed Alnus cover alters N: P stoichiometry and intensifies P limitation in subarctic
3253 streams. *Biogeochemistry*, 153(2), 155-176.
- 3254 Dwire, K. A., & Kauffman, J. B. (2003). Fire and riparian ecosystems in landscapes of the western USA.
3255 *Forest Ecology and Management*, 178(1-2), 61-74.
- 3256 Ebersole, J. L., Liss, W. J., & Frissell, C. A. (2001). Relationship between stream temperature, thermal
3257 refugia and rainbow trout *Oncorhynchus mykiss* abundance in arid-land streams in the northwestern
3258 United States. *Ecology of freshwater fish*, 10(1), 1-10.
- 3259 Ehinger, W.J., W.D. Bretherton, S.M. Estrella, G. Stewart, D.E. Schuett-Hames, and S.A. Nelson. 2021.
3260 Effectiveness of Forest Practices Buffer Prescriptions on Perennial Non-fish-bearing Streams on Marine
3261 Sedimentary Lithologies in Western Washington. Cooperative Monitoring, Evaluation, and Research
3262 Committee Report CMER 2021.08.24, Washington State Forest Practices Adaptive Management
3263 Program, Washington Department of Natural Resources, Olympia, WA.
- 3264 Everett, R., Schellhaas, R., Ohlson, P., Spurbeck, D., & Keenum, D. (2003). Continuity in fire disturbance
3265 between riparian and adjacent sideslope Douglas-fir forests. *Forest Ecology and Management*, 175(1-3),
3266 31-47.
- 3267 Fox, M. J. (2001). A new look at the quantities and volumes of instream wood in forested basins within
3268 Washington State (Doctoral dissertation, University of Washington).
- 3269 Fox, M., & Bolton, S. (2007). A regional and geomorphic reference for quantities and volumes of instream
3270 wood in unmanaged forested basins of Washington State. *North American Journal of Fisheries
3271 Management*, 27(1), 342-359.
- 3272 Fox, M., Bolton, S., & Conquest, L. (2003). 14Reference conditions for instream wood in western
3273 Washington. University of Washington Press: Seattle, WA, 361-393.
- 3274 Fratkin, M. M., Segura, C., & Bywater-Reyes, S. (2020). The influence of lithology on channel geometry
3275 and bed sediment organization in mountainous hillslope-coupled streams. *Earth Surface Processes and
3276 Landforms*, 45(10), 2365-2379.

- 3277 Fredriksen, R. L. (1975). Nitrogen, phosphorus and particulate matter budgets of five coniferous forest
3278 ecosystems in the western Cascades Range, Oregon.
- 3279 Gomi, T., Dan Moore, R., & Hassan, M. A. (2005). Suspended sediment dynamics in small forest streams
3280 of the Pacific Northwest 1. *JAWRA Journal of the American Water Resources Association*, 41(4), 877-898.
- 3281 Gomi, T., Sidle, R. C., Bryant, M. D., & Woodsmith, R. D. (2001). The characteristics of woody debris and
3282 sediment distribution in headwater streams, southeastern Alaska. *Canadian Journal of Forest
3283 Research*, 31(8), 1386-1399.
- 3284 Gravelle, J. A., & Link, T. E. (2007). Influence of timber harvesting on headwater peak stream
3285 temperatures in a northern Idaho watershed. *Forest Science*, 53(2), 189-205.
- 3286 Gravelle, J. A., Ice, G., Link, T. E., & Cook, D. L. (2009). Nutrient concentration dynamics in an inland
3287 Pacific Northwest watershed before and after timber harvest. *Forest Ecology and Management*, 257(8),
3288 1663-1675.
- 3289 Gregory, S. (1990). *Riparian management guide: Willamette National Forest*. US Department of
3290 Agriculture, Forest Service, Pacific Northwest Region.
- 3291 Gresswell, R. E., Barton, B. A., & Kershner, J. L. (Eds.). (1989). *Practical approaches to riparian resource
3292 management: an educational workshop: May 8-11, 1989, Billings, Montana*. US Bureau of Land
3293 Management.
- 3294 Groom, J. D., Dent, L., Madsen, L. J., & Fleuret, J. (2011b). Response of western Oregon (USA) stream
3295 temperatures to contemporary forest management. *Forest Ecology and Management*, 262(8), 1618-
3296 1629.
- 3297 Groom, J. D., L. Dent, and L. J. Madsen. (2011a). Stream temperature change detection for state and
3298 private forests in the Oregon Coast Range, *Water Resour. Res.* 47, W01501
- 3299 Guenther, S. M., Gomi, T., & Moore, R. D. (2014). Stream and bed temperature variability in a coastal
3300 headwater catchment: influences of surface-subsurface interactions and partial-retention forest
3301 harvesting. *Hydrological Processes*, 28(3), 1238-1249.
- 3302 Harmon, M. E., Franklin, J. F., Swanson, F. J., Sollins, P., Gregory, S. V., Lattin, J. D., ... & Cummins, K. W.
3303 (1986). Ecology of coarse woody debris in temperate ecosystems. *Advances in ecological research*, 15,
3304 133-302.
- 3305 Hart, Stephanie K., David E. Hibbs, and Steven S. Perakis. (2013) "Riparian litter inputs to streams in the
3306 central Oregon Coast Range." *Freshwater Science* 32.1 (2013): 343-358.
- 3307 Hartman, G. F., & Scrivener, J. C. (1990). Impacts of forestry practices on a coastal stream ecosystem,
3308 Carnation Creek, British Columbia.
- 3309 Hassan, M. A., Church, M., Lisle, T. E., Brardinoni, F., Benda, L., & Grant, G. E. (2005). Sediment transport
3310 and channel morphology of small, forested streams 1. *Jawra journal of the american water resources
3311 association*, 41(4), 853-876.

3312 Hatten, J. A., Segura, C., Bladon, K. D., Hale, V. C., Ice, G. G., & Stednick, J. D. (2018). Effects of
3313 contemporary forest harvesting on suspended sediment in the Oregon Coast Range: Alsea Watershed
3314 Study Revisited. *Forest Ecology and Management*, 408, 238-248.

3315 Hoffmann, C. C., Kjaergaard, C., Uusi-Kämppe, J., Hansen, H. C. B., & Kronvang, B. (2009). Phosphorus
3316 Retention in Riparian Buffers: Review of Their Efficiency. *Journal of Environmental Quality*, 38(5), 1942–
3317 1955. <https://doi.org/10.2134/jeq2008.0087>

3318 Hough-Snee, N., Kasprak, A., Rossi, R. K., Bouwes, N., Roper, B. B., & Wheaton, J. M. (2016).
3319 Hydrogeomorphic and Biotic Drivers of Instream Wood Differ Across Sub-basins of the Columbia River
3320 Basin, USA. *River Research and Applications*, 32(6), 1302–1315. <https://doi.org/10.1002/rra.2968>

3321 Hunter, M. A., & Quinn, T. (2009). Summer water temperatures in alluvial and bedrock channels of the
3322 Olympic Peninsula. *Western Journal of Applied Forestry*, 24(2), 103-108.

3323 Hyatt, T. L., & Naiman, R. J. (2001). The residence time of large woody debris in the Queets River,
3324 Washington, USA. *Ecological Applications*, 11(1), 191-202.

3325 Jackson, C. R., C. A. Sturm, and J. M. Ward. 2001. Timber harvest impacts on small headwater stream
3326 channels in the coast ranges of Washington. *JAWRA Journal of the American Water Resources
3327 Association* 37(6):1533-1549.

3328 Jackson, K. J., & Wohl, E. (2015). Instream wood loads in montane forest streams of the Colorado Front
3329 Range, USA. *Geomorphology*, 234, 161-170.

3330 Janisch, J. E., Wondzell, S. M., & Ehinger, W. J. (2012). Headwater stream temperature: Interpreting
3331 response after logging, with and without riparian buffers, Washington, USA. *Forest Ecology and
3332 Management*, 270, 302-313.

3333 Johnson, S. L., & Jones, J. A. (2000). Stream temperature responses to forest harvest and debris flows in
3334 western Cascades, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences*, 57(S2), 30-39.

3335 Karwan, D. L., Gravelle, J. A., & Hubbart, J. A. (2007). Effects of timber harvest on suspended sediment
3336 loads in Mica Creek, Idaho. *Forest Science*, 53(2), 181-188.

3337 Kaylor, M. J., Warren, D. R., & Kiffney, P. M. (2017). Long-term effects of riparian forest harvest on light in
3338 Pacific Northwest (USA) streams. *Freshwater Science*, 36(1), 1-13.

3339 Kibler, K. M., Skaugset, A., Ganio, L. M., & Huso, M. M. (2013). Effect of contemporary forest harvesting
3340 practices on headwater stream temperatures: Initial response of the Hinkle Creek catchment, Pacific
3341 Northwest, USA. *Forest ecology and management*, 310, 680-691.

3342 Kiffney, P., and J. Richardson. 2010. Organic matter inputs into headwater streams of southwestern
3343 British Columbia as a function of riparian reserves and time since harvesting. *Forest Ecology and
3344 Management*. 260:1931-1942.

3345 Knight, S.M. (1990). Forest harvesting impacts on coarse woody debris and channel form in central
3346 Oregon streams. M.S. Thesis. Oregon State University, Corvallis, Oregon.

3347 Lewis, D. D. (1998, April). Naive (Bayes) at forty: The independence assumption in information retrieval.
3348 In *European conference on machine learning* (pp. 4-15). Berlin, Heidelberg: Springer Berlin Heidelberg.

3349 Liquori, M. K. (2006). POST-HARVEST RIPARIAN BUFFER RESPONSE: IMPLICATIONS FOR WOOD
3350 RECRUITMENT MODELING AND BUFFER DESIGN 1. JAWRA Journal of the American Water Resources
3351 Association, 42(1), 177-189.

3352 Litschert, S. E., & MacDonald, L. H. (2009). Frequency and characteristics of sediment delivery pathways
3353 from forest harvest units to streams. *Forest Ecology and Management*, 259(2), 143-150.

3354 Macdonald, J. S., Beaudry, P. G., MacIsaac, E. A., & Herunter, H. E. (2003). The effects of forest harvesting
3355 and best management practices on streamflow and suspended sediment concentrations during
3356 snowmelt in headwater streams in sub-boreal forests of British Columbia, Canada. *Canadian Journal of
3357 Forest Research*, 33(8), 1397-1407.

3358 Macdonald, J. S., MacIsaac, E. A., & Herunter, H. E. (2003b). The effect of variable-retention riparian
3359 buffer zones on water temperatures in small headwater streams in sub-boreal forest ecosystems of
3360 British Columbia. *Canadian journal of forest research*, 33(8), 1371-1382.

3361 Martin, D. J., & Grotefendt, R. A. (2007). Stand mortality in buffer strips and the supply of woody debris
3362 to streams in Southeast Alaska. *Canadian Journal of Forest Research*, 37(1), 36-49.

3363 Martin, D. J., Kroll, A. J., & Knoth, J. L. (2021). An evidence-based review of the effectiveness of riparian
3364 buffers to maintain stream temperature and stream-associated amphibian populations in the Pacific
3365 Northwest of Canada and the United States. *Forest Ecology and Management*, 491, 119190.

3366 May, C. L., & Gresswell, R. E. (2003). Large wood recruitment and redistribution in headwater streams in
3367 the southern Oregon Coast Range, USA. *Canadian Journal of Forest Research*, 33(8), 1352-1362.

3368 McDade, M, F. Swanson, W. McKee, J. Franklin and J. Van Sickle. (1990). Source distances for coarse
3369 woody debris entering small streams in western Oregon and Washington. *Canadian Journal of Forest
3370 Resources* 20, 326-330

3371 McIntyre, A.P., M.P. Hayes, W.J. Ehinger, S.M. Estrella, D. Schuett-Hames, and T. Quinn (technical
3372 coordinators). 2018. Effectiveness of Experimental Riparian Buffers on Perennial Non-fish-bearing
3373 Streams on Competent Lithologies in Western Washington. Cooperative Monitoring, Evaluation and
3374 Research Report CMER 18-100, Washington State Forest Practices Adaptive Management Program,
3375 Washington Department of Natural Resources, Olympia, WA.

3376 McIntyre, A.P., M.P. Hayes, W.J. Ehinger, S.M. Estrella, D.E. Schuett-Hames, R. Ojala-Barbour, G. Stewart
3377 and T. Quinn (technical coordinators). 2021. Effectiveness of experimental riparian buffers on perennial
3378 non-fish-bearing streams on competent lithologies in western Washington – Phase 2 (9 years after
3379 harvest). Cooperative Monitoring, Evaluation and Research Report CMER 2021.07.27, Washington State
3380 Forest Practices Adaptive Management Program, Washington Department of Natural Resources,
3381 Olympia, WA.

3382 Meleason, M. A., Gregory, S. V., & Bolte, J. P. (2003). Implications of riparian management strategies on
3383 wood in streams of the Pacific Northwest. *Ecological Applications*, 13(5), 1212-1221.

3384 Merschel, A. G., Spies, T. A., & Heyerdahl, E. K. (2014). Mixed-conifer forests of central Oregon: effects of
3385 logging and fire exclusion vary with environment. *Ecological Applications*, 24(7), 1670-1688.

3386 Mueller, E. R., & Pitlick, J. (2013). Sediment supply and channel morphology in mountain river systems: 1.
3387 Relative importance of lithology, topography, and climate. *Journal of Geophysical Research: Earth*
3388 *Surface*, 118(4), 2325-2342.

3389 Murray, G. L. D., Edmonds, R. L., & Marra, J. L. (2000). Influence of partial harvesting on stream
3390 temperatures, chemistry, and turbidity in forests on the western Olympic Peninsula,
3391 Washington. *Northwest science.*, 74(2), 151-164.

3392 Naiman, R. J., Fetherston, K. L., McKay, S. J., & Chen, J. (1998). Riparian forests. *River ecology and*
3393 *management: lessons from the Pacific Coastal Ecoregion*, 289-323.

3394 Nowakowski, A. L., & Wohl, E. (2008). Influences on wood load in mountain streams of the Bighorn
3395 National Forest, Wyoming, USA. *Environmental Management*, 42(4), 557-571.

3396 Polyakov, V., Fares, A., & Ryder, M. H. (2005). Precision riparian buffers for the control of nonpoint source
3397 pollutant loading into surface water: A review. *Environmental Reviews*, 13(3), 129-144.

3398 Puntenney-Desmond, K. C., Bladon, K. D., & Silins, U. (2020). Runoff and sediment production from
3399 harvested hillslopes and the riparian area during high intensity rainfall events. *Journal of Hydrology*, 582,
3400 124452.

3401 Quinn, T., G.F. Wilhere, and K.L. Krueger, technical editors. (2020). *Riparian Ecosystems, Volume 1:*
3402 *Science Synthesis and Management Implications. Habitat Program, Washington Department of Fish and*
3403 *Wildlife, Olympia.*

3404 Rachels, A. A., Bladon, K. D., Bywater-Reyes, S., & Hatten, J. A. (2020). Quantifying effects of forest
3405 harvesting on sources of suspended sediment to an Oregon Coast Range headwater stream. *Forest*
3406 *Ecology and Management*, 466, 118123.

3407 Reid, D. A., & Hassan, M. A. (2020). Response of in-stream wood to riparian timber harvesting: Field
3408 observations and long-term projections. *Water Resources Research*, 56(8), e2020WR027077.

3409 Reiter, M., Bilby, R. E., Beech, S., & Heffner, J. (2015). Stream temperature patterns over 35 years in a
3410 managed forest of western Washington. *JAWRA Journal of the American Water Resources*
3411 *Association*, 51(5), 1418-1435.

3412 Reiter, M., Heffner, J. T., Beech, S., Turner, T., & Bilby, R. E. (2009). Temporal and Spatial Turbidity Patterns
3413 Over 30 Years in a Managed Forest of Western Washington 1. *JAWRA Journal of the American Water*
3414 *Resources Association*, 45(3), 793-808.

3415 Reiter, M., Johnson, S. L., Homyack, J., Jones, J. E., & James, P. L. (2020). Summer stream temperature
3416 changes following forest harvest in the headwaters of the Trask River watershed, Oregon Coast
3417 Range. *Ecohydrology*, 13(3), e2178.

3418 Roni, P., Beechie, T., Pess, G., & Hanson, K. (2015). Wood placement in river restoration: fact, fiction, and
3419 future direction. *Canadian Journal of Fisheries and Aquatic Sciences*, 72(3), 466-478.
3420 <https://doi.org/10.1139/cjfas-2014-0344>

3421 Roon, D. A., Dunham, J. B., & Groom, J. D. (2021a). Shade, light, and stream temperature responses to
3422 riparian thinning in second-growth redwood forests of northern California. *PloS One*, 16(2), e0246822–
3423 e0246822. <https://doi.org/10.1371/journal.pone.0246822>

3424 Roon, D. A., Dunham, J. B., & Torgersen, C. E. (2021b). A riverscape approach reveals downstream
3425 propagation of stream thermal responses to riparian thinning at multiple scales. *Ecosphere*, 12(10),
3426 e03775.

3427 Safeeq, M., Grant, G. E., Lewis, S. L., & Hayes, S. K. (2020). Disentangling effects of forest harvest on long-
3428 term hydrologic and sediment dynamics, western Cascades, Oregon. *Journal of Hydrology*, 580, 124259.

3429 Salo, E. O., & Cundy, T. W. (1986, February). Streamside management: forestry and fishery interactions.
3430 In *Proceedings of a conference sponsored by the College of Forest Resources, University of Washington*
3431 *and others, and held at the University of Washington*.

3432 Schuett-Hames, D., & Stewart, G. (2019a). Post-Harvest Change in Stand Structure, Tree Mortality and
3433 Tree Fall in Eastern Washington Riparian Buffers. Cooperative Monitoring Evaluation and Research
3434 Report. Washington State Forest Practices Adaptive Management Program. Washington Department of
3435 Natural Resources, Olympia, WA.

3436 Schuett-Hames, D. & Stewart, G. (2019b). Changes in stand structure, buffer tree mortality and riparian-
3437 associated functions 10 years after timber harvest adjacent to non-fish-bearing perennial streams in
3438 western Washington. Cooperative Monitoring Evaluation and Research Report. Washington State Forest
3439 Practices Adaptive Management Program. Washington Department of Natural Resources, Olympia, WA.

3440 Schuett-Hames, D., Martin, D., Mendoza, C., Flitcroft, R., & Haemmerle, H., (2015). Westside Type F
3441 Riparian Prescription Monitoring Project Technical Writing and Implementation Group (TWIG).
3442 Cooperative Monitoring Evaluation and Research Report. Washington State Forest Practices Adaptive
3443 Management Program. Washington Department of Natural Resources, Olympia, WA.

3444 Schuett-Hames, D., Roorbach, A., & Conrad, R. (2011). Results of the Westside Type N Buffer
3445 Characteristics, Integrity and Function Study Final Report. Cooperative Monitoring Evaluation and
3446 Research Report, CMER 12-1201, Washington Department of Natural Resources, Olympia, WA.

3447 Shah, N. W., & Nisbet, T. R. (2019a). The effects of forest clearance for peatland restoration on water
3448 quality. *Science of the Total Environment*, 693, 133617.

3449 Sievers, M., Hale, R., & Morrongiello, J. R. (2017). Do trout respond to riparian change? A meta-analysis
3450 with implications for restoration and management. *Freshwater Biology*, 62(3), 445–457.
3451 <https://doi.org/10.1111/fwb.12888>

3452 Six, L. J., Bilby, R. E., Reiter, M., James, P., & Villarin, L. (2022). Effects of current forest practices on
3453 organic matter dynamics in headwater streams at the Trask river watershed, Oregon. *Trees, Forests and*
3454 *People*, 8, 100233.

3455 Sobota, D. J., Gregory, S. V., & Sickle, J. V. (2006). Riparian tree fall directionality and modeling large
3456 wood recruitment to streams. *Canadian Journal of Forest Research*, 36(5), 1243–1254.
3457 <https://doi.org/10.1139/x06-022>

3458 Sugden, B. D., Steiner, R., & Jones, J. E. (2019). Streamside management zone effectiveness for water
3459 temperature control in Western Montana. *International Journal of Forest Engineering*, 30(2), 87-98.

3460 Sullivan, P. F., Neale, M. C., & Kendler, K. S. (2000). Genetic epidemiology of major depression: review
3461 and meta-analysis. *American journal of psychiatry*, 157(10), 1552-1562.

3462 Swanson, F. J., & Dyrness, C. T. (1975). Impact of clear-cutting and road construction on soil erosion by
3463 landslides in the western Cascade Range, Oregon. *Geology*, 3(7), 393-396.

3464 Swanson, F. J., Gregory, S. V., Sedell, J. R., & Campbell, A. G. (1982). Land-water interactions: the riparian
3465 zone.

3466 Swartz, A., Roon, D., Reiter, M., & Warren, D. (2020). Stream temperature responses to experimental
3467 riparian canopy gaps along forested headwaters in western Oregon. *Forest Ecology and
3468 Management*, 474, 118354.

3469 Sweeney, B. W., & Newbold, J. D. (2014). Streamside forest buffer width needed to protect stream water
3470 quality, habitat, and organisms: a literature review. *JAWRA Journal of the American Water Resources
3471 Association*, 50(3), 560-584.

3472 Teply, M., McGreer, D., & Ceder, K. (2014). Using simulation models to develop riparian buffer strip
3473 prescriptions. *Journal of Forestry*, 112(3), 302-311.

3474 Teply, M., McGreer, D., Schult, D., & Seymour, P. (2007). Simulating the effects of forest management on
3475 large woody debris in streams in northern Idaho. *Western Journal of Applied Forestry*, 22(2), 81-87.
3476 <https://doi.org/10.1093/wjaf/22.2.81>

3477 Vanderbilt, K. L., Lajtha, K., & Swanson, F. J. (2003). Biogeochemistry of unpolluted forested watersheds
3478 in the Oregon Cascades: temporal patterns of precipitation and stream nitrogen
3479 fluxes. *Biogeochemistry*, 62(1), 87-117.

3480 Warren, D. R., Keeton, W. S., Bechtold, H. A., & Rosi-Marshall, E. J. (2013). Comparing streambed light
3481 availability and canopy cover in streams with old-growth versus early-mature riparian forests in western
3482 Oregon. *Aquatic sciences*, 75(4), 547-558.

3483 Wing, M. G., & Skaugset, A. (2002). Relationships of channel characteristics, land ownership, and land
3484 use patterns to large woody debris in western Oregon streams. *Canadian Journal of Fisheries and
3485 Aquatic Sciences*, 59(5), 796-807.

3486 Wise, E. K. (2010). Tree ring record of streamflow and drought in the upper Snake River. *Water Resources
3487 Research*, 46(11).

3488 Wissmar, R.C., Beer, W.N. & Timm, R.K. Spatially explicit estimates of erosion-risk indices and variable
3489 riparian buffer widths in watersheds. *Aquat. Sci.* 66, 446-455 (2004). [https://doi.org/10.1007/s00027-
3490 004-0714-9](https://doi.org/10.1007/s00027-004-0714-9)

3491 Yang, Y., Hart, S. C., McCorkle, E. P., Stacy, E. M., Barnes, M. E., Hunsaker, C. T., ... & Berhe, A. A. (2021).
3492 Stream water chemistry in mixed-conifer headwater basins: role of water sources, seasonality,
3493 watershed characteristics, and disturbances. *Ecosystems*, 24(8), 1853-1874.

3494 Yeung, A. C., Stenroth, K., & Richardson, J. S. (2019). Modelling biophysical controls on stream organic
3495 matter standing stocks under a range of forest harvesting impacts. *Limnologica*, 78, 125714.

3496 Zhang, X., & Shu, C. W. (2010). On positivity-preserving high order discontinuous Galerkin schemes for
3497 compressible Euler equations on rectangular meshes. *Journal of Computational Physics*, 229(23), 8918-
3498 8934.

3499

3500

3501

3502

3503

3504

3505

3506

3507

3508

3509

3510

3511

3512

3513

3514

3515

3516

3517

3518

3519

3520 Appendix I

3521

3522 Table A-1. List of treatments, variables, metrics, and results from publications reviewed for information on litter, organic matter, and
 3523 nutrient inputs.

Reference	Treatment	Variables	Metrics	Notes	Results
Anderson et al., 2007	Upland stands either thinned to 198 TPA or unthinned and ranged from 500-865 TPA. Within thinned stands, 10% of the area was harvested to create patch openings. streamside buffers ranged in width from <5 m to 150 m.	Microsite, microclimate, stand structure, canopy cover	Microsite and microclimate data (humidity, temperature sensors). Stand basal area. Canopy cover was estimated through photographic techniques.	Many of the reported differences in temperature and humidity were considerable but not significant. Results for changes in upland areas not reported here.	Subtle microclimatic changes as mean temperature maxima in treated stands were 1 to 4°C higher than in untreated stands. Buffer widths greater than or equal to 15 m experienced a daily maximum air temperature above stream center of less than 1°C greater than untreated stands. Daily minimum relative humidity for buffers 15 m or greater was less than 5 percent lower than for unthinned stands. Air temperatures were significantly higher in patch openings (+6 to +9°C), and within buffers adjacent to patch openings (+3.5°C), than in untreated stands.
Bilby & Heffner, 2016	Various wind speeds for young and old-growth conifer and deciduous forests. Distance of litter delivery.	Litter input	Models were developed with site characteristics and litter release experiments from sites along Humphrey Creek in the cascade mountains of western Washington.	Wind speeds, direction, and litter release data were collected for only one year in one area of western Washington.	The results of the linear mixed model developed by the authors showed the strongest relationship for recruitment distance was with wind speed (p<0.0001). Using this relationship the authors estimated that the effective delivery area could be increased by 67-81% by doubling wind speed. The other significant relationship was with stand age for needles (not alder leaves). Needles released from mature stands traveled further distances. This is likely due to the higher height of the canopy in the mature stands.
Deval et al., 2021	clearcut to stream, 50% shade retention, with site management operations including pile burning and competition release herbicide application.	Changes in nitrogen and phosphorus compounds.	monthly grab samples from multiple flume sites pre- and post-harvest, laboratory chemical analysis	Data was compared from pre-harvest to post experimental harvest (PH-I), and post operational harvest (PH-II)	The response in NO3 + NO2 concentrations was negligible at all treatment sites following the road construction activities. However, NO3 + NO2 concentrations during the PH-I period increased significantly (p < 0.001) at all treatment sites. Similar to the PH-I period, all watersheds experienced significant increases in NO3 + NO2 concentration during the PH-II treatment period. Overall, the cumulative mean NO3 + NO2 load from all watersheds followed an increasing trend with initial signs of recovery in one treatment watershed after 2014. Mean monthly TP concentrations showed no significant changes in the concentrations

					during the post-road and PH-I treatment periods. However, a statistically significant increase in TP concentrations ($p < 0.001$) occurred at all sites, including the downstream cumulative sites, during PH-II. Generally, OP concentrations throughout the study remained near the minimum detectable concentrations
Gravelle et al., 2009	clearcut to stream, 50% shade retention, uncut reference	Changes in nitrogen and phosphorus compounds.	monthly grab samples from multiple flume sites pre- and post-harvest, laboratory chemical analysis	Data was compared in three treatment periods: pre-harvest, under road construction, post-harvest.	Results showed significant increases in monthly mean NO ₃ and NO ₂ following clear-cut harvest treatments relative to the pre-harvest, and road construction periods. Monthly nitrate responses showed progressively increasing concentrations for 3 years after harvest before declining. Significant increases in NO ₃ and NO ₂ concentrations were also found further downstream but at values lower than those immediately downstream from harvest treatments. No significant changes of in-stream concentration of any other nutrient recorded were found between time periods and treatments except for one downstream site that showed a small increase in orthophosphate by 0.01 mg P L ⁻¹ .
Hart et al., 2013	(1) a no cut or fence control; (2) cut and remove a 5 x 8 m section adjacent to stream for plants < 10 cm DBH and >12 cm; and (3) 5 m fence extending underground and parallel to the stream to block litter moving downslope from reaching stream	Litter inputs, vegetation composition, topography, litter chemistry	Litter collected with lateral and vertical traps. Litter was sorted by type, time of fall, spatial source, and quantified by weight. Vegetation, LW, and Site characteristics were quantified for each plot.	This study took place within 5 contiguous watersheds located in the central Coast Range of Oregon.	Deciduous forests dominated by red alder delivered greater vertical and lateral inputs to streams than did coniferous forests dominated by Douglas-fir by 110 g/m ² (28.6–191.6) and 46 g/m (1.2–94.5), respectively. Annual lateral litter input increased with slope at deciduous sites ($R^2 = 0.4073$, $p = 0.0771$) but not at coniferous sites ($R^2 = 0.1863$, $p = 0.2855$). Total nitrogen flux to streams at deciduous sites was twice as much as recorded at coniferous sites. However, the nitrogen flux had a seasonal effect with the majority of N flux occurring in autumn at the deciduous sites. The authors of this study conclude by suggesting management in riparian areas consider utilizing deciduous species such as red alder for greater total N input to aquatic and terrestrial ecosystems with increased shade and large woody debris provided by coniferous species.
Kiffney & Richardson, 2010	clearcut to stream, 10 m buffer, 30 m buffer, uncut control	Litter inputs.	Litter was separated into broadleaf deciduous, twig, needles, and other (seeds, cones, and moss) categories following collection and subsequently dried and weighed using a microbalance.	Sites were measured over an 8-year period and included clear-cut (n=3), 10-m buffered reserve (n=3), 30-m buffered reserve (n=2), and uncut control (n=2) treatments.	Inputs consisting of needles and twigs were significantly lower adjacent to clearcuts compared to other treatments, while deciduous inputs were higher in clearcuts compared to other treatments. For example, one year post-treatment, needle inputs were 56x higher during the Fall into control and buffered treatments than into the clearcut. Needle inputs remained 6x higher in the buffer and control sites through year 7, and 3-6x higher in year 8 than in the clearcut sites. Twig inputs into the control and buffered sites were ~25x higher than in the clearcut sites in the first year after treatment. There was no significant difference in treatment for deciduous litter but a trend of increasing deciduous litter input in the clear cut was observed in the data. The linear relationship between

					reserve width and litter inputs was strongest in the first year after treatment, explaining ~57% of the variation, but the relationship could only explain ~17% of the variation in litter input by buffer width by year 8 (i.e., the relationship degraded over time).
McIntyre et al., 2018	(1) unharvested reference, (2) 100% treatment, a two-sided 50-ft riparian buffer along the entire Riparian Management Zone (RMZ), (3) FP treatment, a two-sided 50-ft riparian buffer along at least 50% of the RMZ (4) 0% treatment, clearcut to stream edge (no-buffer).	Litter inputs from litter traps situated along channel	Sorted by litter type (conifer needles, deciduous leaves, woody components, etc.). Compared between treatments by dry weight.	Authors of the study identify a lack of information on local meteorology as a primary limitation to the study. This, the authors suggest, would have allowed for a more detailed analysis including information on hydrologic mass balance.	Showed a decrease in TOTAL litterfall input in the FP (P = 0.0034) and 0% (P = 0.0001) treatments between pre- and post-treatment periods. LEAF litterfall (deciduous and conifer leaves combined) input decreased in the FP (P = 0.0114) and 0% (P <0.0001) treatments in the post-treatment period. In addition, CONIF (conifer needles and scales) litterfall input decreased in the FP (P = 0.0437) and 0% (P <0.0001) treatments, DECID (deciduous leaves) in the 0% (P <0.0001) treatment, WOOD (twigs and cones) in the FP (P = 0.0044) and 0% (P = 0.0153) treatments, and MISC (e.g., moss and flowers) in the 0% (P = 0.0422) treatment. Results for comparison of the post-harvest effects between treatments showed LEAF litterfall input decreased in the 0% treatment relative to the reference (P = 0.0040), 100% (P = 0.0008), and FP (P = 0.0267) treatments. Likewise, there was a decrease in DECID litterfall input in the 0% treatment relative to the Reference (P = 0.0001), 100% (P <0.0001), and FP (P = 0.0015) treatments. Statistical differences were only detected for deciduous inputs between the 0% treatment and the other treatments.
McIntyre et al., 2021	<u>1) unharvested reference, 2) 100% treatment, a two-sided 50-ft riparian buffer along the entire RMZ, 3) FP treatment a two-sided 50-ft riparian buffer along at least 50% of the RMZ, (4) 0% treatment, clearcut to stream edge (no-buffer).</u>	stream discharge, nitrogen export		Type N (non-fish-bearing streams). Hard-Rock study.	Discharge increased by 5-7% on average in the 100% treatments while increasing between 26-66% in the FP and 0% treatments Results for harvest effects on total Nitrogen export showed significant (P <0.05) treatment effects were present in the FP treatment and in the 0% treatment in the post-harvest (2-years immediately following harvest) and extended periods (7 and 8 years post-harvest) relative to the reference sites, Analysis showed an increase in total-N export of 5.73 (P = 0.121), 10.85 (P = 0.006), and 15.94 (P = 0.000) kg/ha/yr post-harvest in the 100%, FP, and 0% treatments, respectively, and of 6.20 (P = 0.095), 5.34 (P = 0.147), and 8.49 (P = 0.026) kg/ha/yr in the extended period. The authors conclude that the 100% treatment was generally the most effective in minimizing changes in total-N from pre-harvest conditions, the FP was intermediate, and the 0% treatment was least effective. At the end of the study (8 years), only one site had recovered to pre-harvest nitrate-N levels.

Murray et al., 2000	7% and 33% watershed upland harvest. Harvest extended to stream channel.	stream chemistry, stream temperatures, sediment input	Chemistry and pH tested on water grab samples; Daily max, min, and average temperatures collected with Stowaway dataloggers; Sediment change detected with turbidity meters.	Results reflect differences in stream conditions 11-15 years post-harvest only. No data collected in first decade following treatment.	10-15 years post-harvest mean maximum daily summer temperatures were still significantly higher (15.4 °C) and mean maximum daily winter temperatures were lower (3.7 °C) than in the reference streams (12.1 °C and 6.0 °C) respectively. Also, winter minimum temperatures for one of the harvested watersheds reached 1.2 °C compared to a winter minimum of 6 °C There were no significant differences in stream chemistry with the exception of calcium and magnesium being consistently higher in the unharvested reference watersheds. No detectable difference in turbidity between treatment and reference watershed streams 10-5 years post-treatment. The stream temperature changes were significant but did not exceed the 16 °C threshold used as a standard for salmonid habitat.
Six et al., 2022	Clearcut with no leave trees or retention buffer (CC), clearcut with leave trees (CC w/LT; retention of 5 trees per hectare/2 trees per acre), and clearcut with 15 m wide retention buffer (CC c/B) and two uncut references (REF 1, and 2) along headwater streams	Litter input, LW recruitment	litter traps, in-stream LW volume, weight, and counts.	No replication of treatment sites. Data was analyzed with descriptive and graphical representation only.	Results showed a reduction of canopy cover from 91.4% to 34.4% in the clearcut treatment with no leave trees, from 89.8% to 76.1% in the clearcut treatment with leave trees, and from 89.5% to 86.9% in the clearcut treatment with the 15 m retention buffer. Post harvest litter delivery decreased for the clearcut with no leave trees but increased for both the clearcut with leave tree and clear cut with retention buffer.
Vanderbilt et al., 2003	Datasets (ranging from 20-30 years) from six watersheds in the H.J. Andrews Experimental Watershed.	Nitrogen concentration in streams, precipitation patterns	regression analysis of annual N inputs and outputs with annual precipitation and stream discharge to analyze patterns.	These results come from a coastal climate of western Oregon. The authors warn that the controls on in stream N concentrations will likely differ in different regions.	Total annual discharge was a positive predictor of annual DON export in all watersheds with r2 values ranging from 0.42 to 0.79. In contrast, significant relationships between total annual discharge and annual export of NO3-N, NH4-N, and PON were not found in all watersheds. DON concentrations increased in the fall in every watershed. The increase in concentration began in July or August with the earliest rain events, and peak DON concentrations occurred in October through December. DON concentrations then declined during the winter months. The authors conclude that total annual stream discharge was a positive predictor of DON output suggesting a relationship to precipitation.

Yang et al., 2021	Young stands with high shrub cover (> 50%) masticated to < 10% shrub cover. trees removed to a target basal area range of 27–55 m ² ha ⁻¹ .	Drought, nutrients, dissolved organic carbon	Stream water samples grab samples and chemical analysis	Because of difficulties with accessibility due to weather-related phenomena (particularly during winter months), snowmelt and soil samples were restricted to the lower elevation site.	Drought alone altered DOC in stream water, and DOC:DON in soil solution in unthinned (control) watersheds. The volume-weighted concentration of DOC was 62% lower, and DOC:DON was 82% lower in stream water in years during drought than in years prior to drought. Drought combined with thinning altered DOC and DIN in stream water, and DON and TDN in soil solution. For stream water, volume-weighted concentrations of DOC were 66- 94% higher in thinned watersheds than in control watersheds for all three consecutive drought years following thinning. No differences in DOC concentrations were found between thinned and control watersheds before thinning. Watershed characteristics inconsistently explained the variation in volume-weighted mean annual values of stream water chemistry among different watersheds
Yeung et al., 2019	Range of forest harvest intensities	Litter inputs, CPOM in streams	stream temperature, streamflow, litter traps, CPOM decay rates	Authors point out that model results are primarily applicable to stream reaches similar to those used in the study and may not be suitable for streams where large wood is a dominant structure retaining CPOM.	The simulation predicted that litter input reduction from timber harvest was the strongest control on CPOM in streams relative to streamflow and temperature variability. The effects of litterfall reduction were at least an order of magnitude higher than streamflow increases in depleting in-stream CPOM. Significant CPOM depletions were most likely when there was a 50% or greater reduction in litterfall following harvest. The caveat of this study is that it did not include LW dynamics in preserving CPOM post-harvest. As other studies have shown, harvest can increase in-stream LW, and in-stream LW can act as a catchment for CPOM.

3524

3525 Table A-2. List of treatments, variables, metrics, and results from publications reviewed for information on large wood (LW), wood loads, and wood recruitment.

Reference	Treatment	Variables	Metrics	Notes	Results
Anderson & Meleason, 2009	Buffer averaging 69 m adjacent to thinning and a 0.4 patch opening; variable width buffer averaging 22 m adjacent to thinning and a 0.4 patch opening.	Instream wood load, understory vegetation cover	Percent cover of LW in streams and in riparian area, %cover shrubs, herbs, moss.		LW changes were non-significant, decrease in treatment reaches with greatest pre-treatment values 5 years post-treatment caused homogenization of LW. Gaps (patch openings) showed the highest changes increase in herbaceous cover, decrease in shrub cover. Moss cover increased in thinned areas but decreased in gaps. LW and vegetation changes insensitive to treatment buffers > 15 m.

Bahuguna et al., 2010	Two buffer widths on each side of the stream (10 m and 30 m) with upland clearcuts, and an unharvested control.	LW, Stand Structure, mortality	Strip plot sampling method running parallel to the stream to collect data on stand metrics.	Experimental design included 3 replicates of each treatment. Data was collected annually for one year pre- and 8 years post-treatment. Vancouver, B.C.	Following harvest, 11% of initially standing timber was blown down in the first and second years in the 10 m buffer, compared to 4% in the 30 m buffer, and 1% in the unharvested controls. Small diameter trees were significantly more represented in streams - 77% of LW was in the 10 cm - 20 cm diameter class while the mean diameter of standing trees in riparian buffers was 30 cm. By 8 years post-harvest, a significant amount of annual mortality occurred in the unharvested control at 30%, compared to 15% in both 30 m and 10 m buffers.
Benda et al., 2016	Simulated treatments of single or double entry thinning with and without a 10-m no cut buffer, with and without mechanical tipping of stems into streams. Thinning encompassed 5-20 % thinning.	instream LW volume	ORGANON growth models simulated secondary forest growth. The model was run for 100 years in 5-year time steps.	used the reach scale wood model (RSWM) developed for the Alcea watershed in central coastal Oregon. Data was sourced from FIA.	Single entry thinning reduced in-stream wood by 33 and 66% after a century, relative to reference streams when one and both sides of the channel were harvested. Adding a 10 m buffer reduced total loss to 7 and 14%. Mechanical tipping of 14 and 12% of cut stems were sufficient in offsetting the loss of instream wood without and with buffers. Double entry thinning without a buffer resulted in 42 and 84% loss of in stream wood relative to the reference streams when one or both sides of the channel were harvested. Adding a 10 m buffer changed reductions of in stream wood to 11 and 22% for one- and two-sided channel harvest. To offset the total predicted reduction of in stream wood for the double entry thinning would require tipping of 10 and 7% of cut stems without and with 10 m buffers.
Burton et al., 2016	70-m buffer representative of one site potential tree, 15-m buffer, 6-m buffer. Outside of buffer, all treatment stands were thinned first to 200 trees per hectare (tph), then again to 85 tph ~ 10 years later. Uncut reference was ~400 tph.	LW recruitment, In-stream wood volume, biomass, and	LW volume, LW characteristics and source evidence, reach and stream characteristics.	Wood surveys were carried out at four times during the study: (1) prior to the first thinning, (2) five years after the first thinning, (3) 9-13 years after the first thinning and just prior to the second thinning, and (4) one year after the second thinning.	In-stream wood volume increased significantly with drainage basin area; for every 1-ha increase in drainage basin area, wood volume increased by 0.63%. LW volume was slightly higher in the streams adjacent to 6 m buffers than in streams bordered by 15 and 70 m buffers. The higher volume of wood in the 6 m buffers began 5 years after the first harvest and maintained through 1 year after the second harvest (end of study). . 82% to 85% of all wood inputs (early- and late-stage decay) were sourced from within 15 m of the streams (90% of early-stage decay wood could be sourced, only 45% of late-stage decay wood could be sourced).

Chen et al., 2005	All harvested streams were clearcut to stream edge. Wildfire streams had no post-fire harvest	Instream wood load, biomass, carbon pool	LW count, volume, decay class, size		LW volume, biomass, and carbon pools were significantly higher in streams adjacent to areas recently disturbed by timber harvest (~10 years) or wildfire (~40 years) than in streams passing through old-growth forests. There was no significant difference in in-stream LW between old-growth riparian areas and areas harvested > 30 years ago. The wildfire sites had significantly higher LW values than both the harvested sites. The authors conclude: (1) LWD input in old growth forested streams was relatively stable based on statistical significance. They also speculate: (1) timber harvesting activities would cause a short-term increase of LWD stocks and might greatly reduce LWD loadings over a long-term, and (2) wildfire disturbance would delay LWD recruitment because not all burnt trees would fall in the stream immediately after the wildfire, based on trends in, and extrapolation of the data.
Chen et al., 2006	A total of 35 sites with stream orders ranging from 1-5 (grouped into 4 stream size categories (I = first order; II = second to third order; III = third to fourth order; IV = fourth to fifth order) were selected to measure spatial distribution and variability of LW characteristics	LW, defined as having a diameter of > 0.1 m and a length > 1.0 m.	LW size, volume, density, and biomass. Multiple stream channel features obtained from readily available physiographic and forest cover data.	Study sites were selected based on the following criteria. (1) the streams were in areas of intact mature riparian forests (>80 years); (2) the stream side forests were not disturbed by human activities, such as harvesting, road building; (3) the streams were not salvaged.	Results from this study show that LW size, volume, and biomass generally increased with increasing stream size. For example, the mean LWD diameter in stream size I (16.4 cm) was lower than that in stream size III (20.6 cm) and IV (20.5 cm), respectively. Mean LW length also increases with stream size from 2.3 m in size I, 2.9 m in size II, 3.1 m in size III, and 3.9 m in size IV. Stream IV had the highest mean volume (0.18 m ³), significantly higher than stream size I (0.06 m ³). LW density (pieces per 100 m ² of stream area), however, decreased as stream size increased. For example, LW density (defined as piece numbers per 100 m ²) numbers were 19, 17, 12, and 4 for stream size I, II, III, and IV respectively. Increases in channel bank full width (R ² = 0.52) and stream area (R ² = 0.58) was found to be strongly inversely correlated with LW density.
Ehinger et al., 2021	1) Buffers encompassing the full width (50 feet), 2) <50ft buffers, 3) Unbuffered, harvested to the edge of the			Soft Rock study. Only descriptive statistics were applied for changes in stand structure and wood loading.	There was little post-harvest large wood input in reference sites: an average of 4.3 pieces and 0.34 m ³ of combined in- and over-channel volume per 100 m of channel. In contrast, the full buffer sites and <50 ft buffer sites received an average of 23 and 10 pieces/100 m and 2.3 and 0.7 m ³ /100 m of large wood, respectively. Piece counts remained stable in the reference sites through year 3 post-harvest, increased in the

	channel, and 4) Reference sites in unharvested forests.			Small sample sizes.	full buffer and unbuffered sites (8 and 13%, respectively), and decreased in the <50 ft buffers (-15%).
Fox & Bolton, 2007	LW values from 150 stream segments located in unmanaged watersheds, across all of Washington State	Instream LW, geomorphology, forest zone, disturbance regimes	Descriptive statistics for LW volume and quantity, channel geomorphology, forest habitat type, disturbance regimes.	the authors warn that these values for reference conditions are only applicable to streams with bank-full widths 1-100 m, gradients 0.1%-47%, elevations 91-1,906 m, drainage areas 0.4-325 km ² , glacial and rain- or snow-dominated origins, forest types common to the Pacific Northwest.	Results showed that in-stream wood volume increased with drainage area and as streams became less confined. Bank full width (BFW) was the single greatest predictor of in-stream wood volumes relative to other predictor variables. However, this result comes with the caveat that other processes and geomorphologies (e.g., channel bed form, gradient, confinement) are also important in the mechanisms for wood recruitment, modeling in this study showed too much inconsistency with these predictor variables to draw strong conclusions. In-stream wood volume also increased with adjacent riparian timber age as determined by the last stand replacing fire. The authors developed thresholds for expected "key piece volume (m ³)" (pieces with independent stability) of wood for three BFW classes (20-30 m, >30 – 50 m, > 50 m width) per 100 m stream length for streams with BFW greater than 20 m. From percentile distributions the authors recommend minimum volumes, defined by the 25th percentiles, of approximately 9.7 m ³ for the 20- to 30-m BFW class, 10.5 m ³ for the 30- to 50-m BFW class, and 10.7 m ³ for channels greater than 50 m BFW per 100 m length of stream.
Gomi et al., 2001	Five management or disturbance regimes: old growth (OG), recent clear-cut (CC; 3 years), young conifer forest (YC; 37 years after clear-cut), young alder (YA; 30 years after clear-cut), and recent landslide and debris flow channels (LS)	LW quantity and distribution, sediment quantity and distribution, landslide frequency, harvest intensities	LW counts, LW characteristics, stream characteristics.	Results are highly variable among treatments	in-channel numbers of LW pieces were significantly higher in YC and CC sites when compared to OG, YA, and LS sites. The number of LW pieces was highest in YC streams even though logging concluded 3 decades prior to sampling. LW volume per 100 m of stream length in YC was twice that in OG. The total volume of LW per 100 m associated with CC channels was half that in OG channels. The authors conclude (i) inputs of logging slash and unmerchantable logs significantly increase the abundance of in-channel woody debris; (ii) in the absence of landslides or debris flows, these woody materials remain in the channel 50–100 years after logging.

Hough-Snee et al., 2016	In-stream wood volume and frequency were quantified across multiple sub basins.	LW frequency and volume, hydrologic and geomorphic attributes	Models were calibrated with site characteristics from multiple riparian stands in the Columbia River Basin.	Results show a high level of variability between sub basins studied. The overall model shows site (watershed) was an important predictor.	In stream wood volume and frequency were distinctly different across all seven sub-basins. According to random forest (RF) models, mean annual precipitation, riparian large tree cover, and individual watershed were the three most important predictors of wood volume and frequency, overall. Sinuosity and measures of streamflow and stream power were relatively weak predictors of wood volume and frequency. Final RF models explained 43.5% of the variance in volume and 42.0% of the variance in frequency of in stream wood loads. Depending on the sub basin wood volume and frequency was positively correlated with forest cover, watershed area, large tree cover, 25-year flood event stream power, riparian conifer cover, and precipitation. Negative correlations, depending on sub basin, of wood volume and frequency with baseflow discharge, riparian woody cover, watershed area, and large tree cover. Given the heterogeneous results across all sub-basins studied, the authors conclude by emphasizing the importance of incorporating local data and context when building wood models to inform future management decisions.
Hyatt & Naiman, 2001	LW data was collected from multiple sites in the Queets River Watershed.	LW in stream and in riparian forests.	Increment cores from in-stream LW were cross-dated to estimate the time LW was recruited. LW pieces in decay were dated using carbon-dating. A depletion curve was fitted for LW recruited between 1599 and 1997.	The depletion constant was developed for a large, mostly alluvial river and should probably not be applied to smaller streams	Results from this study indicate that the half-life of stream LW to be approximately 20 years, suggesting that current LW will either be exported, broken down, or buried within 3 to 5 decades (for conifers). Hardwoods were better represented in riparian forests than as in-stream LW, and conversely, conifers were better represented as in-stream LW than in adjacent forests suggesting that LW originating from hardwoods is depleted faster than conifers.
Jackson & Wohl, 2015	In-stream wood volume and frequency were quantified along 33 pool-riffle or plane-bed stream reaches in the Arapaho and Roosevelt National Forests in Colorado.	Sediment storage, channel geometry, in-stream wood load, and forest stand characteristics	Wood loads, wood jam volumes, log jam frequencies, residual pool volume, and fine sediment storage around wood, stand age, and	Old growth defined as forests ≥ 200 years. Age range of young forests not reported. Sample sizes include 10 old-growth and	Results indicated that channel wood load (OG = $304.4 + 161.1$; $Y = 197.8 + 245.5 \text{ m}^3 / \text{ha}$), floodplain wood load (OG = $109.4 + 80$; $Y = 47.1 + 52.8 \text{ m}^3 / \text{ha}$), and total wood load (OG = $154.7 + 64.1$; $Y = 87.8 + 100.6 \text{ m}^3 / \text{ha}$) per 100 m length of stream and per unit surface area were significantly larger in streams of old-growth forests than in young forests. Streams in old-growth forests also had significantly more wood in jams, and more total wood jams per unit length of channel than in younger forests (jam wood volume: OG = $7.10 + 6.9 \text{ m}^3$; $Y =$

			disturbance history.	23 younger forests.	1.71 + 2.81 m ³). Although wood load in streams draining from pine beetle infested forests did not differ significantly from healthy forests, best subset regression (following principal component analysis) indicated that elevation, stand age, and pine beetle infestation were the best predictors of wood load in channels and on floodplains.
Jackson et al., 2001	3 unthinned riparian buffers; 1 with a partial buffer; 1 with a buffer of non-merchantable trees; and 6 were clearcut to the stream edge. Buffers ranged from 15 to 21 m wide, partial buffers were as thin as 2.3 m.	Instream LW, particle size, surface roughness	LW as functional and nonfunctional (not altering flow hydraulics). Particle size distributions.	Data collected for only 1-year pre- and 1-month post-harvest. These results only describe immediate effects of harvest on stream conditions.	Increased slash debris (LW) provided shade for the harvested streams but trapped sediments and prevented fluvial transport. The percentage of fine particles increased from 12 to 44% because of bank failure and increased surface roughness. This was a short-term study on small headwater streams. Sediment and LW conditions in the unharvested and buffered streams remained relatively unchanged during the study.
Liquori, 2006	Data were collected from 20 riparian buffer sites that had all been clearcut within three years of sampling with standard no-cut 25 ft or 50-100 ft buffers for non-fish-bearing and fish-bearing streams, respectively.	Tree and tree fall characteristics, Site characteristics	Tree characteristic data estimated cause of mortality, and distance to the stream. Tree recruitment probability curves were developed as a function of tree height.		Within no-cut buffers windthrow caused mortality was up to 3 times greater than competition induced mortality for 3 years following treatment Tree fall direction was heavily biased towards the channel regardless of channel or buffer orientation and tree fall probability was highest in the outer areas of the buffers (adjacent to the harvest area). Tree fall rates and direction were also heavily biased by species with western hemlock and Pacific silver fir having the highest fall rates compared to Douglas-fir, western red cedar, and red alder.

<p>Martin & Grotefendt, 2007</p>	<p>Buffer widths a minimum of 20 m. Multiple buffer widths and harvest intensities.</p>	<p>Instream wood load, stand mortality</p>	<p>Counts of downed wood, tree stumps, stand characteristics, instream wood from aerial photographs taken post-logging</p>	<p>Stand and stream characteristic, and LW data was surveyed from aerial photographs.</p>	<p>Results showed significantly higher mortality, significantly lower stand density, and a significantly higher proportion of LW recruitment from the buffer zones of the treatment sites than in the reference sites. Differences in mortality for the treatment sites were similar to the reference sites for the first 0-10 m from the stream (22% increase). However, mortality in the outer half of the buffers (10-20 m) from the stream in the treatment sites was more than double (120% increase) what was observed in the reference sites. This caused a change in the LW recruitment source distance curves, with a larger proportion of LW recruitment coming from greater distances in logged watersheds. LW recruitment based on the proportion of stand recruited (PSR) was significantly higher in the buffered units compared to the reference units. However, PSR from the inner 0-20 m was only 17% greater in the buffer units than in the reference units; while PSR of the outer unit (10 – 20 m) was more than double in the buffered units than in the reference units. The researchers conclude that the increase in mortality was caused by an increased susceptibility to windthrow. They estimate that future recruitment potential from the logged sites diminished by 10% relative to the unlogged reference sites.</p>
<p>May & Gresswell, 2003</p>	<p>Survey of LW in three second-order streams and the mainstem of the North Fork of Cherry creek.</p>	<p>LW, delivery mechanism</p>	<p>LW > 20 cm diameter, and >2 m length was categorized by 4 delivery mechanisms, Delivery process, disturbance type, and channel characteristics.</p>	<p>Although mean age of Douglas-fir trees was identified to be excess of 300 years old, further information on differences in stand structure or development stage between sites are not included.</p>	<p>Processes of slope instability were shown to be important conveyors of wood from upland forests to small colluvial channels. In the larger alluvial channels, windthrow was found to be the dominant recruitment process from adjacent riparian area. 80% of total wood pieces and 80% of total wood volume recruited to colluvial streams originated from trees rooted within 50 m of the channel. In the alluvial channel, 80% of the pieces of wood and 50% of the total volume originated from trees which came from 30 m of the channel. The primary function of wood in colluvial channels was sediment storage (40%) and small wood storage (20%). The primary function of wood in alluvial channels is bank scour (26%), stream bed scour (26%), and sediment storage (14%).</p>
<p>McIntyre et al., 2021</p>	<p>(1) unharvested reference, (2) 100% treatment, a two-sided 50-ft riparian buffer along the entire Riparian Management Zone</p>			<p>Hard Rock Study Physical constraints such as a lack of suitable low gradient reaches and/or issues with</p>	<p>Large wood recruitment to the channel was greater in the 100% and FPB RMZs than in the reference for each pre- to post-harvest time interval. Eight years post-harvest mean recruitment of large wood volume was two to nearly three times greater in 100% and FPB RMZs than in the references. Annual LW recruitment rates were greatest during the first two years, then decreased. However, these differences were</p>

	(RMZ), (2) FP treatment a two-sided 50-ft riparian buffer along at least 50% of the RMZ, (3) 0% treatment, clearcut to stream edge (no-buffer).			accessibility related to weather limited downstream measurements of exports to just eight sites.	not significant between any treatment comparisons, likely due to the high variability in the data. Mean LW loading (pieces per meter of stream) differed significantly between treatments in the magnitude of change overtime. Results showed a 66% ($P < 0.001$), 44% ($P = 0.05$) and 47% ($P = 0.01$) increase in mean large wood density in the 100%, FP and 0% treatments, respectively, in the first 2 years post-harvest compared with the pre-harvest period and after controlling for temporal changes in the references. Five years post-treatment the FP continued to increase 42% ($P = 0.08$), and again 8 years post-treatment (41%; $P = 0.09$). From 2-8 years post-harvest LW density in the 100% treatment stabilized and began to decrease in the 0% treatment.
Meleason et al., 2003	Multiple buffer widths and upland harvest intensities	Change in instream wood load over time	Simulation metrics for forest growth, tree breakage, and in-channel process	A potential limitation of growth models in that they lack the ability to predict responses to novel climatic conditions different than those of the past.	Simulation results predicted clear-cut to stream accumulated little LW immediately following treatment and little change over time. Maximum in-stream LW loads were predicted for streams with no-cut buffers >30 m for 500-year-old forests (500 years post treatment). Streams with 6 m wide buffers predicted only 32% of pre-harvest standing LW loads after 240 years. Forest plantations with > 10 m buffer widths contributed minimal LW to the stream from outside the buffer zone.
Nowakowski & Wohl, 2008	History of regulated and unregulated timber harvest practices.	Instream wood volume	LW volume, LW characteristics source evidence, buffer widths, reach and stream characteristics.		In-stream LW was 2-3 times lower in a watershed with a history (>100 years) of timber harvest (1.1 m ³ /100 m) when compared to unmanaged reference watersheds (3.3 m ³ /100 m). Valley characteristics (elevation, forest type, forest stand density, etc.) consistently explained more of the variability in wood load (42-80%) than channel characteristics (21-33%; reach gradient, channel width, etc.). Across all streams, the highest explanatory power of all models tested produced land use (managed vs unmanaged), and basal area as a significant predictor of wood loads ($r^2 = 0.8048$). For the unmanaged watershed the model produced stream valley sideslope gradient as the single best predictor of wood load ($r^2 = 0.5748$). Shear stress was the best predictor of wood load in the managed watersheds ($r^2 = 0.2403$). When the significant valley and channel characteristics of the managed and unmanaged watersheds were controlled for, the significant difference in wood loads between managed and unmanaged watersheds were enhanced ($p = 0.0006$). Managed watersheds

					(1.1 m ³ /100 m) had, on average, 2-3 times lower in-stream wood loads than unmanaged (3.3 m ³ /100 m) watersheds.
Reid & Hassan, 2020	Clearcut to stream and buffer widths that range from 1-70 m. Models were developed for 3 harvest scenarios (1: no-harvest; 2 partial loss of riparian forests; 3 intensive harvest in the riparian zone)	Instream LW	Models were calibrated with long-term data for site and LW characteristics in treatment reaches dating back to 1973.	One caveat of this model is it doesn't account for as much variability on stream configuration or valley morphologies that are likely to affect LW storage.	Results of the model show evidence that wood storage in streams of harvested reaches its minimum value in 50 years or more following loss of LW input, decay, and export of current stock. Recovery of LW volume in-streams following harvest is estimated to take approximately 150-200 years. The pattern and intensity of the harvesting operation had little effect on LW loss and recovery times but did affect the estimated magnitude of LW volume loss in the first 50 – 80 years. The authors conclude that the results show evidence that timber harvest has a long-term effect on LW storage and loading dynamics even with protective buffers. However, buffers can ameliorate the magnitude of LW loss during the recovery period.
Schuett-Hames & Stewart, 2019a	Buffer prescriptions for standard shade rule (a 30-ft no-cut buffer width, and thinning 30-75 ft from the stream), and all available shade rule (requires retention of all shade providing trees in this area) for eastern Washington.	LW recruitment, instream wood volume, mortality, stand structure	LW volume, LW characteristics, LW source evidence, reach and stream characteristics, basin metrics, stand metrics	Short-term study. Results only for 5 years post-harvest. The authors note that LW recruitment is a process that can change over decadal time scales.	Results showed cumulative wood recruitment from tree fall over the five-year post-harvest interval was highest in the standard shade rule (SR) group, lower in the all-available-shade rule (AAS) group and lowest in the reference (REF) group. The SR and AAS rates by volume were nearly 300% and 50% higher than the REF rates, respectively. Most recruiting fallen trees originated in the first 30 feet (76%, 72%, and 64% for the REF, AAS and SR groups, respectively), while the proportion from the inner zone (30–75 feet from the stream) was ~10% greater for the SR group compared to the AAS and REF groups.
Schuett-Hames et al., 2011; Schuett-Hames & Stewart, 2019b	Clearcut to stream with 30-foot equipment exclusion zone, and 50-foot no-cut buffers	LW, mortality, stand structure, canopy cover	QMD, basal area, tree fall rates, instream LW counts and volume, canopy percentage from densiometer.	1) Substantial variability among sites. 2) Due to scale of study, results only applicable to immediate vicinity of buffer treatment.	10 years post treatment, 50-foot buffer mortality stabilized, cumulative 14.1% reduction in basal area; Reference stands increased in basal area by 2.7% over the 10 years. 10-year cumulative LW recruitment into channels were double that of the reference stands 10-year canopy cover of the 50-foot buffer recovered to similar percentages as the reference stands 10-year cumulative canopy cover of CC was 71.5% due to ingrowth of dense shrubs, saplings and herbaceous plants.

Sobota et al., 2006	Data was collected at 15 riparian sites throughout the Pacific Northwest and the Intermountain West	Tree characteristics, forest structural variables and topographic features	Stand density, basal area, and dominant tree species by basal area; Active channel width and valley floor width.	Bias in landform types between slope categories. Effects of catastrophic disturbance regimes in large rivers not included in model.	The strongest correlations of tree fall direction were with valley constraint. When grouped by species, the individual trees showed a stronger tendency to fall towards the stream when hillslopes were >40%. When field data was integrated into the recruitment model, results showed that stream reaches with steep side slopes (>40%) were 1.5 to 2.4 times more likely to recruit LW into streams than in moderately sloped (< 40%) reaches. The authors warn that while side slope categories (>40%, <40%) was the strongest predictor of tree fall direction in this study, they believe the differences in tree fall direction between these categories mainly characterized differences between fluvial (88% of moderate slope sites) and hillslope landforms (71% of steep slope sites). They suggest that the implications from this study are most applicable to small- to medium-size streams (second- to fourth-order) in mountainous regions where sustained large wood recruitment from riparian forest mortality is the significant management concern.
Teply et al., 2007	25-ft no-cut buffer, with additional 50-foot requiring 88 trees per acre.	Instream wood load	Simulation metrics for forest growth, tree breakage, and in-channel process	The simulation evaluated both a harvest and a no-harvest scenario to predict mean in-stream LW loads after 30, 60, and 100 years	Simulation results predict a 25-foot no-cut buffer, with an additional 50-foot (25–75 feet from the high watermark) zone requiring retention of 88-trees-per-acre were sufficient in maintaining no significant change in in-stream LW loading relative to unharvested reference streams.
Wing & Skaugset, 2002	LW loads and site characteristics were collected from 3793 stream reaches in western Oregon State (west of Cascade crest).	LW pieces, LW key pieces, LW volume	LW abundance, land use history, land ownership, site level attributes	Results presented here are only for forested streams ("tree 3" in text). Landownership was the strongest predictor in some models, but this included multiple areas of unforested reaches.	For in stream LW volume, stream gradient was the most important explanatory variable with the split occurring for stream reaches with gradients less than 4.7% averaging 11.5 m ³ , which was less than half of the average found at higher gradient reaches (25.2 m ³); in this model the stream gradient split explained 11% of the variation observed of instream LW volume. For LW pieces in forested stream reaches, bankfull channel width was the most important explanatory variable with the split occurring for stream channels less than 12.2 m wide. LW pieces for streams <12.2 m wide averaged 11.1 LW pieces per reach while larger channels averaged 4.9 pieces per reach; in this model the BFW split explained 7% of the variation in LW pieces found in forested streams. For key LW pieces (logs at least 0.60 m in diameter and 10 m long) in forested reaches, stream gradient was again the most important explanatory variable with the split occurring at a

					gradient of 4.9%. The streams with a gradient < 4.9% averaged 0.5 key LW pieces per reach while streams with higher gradients averaged 0.9 key LW pieces per reach; in this model stream gradient explained 8% of the variation in key LW pieces found in streams. Lithology caused second, third or fourth level splits after stream gradient or BFW.
--	--	--	--	--	--

3526

3527 Table A-3 List of treatments, variables, metrics, and results from publications reviewed for information on sediment inputs and source.

Reference	Treatment	Variables	Metrics	Notes	Results
Bywater-Reyes et al., 2017	Harvest had a mixture of intensities including clearcut to stream and clearcut with 15 m buffers.	Sediment concentration, basin lithology, geomorphology	Channel, stream, and riparian area characteristics sourced from a mixture of LiDAR and management data.	This study analyzed 6 years of data from the Trask River Watershed in Northeastern Oregon and included data from harvested and unharvested sub-catchments underlain by heterogenous lithologies.	Results from this study indicate that site lithology was a first order control over suspended sediment yield (SSY) with SSY varying by an order of magnitude across lithologies observed. Specifically, SSY was greater in catchments underlain by Siletz Volcanics (r = 0.6), the Trask River Formation (r = 0.4), and landslide deposits. In contrast, the site effect had a strong negative correlation with percent area underlain by diabase (r = 0.7), with the lowest SSY associated with 100% diabase independent of whether earthflow terrain was present. Sites with low SSY and underlain by more resistant lithologies were also resistant to harvest-related increases in SSY. The authors conclude that sites underlain with a friable lithology (e.g., sedimentary formations) had SSYs an order of magnitude higher, on average, following harvest than those on more resistant lithologies (intrusive rocks).
Bywater-Reyes et al., 2018	long-term data (60 years) of sediment, discharge, weather, and disturbance.	Sediment yield, discharge history, physiography.	suspended sediment concentration involved using either vertically integrated storm-based grab samples, or discharge-proportional composite samples.	The authors caution that the high variability of sediment yield over space and time (~0.2 - ~953 t/km ²) indicates that the factors tested in this study should be tested more broadly to investigate their utility to forest managers.	The results of this study show that watershed slope variability combined with cumulative annual discharge explained 67% of the variation in annual sediment yield across the approximately 60-year data set. The results, however, show that annual sediment yields also moderately correlated with many other physiographic variables and the authors caution that the strong relationship with watershed slope variability is likely a proxy for many processes, encompassing multiple catchment For the relationships between disturbance and sediment yield the authors conclude that the few anomalous years of high sediment yield occurred in watersheds with high slope variability and within a decade of forest management and a large flood event.

<p>Hatten et al., 2018</p>	<p>Data from pre restriction and post Oregon BMPs prescriptions for non-fish-bearing<u>non-fish-bearing</u> streams. BMPs: no buffer in non-fish-bearing streams with equipment exclusion zones, and a 15 m no-cut-buffer in fish-bearing streams</p>	<p>suspended sediment concentrations (SSC)</p>	<p>suspended sediment, stream discharge, and daily precipitation</p>	<p>Phase I harvest: 2009 harvest of upper half of watershed. Phase II harvest: 2015 harvest of lower half of watershed.</p>	<p>Methods used in 1966 to harvest the same watershed (no buffer, road construction, broadcast burning) resulted in an approximate 2.8-fold increase in SSC from pre- to post-Harvest. In the contemporary study both the mean and maximum SSC were greater in the reference catchments (FCG and DCG) compared to the harvested catchment (NBLG) across all water years. In NBLG the mean SSC was 32 mg L⁻¹ (~63%) lower after the Phase I harvest and 28.3 mg L⁻¹ (~55%) lower after the Phase II harvest when compared to the pre-harvest concentrations. Compared to the reference watersheds, the mean SSC was 1.5-times greater in FCG (reference) compared to NBLG during the pre-harvest period. After Phase I harvest the mean SSC in FCG was 3.1-times greater and after Phase II harvest was 2.9-times greater when compared to the SSC in the harvested watershed. The authors conclude that contemporary harvesting practices (i.e., stream buffers, smaller harvest units, no broadcast burning, leaving material in channels) were shown to sufficiently mitigate sediment delivery to streams, especially when compared to historic practices.</p>
<p>Karwan et al., 2007</p>	<p>clearcut of the watershed area of by 50%, partial cut of 50% canopy removal, timber road construction Riparian zone harvest followed Idaho FPA rules.</p>	<p>Total suspended solid (TSS) yields</p>	<p>Monthly total suspended solid readings from multiple flume locations for pre-, and post-harvest, and pre- and post-road construction.</p>		<p>A significant and immediate impact of harvest on monthly sediment loads in the clear-cut watershed ($p = 0.00011$), and a marginally significant impact of harvest on monthly sediment loads in the partial-cut ($p = 0.081$) were observed. Total sediment load from the clearcut over the immediate harvest interval (1-year post-harvest) exceeded predicted load by 152%; however, individual monthly loads varied around this amount. The largest increases in percentage and magnitude occurred during snowmelt months, namely April 2002 (560%) and May 2002 (171%). Neither treatment showed a statistical difference in TSS during the recovery time, 2-4 years post-harvest (clearcut: $p = 0.2336$; partial-cut: $p = 0.1739$) compared to the control watersheds. Road construction in both watersheds did not result in statistically significant impacts on monthly sediment loads in either treated watershed during the immediate or recovery time intervals.</p>

Litschert & MacDonald, 2009	Data collected from 4 NF of Nort CA. ~200 harvest sites near riparian zones with 90 m and 45 m buffer widths.	Sediment delivery pathway frequency and characteristics.	Pathway length, width, origins, and connectivity of sediment delivery pathways to streams.	Authors mention a caveat to the results of the study in that there is a potential of underestimating the frequency of rills and sediment plumes as sites recover.	Only 19 of the 200 harvest units had sediment development pathways and only 6 of those were connected to streams and five of those originated from skid trails. Pathway length was significantly related to mean annual precipitation, cosine of the aspect, elevation, and hillslope gradient.
Macdonald et al., 2003a	low-retention = removed all timber >15 cm DBH for pine and > 20 cm DBH for spruce within 20 m of the stream; high-retention = removed all timber > 30 cm within 20 m of the stream.	suspended sediment yields, stream discharge	Discharge rate and total suspended sediments (TSS) collected using Parshall flumes	Only 1-year pre-harvest data was collected to generate predicted TSS and discharge values post-harvest.	Immediately following harvest, TSS concentrations and discharge rates increased above predicted values for both treatment streams. Increased TSS persisted for two-years post-harvest in the high-retention treatment, and for 3-years in the low-retention. This study shows evidence that harvest intensity (low vs. high retention) is proportional to the increase in stream discharge, TSS concentrations, and recovery time to pre-harvest levels. The authors speculate that the treatment areas may have accumulated more snow (e.g., more exposed area below canopy) than in the control reaches leading to the increase in discharge.
McIntyre et al., 2021	1) unharvested reference, 2) 100% treatment, a two-sided 50-ft riparian buffer along the entire RMZ, 3) FP treatment a two-sided 50-ft riparian buffer along at least 50% of the RMZ, (4) 0% treatment, clearcut to	stream discharge, turbidity, and suspended sediment export.		Type N (non-fish-bearing streams). Hard-Rock study.	Discharge increased by 5-7% on average in the 100% treatments while increasing between 26-66% in the FP and 0% treatments. Results for water turbidity and suspended sediment export (SSE) were stochastic in nature and the relationships between SSE export and treatment effects were not strong enough to confidently draw conclusions. The authors conclude that timber harvest did not change the magnitude of sediment export for any buffer treatment.

	stream edge (no-buffer).				
Mueller & Pitlick, 2013	The study used sediment concentration data from 83 drainage basins in Idaho and Wyoming.	Sediment concentration, basin lithology, geomorphology	Sediment concentration distribution, geomorphology, and weather data from multiple sources.		The strongest correlation of in stream sediment supply was with lithology relative softness. Bankfull sediment concentrations increased by as much as 100-fold as basin lithology became dominated by softer sedimentary and volcanic rock. Relief (elevation), basin sideslope, and drainage density showed little correlation strength with bankfull sediment supply.
Puntenney-Desmond et al., 2020	Variable retention buffers with clearcut.	surface and subsurface runoff rates, sediment.	Simulation metrics calibrated with runoff and sediment samples from sample area. Precipitation calibrated for 100-year-rain events.	Differences in sediment yield not statistically significant.	Surface and shallow subsurface runoff rates were greatest in the buffer areas than in the harvested areas or in the harvest-buffer interfaces especially during dry conditions. The authors speculate this was likely due to the greater soil porosity in the disturbed, harvested areas. Sediment concentration in the runoff, however, was approximately 15.8 times higher for the harvested area than in the riparian buffer, and 4.2 times greater than in the harvest-buffer interface. Total sediment yields from the harvested area (runoff + sediment concentration) were approximately 2 times greater than in the buffer areas, and 1.2 times greater in the harvest-buffer interface, however this difference was not significant.
Rachels et al., 2020	harvested following the current Oregon Forest Practices Act policies and BMPs	proportion of sediment from sources	Sediment collected in traps; sourced using chemical analysis	limited sample size (1 treatment, 1 paired reference watershed) and does not incorporate the effects of different watershed physiography on sediment erosion.	The proportion of suspended sediment sources were similar in the harvested (90.3 + 3.4% from stream bank; 7.1 + 3.1% from hillslope) and unharvest (93.1 + 1.8% from streambank; 6.9 + 1.8% from hillslope) watersheds. In the harvested watersheds the sediment mass eroded from the general harvest areas (96.5 + 57.0 g) was approximately 10 times greater than the amount trapped in the riparian buffer (9.1 + 1.9 g), and 4.6 times greater than the amount of sediment collected from the unharvested hillslope (21.0 + 3.3 g).

Safeeq et al., 2020	Long term (51 years) effects of clearcut to stream followed by broadcast burn.	streamflow, sediment transport	Historical streamflow data, precipitation data, sediment grab samples for bedload and suspended sediment.	Data compared one treatment watershed and one control watershed across 51+ years.	The results for post-treatment sediment yields showed suspended load declined to pre-treatment levels in the first two decades following treatment, bedload remained elevated, causing the bedload proportion of the total load to increase through time. Changes in streamflow alone account for 477 Mg/km ² (10%) of the suspended load and 113 Mg/km ² (5%) of the bedload over the post-treatment period. Increase in suspended sediment yield due to increase in sediment supply is 84% of the measured post-treatment total suspended sediment yield. In terms of bedload, 93% of the total measured bedload yield during the posttreatment period can be attributed to an increase in sediment supply. The authors conclude that Following harvest, changes on streamflow alone was estimated in being responsible for < 10% of the resulting suspended sediment transported into streams, while the increase in sediment supply due to harvest disturbance was responsible for >90%.
Wise, 2010	Streamflow patterns derived from instrumental data and from reconstructed tree-ring chronologies were compared with other previously reconstructed rivers in similar climates.	Streamflow	Dendrochronology, historical data records, seasonal patterns	The reconstruction model developed for the analysis explained 62% of the variance in the instrumental record after adjustment for degrees of freedom.	Results showed evidence that droughts of the recent past are not yet as severe, in terms of overall magnitude, as a 30-year extended period of drought discovered in the mid-1600s. However, in terms of number of individual years of < 60% mean-flow (i.e., low-flow years), the period from 1977-2001 were the most severe. Considering the frequency of consecutive drought years, the longest (7-year-droughts), occurred in the early 17th and 18th centuries. However, the 5-year drought period from 2000-2004 was the second driest period over the 415-year period examined.
Wissmar et al., 2004	Data sourced from management records and geospatial data to identify high erosion-risk areas.	Sediment, weather, stand characteristics, landscape factors	unstable soils, immature forests, roads, critical slopes for land failure, and rain-on-snow events		The highest-risk areas contained a combination of all landscape cover factor combinations (rain-on-snow zone, critical failure slope, unstable soil, immature forests, and roaded areas). The lowest risk categories contained only rain-on-snow zones, and critical failure slopes. Roaded areas and unstable soils were only present in risk categories 3-6.

3529 Table A-4. List of treatments, variables, metrics, and results from publications reviewed for information on shade and stream temperature.

Reference	Treatment	Variables	Metrics	Notes	Results
Bladon et al., 2016	15 m buffer with a minimum of ~3.7 m ² conifer basal area retained for every 300 m length of stream). Historical data with no streamside vegetation maintenance (i.e., no buffer) .	Stream temperature	7-day moving mean stream temperature, daily mean stream temperature, and diel stream temperature fluctuation. Data was recorded with Tidbit data loggers.	The authors caution that the streams in this study have potential for a muted stream temperature response following harvest relative to other regions because of the (1) north-south stream orientation (2) steep catchment and channel slopes, (3) potential increases in groundwater contributions after harvesting.	Under the contemporary Oregon Forest Practices Act there was no significant changes in the 7-day moving mean of daily maximum stream temperature, mean daily stream temperature, and diel stream temperature for 3 years following harvest when analyzed across all sites for all summer months (July – September). There was a significant increase in the 7-day moving maximum temperature from pre- to post-harvest values when data was constrained to the period of July 15 – August 15 by 0.6 ± 0.2 °C. However, when analyzed by individually paired sites and when interannual and site variability was accounted for, no significant changes in stream temperature were observed. The authors caution that these results should not be generalized to areas outside the Oregon coast or to riparian areas of different contexts (see notes).
Bladon et al., 2018	Buffer widths at harvested sites varied but averaged 20 m on either side of streams.	Stream temperature, lithology	the 7-day moving average of daily maximum stream temperature adjacent to and downstream of harvest.	Conducted at 3 paired watershed studies on the coast and western Cascades of Oregon. The pre-harvest relationship in stream temperatures for paired sites were used to create predicted changes in stream temperatures post-harvest. Post-harvest stream temperatures exceeding the predictive temperature interval by more than 95% were reported as significant.	Results showed an increase in stream temperatures beyond the 95% predictive interval (PI) at 7 of the 8 sites within harvest areas. 4 of these 7 sites exceeded the PI between 22 and 100% of the time (all summer months for 3 years following harvest). In the remaining 3 sites, exceedance only occurred between 0 and 15% of the time. There was no evidence of elevated stream temperatures beyond the predicted intervals in any of the downstream sites following harvesting. At the harvested sites there was a strong relationship between stream temperature increases and catchment lithologies, but no statistically significant relationship between stream temperature changes and percent of catchment harvested. Downstream sites showed a strong relationship between stream temperatures and the interaction of harvest percentage and lithology. The greatest temperature increases at downstream sites were in areas with a higher percentage of catchment harvested and were underlain by more resistant lithologies. There was no evidence for increases in stream temperatures in catchments with a high percentage of harvest that were underlain by permeable geology

Cole & Newton, 2013	clearcut to stream, partial buffer (12 m width on predominant sun-side)), Oregon state BMP (15-30 m no-cut buffer both sides)	Stream temperature	Controlled for yearly fluctuations in temperatures by analyzing the difference in stream temperature entering and exiting the reach with digital temperature data loggers	Stream temperature data collected for 2 – years prior and 4 to 5 years following harvest. Unharvested control sites were located downstream of treatment sites. Treatment applied to four small fish-bearing streams.	Results showed the most significant increases in daily maximum, and mean, and diel fluctuations in temperatures post-harvest for all no tree buffers. Changes to daily maxima ranged from -0.11 to 3.84 °C, and changes to daily minimum ranged from -1.12 to 0.49 °C. The no tree buffers also showed small but significant changes below predicted summer minima between -1.12 and -0.49 °C. The partial buffer units varied in their response to treatment exhibiting increases, decreases, and no change from preharvest trends.
Cupp & Lofgren, 2014	the “all available shade” rule (ASR), and the standard rule (SR) in eastern WA. ASR: requires retention of all available shade within 75 feet of the stream. SR: some harvest is allowed within the 75-foot buffer depending on elevation and pre-harvest canopy cover.	Canopy closure, shade measurements, stream temperature	Hand-held densiometer (canopy closure), self-leveling fisheye lens digital camera (shade), temperature data loggers	Sites were between 65-100 years old and were situated along second to fourth order streams with harvest-regenerated or fire-regenerated forests. Reference reaches were located upstream from treatment reaches where harvest was applied.	Results showed post-harvest shade values decreased in SR sites (mean effect of -2.8%, p = 0.002), as did the canopy closure values (mean effect of -4.5%, p < 0.001). Shade and canopy closure values did not significantly change in the ASR sites. Mean shade reduction in the SR treatment sites exceeded the mean shade reduction in the ASR sites by 3%. Canopy closure reduction was also greater in the SR sites than in the ASR sites by a mean of 4%. Site seasonal means of daily maximum stream temperature treatment responses in the first two years following harvest ranged from - 0.7 °C to 0.5 °C in the ASR reaches and from -0.3 to 0.6 in the SR reaches. Site seasonal mean post-harvest background responses in reference reaches ranged from - 0.5 °C to 0.6 °C in the first two years following harvest. Mean daily maximum stream temperature increased 0.16 °C in the SR harvest reaches, whereas stream temperatures in both the ASR sites and in the no-harvest reference reaches increased on average by 0.02 °C.
Ehinger et al., 2021	1) TRT: two-sided 50-ft riparian buffer along at least 50% of the RMZ) Buffers encompassing the full width (50 feet), 2) REF: <50ft buffers, 3) Unbuffered, harvested to the edge of the channel, and 4) Reference sites in unharvested	Canopy closure estimated from densitometer, stream water temperature at 30-minute intervals using StowAway TidBit thermistors		Soft Rock study. Multiple Before-After Control-Impact (MBACI) study design. Because of unstable slopes, total buffer area was 18 to 163% greater than a simple 50-ft buffer along 50% of the stream length.	Mean canopy closure decreased in the treatment sites from 97% in the pre-harvest period to 75%, 68%, and 69% in the first, second, and third post-harvest years, respectively, and was related to the proportion of stream buffered and to post-harvest windthrow within the buffer. The seven-day average temperature response increased by 0.6°C, 0.6°C, and 0.3°C in the first, second, and third post-harvest years, respectively. During and after harvest, mean monthly water temperatures were higher, but equaled or exceeded 16.0°C in 2 treatment sites by up to 1.8°C at one site (for 5 years post-harvest)

	forestsUnharvested Reference.				and by 0.1°C at another (at year 5 post-harvest). None of the three REF sites exceeded 16°C during the study.
Gravelle & Link, 2007	50% of the drainage area clearcut to stream edge, thinned to a 50% target shade removal in Fall 2001, and an unimpacted control. Riparian buffer zones were implemented according to Idaho Forest Practices.	stream temperatures at the headwater streams immediately adjacent to treatments, and downstream in larger fish-bearing streams.	Stream temperature data collected from digital sensors.	for the non-fish-bearing, headwater sites pre-treatment data was only collected one season prior to treatment.	In general, the downstream sites showed a cooling effect between -0.2 and -0.3°C. The estimated cooling effect could not be attributed to any cause (e.g., increase in water yield), but the authors conclude that there was no post-harvest increase in peak summer temperatures at the downstream sites. For streams immediately adjacent to the clearcut treatment (headwater streams) a significant increase in temperature was detected at 2 sites ranging between 0.4 and 1.9°C, while a marginally significant decrease in temperature was detected at the third site (-0.1°C, p = 0.06). At the sites located immediately adjacent to partial cuts, results showed mixed results with decreases in temperature (-0.1°C; non-significant) at one site and significant but minimal changes at another site (0.0-3.0°C) across the individual post-harvest years. Overall, there were minimal to no changes in stream peak temperatures following treatment in the partial-cut riparian areas. Despite slight increases in temperature in 2 of the headwater streams, no increase in stream temperature was detected in the larger downstream fish-bearing streams.
Groom et al., 2011a	Private site FPA rules are 15 and 21 m wide on small and medium fish-bearing streams of limited entry. State sites followed a 52 m wide buffer of limited entry. FPA = 6 m no entry buffer, State = 8 m no entry buffer. Thinning intensity not specified.	Stream temperature	Stream temperature collected with digital temperature sensors within harvested areas before and after treatment.	Eighteen of the 33 sites were on privately owned lands, and the other 15 were on state-managed forest land. Treatment reaches were harvested according to the FPA or FMP and included 26 clear-cuts and 7 partial cuts. All private sites were clear-cut. Seventeen sites were	Pre harvest to post harvest comparison of 2 years of data will detect a temperature change of > 0.3°C. Conversely, harvest to state FMP standards resulted in an 8.6% probability of exceedance that did not significantly differ from all other comparisons. The a-priori and secondary post hoc multi-model comparisons did not indicate that timber harvest increased the probability of PCW exceedance at state sites. The authors point out that the 0.3°C change threshold still lies 1 or 2 orders of magnitude lower than previous findings from studies which took place prior to the enactment of the riparian protection standards. Note: PCW criterion is that anthropogenic activities are not permitted to increase

Commented [WB24]: See Section 2-2.2 of the Soft Rock report. Like the Hard Rock treatments these consisted of the entire Type N basin. Buffer types were assigned only in the Riparian Stand and Wood Recruitment chapter.

Commented [bs25R24]: Thank you for the map and clarification.

				harvested along one stream bank, of which 13 were state forest sites. The remaining 16 sites were harvested along both banks.	stream temperature by more than 0.3 °C above its ambient temperature
Groom et al., 2011b	Private site FPA rules are 15 and 21 m wide on small and medium fish-bearing streams with a 6 m no-cut zone immediately adjacent to the stream. Harvesting is allowed in the remaining RMA to a minimum basal area of 10.0 (small streams) and 22.9 (medium streams) m ² /ha. State sites followed a 52 m wide buffer with an 8 m no cut buffer. Limited harvest is allowed within 30 m of the stream only to create mature forest conditions.	Stream temperature, Shade, canopy cover	Stream temperature collected with digital temperature sensors. Stream temperature data was summarized to provide daily minimum, maximum, mean, and fluctuation for analysis. The temperature data was modeled using mixed-effects linear regression. Shade analysis included trees per hectare, basal area per hectare, vegetation plot blowdown, and tree height. a linear regression analysis of shade data (n = 33) was performed.	A comparison of within site changes in maximum temperatures pre-harvest to post-harvest showed an overall increase at private sites, but not all sites behaved the same and some had decreases in maximum temperatures.	Following harvest, maximum temperatures at private sites increased relative to state sites on average by 0.71 °C. Similarly, mean temperatures increased by 0.37 °C (0.24 - 0.50), minimum temperatures by 0.13 °C (0.03 - 0.23), and diel fluctuation increased by 0.58 °C (0.41 - 0.75) relative to state sites. The average of maximum state site temperature changes = 0.0 °C (range = -0.89 to 2.27 °C). Observed maximum temperature changes at private sites averaged 0.73 °C (range = -0.87 to 2.50 °C) and exhibit a greater frequency of post-harvest increases from 0.5 to 2.5 °C compared to state sites. Private site shade values also appeared to decrease pre-harvest to post-harvest. Private post-harvest shade values differed from pre-harvest values (mean change in Shade from 85% to 78%); however, no difference was found for state site shade values pre-harvest to post-harvest (mean change in Shade from 90% to 89%). Results from this study show that between 68% and 75% of variability in post-harvest shade may be accounted for by basal area within 30 m of the stream, tree height, and potentially blow down. The authors speculate that their results suggest sites with shorter trees have higher post-harvest shade and this may be due to the negative correlation between crown ratios and tree heights.
Guenther et al., 2014	Partial retention (50% removal of basal area including riparian zone) methods resulting in approximately 14% reduction in canopy cover on average	Stream temperature, canopy cover, bed temperature	Bed temperatures, stream temperatures, and near stream shallow groundwater temperatures were collected with thermocouples.		Treated watersheds showed an increase of 1.6 - 3.0 °C in daily maximum stream temperatures during the summer months following harvest. Bed temperatures showed an overall increase in temperature but at lower magnitude averaging around 1 °C for up to 30 cm in depth. Bed temperature increases were higher in areas on downwelling flow than in areas of neutral and upwelling flows.

Hunter & Quinn, 2009	an alluvial study site and a bedrock study site whose overall characteristics were otherwise comparable apart from geomorphology.	Stream temperature, Alluvial depth	Water temperature was recorded at 75-m intervals along each channel during the summers of 2003 and 2004	Small sample sizes, results only from two sites for two summers. Actual numeric values not reported but shown in graphs.	Results from this study show consistent differences in stream temperature response in alluvial versus bedrock channels. Seasonal maximum and minimum average daily temperatures varied less at the alluvial site compared to the bedrock site. Two same-day measurements at each site showed the alluvial site gaining 8% of its flow, as compared to the bedrock site whose flow decreased by approximately 15%. Bedrock sites were shown to have the highest variation in reach-scale water temperatures during low flow.
Janisch et al., 2012	clearcut logging with two riparian buffer designs: a continuous buffer and a patched buffered stream. Buffers were 10-15 m wide.	Stream temperature	Channel and catchment attributes (e.g., BFW, Confinement, slope, FPA, etc.), Stream temperatures were recorded with a Tidbit datalogger in areas persistently submerged.	Separation of treatment streams into "clusters" based on year of treatment and an unbalanced experimental design resulted in small sample sizes. Thus, significant differences between treatments were not analyzed. Instead results presented as "significant" represent a significant increase in temperature different from zero.	In general, timber harvest with fixed-width continuous buffers, or patch buffers resulted in increased mean maximum daily summer stream temperatures in the first year following treatment by an average of 1.5 °C (range 0.2 – 3.6 °C). Mean maximum daily summer temperature increases were higher in the streams adjacent to continuous buffer (1.1 °C; range 0.0 to 2.8°C) than the patch buffered catchments (0.6 °C; range – 0.1 to 1.2°C). However, results were highly variable. Post-treatment temperature changes suggested that treatments (p=0.0019), the number of years after treatment (p=0.0090), and the day of the year (p=0.0007) were all significant effects explaining observed changes in temperature. Wetland area (0.96, p<0.01) and length of surface flow (0.67, p=0.05) were strongly correlated with post-logging temperature changes.
Johnson & Jones, 2000	clearcut to stream, patch cutting followed by debris flows (resulted in the removal of all streamside vegetation) , 450+ yo Doug-fir forest reference.	Stream temperature	long term monitoring of weekly stream temperature max, min, and average. Solar radiation data collected from digital sensors. Air and precipitation temperatures collected from local weather stations.	The experimental design used historic stream temperature data to examine changes in stream temperatures. This required conflating data from 2 different devices.	Removal of streamside vegetation whether by clearcut and burn (CCB), or patch-cut and debris (PCD) flow led to significant increases in mean weekly summer maximum and minimum stream temperatures relative to reference streams in the summer immediately following and for 3-4 years post treatment. The CCB's summer mean weekly maximum stream temperatures ranged from 5.4-6.4°C higher than the reference stream for 4 years following treatment. The PCD's summer mean weekly stream temperatures ranged from 3.5-5.2°C higher than the reference stream for 3 years following treatment. The diurnal fluctuations were significantly higher in both treatment streams (6-8 °C in CCB, and 5-6 °C in PCD) relative to reference stream (1-2°C). Pre-harvest temperatures recovered after 15 years of growth.

					Differences in treatment streams and reference stream temperatures were less than 1.1°C pre-treatment and 30-years post-treatment.
Kaylor et al., 2017	50 years post clearcut to streams, control stands were >300 years old	stream light availability, forest age	Stream bank-full width, wetted width, canopy openness, % red alder, and estimated photosynthetically active radiation (PAR) were quantified at 25-m intervals		PAR reaching streams was on average 1.7 times greater in >300-year-old forests than in 30–100-year-old forests. The greatest differences were in streams with both sides harvested. Mean canopy openness was higher in >300-year-old forests (18%) than in 30–100-year-old forests (8.7%). Space-for-time analysis with reviewed literature estimates that canopy closure and minimum light availability occurs at approximately 30 years and maintains until 100 years.
Kibler et al., 2013	Clearcut to stream	Stream temperature, discharge rate,	Stream temperature and discharge rate were recorded with thermistor gauging stations. Canopy cover was recorded with a densiometer as portion of sky covered with vegetation	Post-harvest data was collected only during the summer and autumn immediately following harvest (i.e., 1 season of post-harvest data). Pre-harvest data was collected for 3 years.	Harvest in treatment watersheds resulted in a significant decrease in stream temperatures ranging from –1.9 to -2.8 °C relative to pre-treatment temperatures. The authors attribute the lack of increased temperatures to the shade provided by woody debris.
Macdonald et al., 2003b	Low-retention – remove all timber >15 or >20 cm DBH for pine or spruce, 20 m of the stream 2) high-retention – remove timber >30 cm DBH 20-30m of stream, and 3) Patch-cut removal of all vegetation in the upper 40% of the watershed.	Stream temperature	Temperature data were recorded with Vemco dataloggers. Canopy cover was estimated with densimeters.		Significant increase in stream temperatures ranging from 4 – 6 °C at five years post-harvest, and increased ranges of diurnal temperature fluctuations for all treatment streams relative to the reference streams. Streams that had summer maximum mean weekly temperatures of 8°C before harvesting had maximum temperatures near 12°C or more following harvesting. Daily ranges of 1.0–1.3°C before harvesting became 2.0–3.0°C following harvesting, high-retention buffer treatment mitigated temperature increases for the first three years. Still, increased mortality (attributed to windthrow) caused a reduction in the canopy that, thus, led to increased stream temperatures equivalent to other treatment streams by year five.

<p>McIntyre et al., 2021</p>	<p>(1) unharvested reference, (2) 100% treatment, a two-sided 50-ft riparian buffer along the entire Riparian Management Zone (RMZ), (2) FP treatment a two-sided 50-ft riparian buffer along at least 50% of the RMZ, (3) 0% treatment, clearcut to stream edge (no-buffer).</p>			<p>Hard Rock Study.</p>	<p>Results for canopy cover showed that riparian cover declined after harvest in all buffer treatments reaching a minimum around 4 years post-harvest (after mortality stabilized). The treatments, ranked from least to most change, were REF, 100%, FP, and 0% for all metrics and across all years. Effective shade results showed decreases of 11, 36, and 74 percent in the 100%, FP, and 0% treatments, respectively. Significant post-harvest decreases in shade were noted for all treatments and all years. Results for stream temperature showed that within treatment mean post-pre-harvest difference in the REF treatment never exceeded 1.0°C. In contrast, mean within treatment difference in the 100% treatment was 2.4°C in 2009 (Post-harvest year 1) but never exceeded 1.0°C in later years. The mean difference in the FP treatment exceeded 1.0°C immediately after harvest then again in 2014–2016 (post-harvest years 6–9) while in the 0% treatment the mean difference was 5.3°C initially, then decreased over time to near, but never below, 0.9°C. Stream temperature increased post-harvest at most locations within all 12 harvested sites and remained elevated in the FP and 0% treatments over much of the nine years post-harvest.</p>
<p>Pollock et al., 2009</p>	<p>A range of harvest from 0 – 100%, < 20 years old regrowth, ~ 40 years old regrowth . Unharvested sites were estimated as being >150-years old</p>	<p>Stream temperature, time since harvest, percent of watershed and stream network harvested.</p>	<p>average daily maximum (ADM), average daily range, seasonal range, average, maximum, and minimum Stream temperatures collected with Tidbit data loggers. Stand age grouped by time since harvest.</p>	<p>tested 3 hypotheses: (1) the condition of the riparian forest immediately upstream of a site primarily controls stream temperature, (2) the condition of the entire riparian forest network affects stream temperature, and (3) the forest condition of the entire basin affects stream temperature.</p>	<p>Results of general temperature patterns showed that average daily maximum (ADM) were strongly correlated with average diurnal fluctuations ($r^2 = 0.87$, $p < 0.001$, $n = 40$), indicating that cool streams also had more stable temperatures. For basin-level harvest effects on stream temperatures. The percentage of the basin harvested explained 39% of the variation in the ADM among subbasins ($r^2 = 0.39$, $p < 0.001$, $n = 40$) and 32% of variation in the average daily range (ADR) ($r^2 = 0.32$, $p < 0.001$, $n = 40$). The median ADM for the unharvested subbasins was 12.8 °C (mean = 12.1 °C), which was significantly lower than 14.5 °C, the median (and average) ADM for the harvested subbasins ($p < 0.001$). Likewise, the median (and average) ADR for the unharvested subbasins was 0.9 °C, which was significantly lower than 1.6 °C, the median ADR (average = 1.7 °C) for the harvested subbasins ($p < 0.001$). Results for the correlations between the riparian network scale forest harvest and stream temperature showed that the total</p>

					<p>percentage of the riparian forest network upstream of temperature loggers harvested explained 33% of the variation in the ADM among subbasins ($r^2 = 0.33$, $p < 0.001$, $n = 40$) and 20% of variation in the ADR ($r^2 = 0.20$, $p = 0.003$, $n = 40$). However, the total percentage of upstream riparian forest harvested within the last 20 years was not significantly correlated to ADM or ADR. Results for near upstream riparian harvest and stream temperature showed either non-significant, or very weakly significant correlations. For example, there were no significant correlations between the percentage of near upstream riparian forest recently clear-cut and ADM temperature ($r^2 = 0.03$, $p = 0.79$, $n = 40$), the ADR of stream temperatures ($r^2 = 0.02$, $p = 0.61$, $n = 40$) or any other stream temperature parameters. The proportion of total harvested near upstream riparian forest (avg = 0.66, SD \pm 0.34, range = 0.0-1.0) was weakly correlated with ADM ($r^2 = 0.12$, $p = 0.02$, $n = 40$) and not significantly correlated with ADR ($r^2 = 0.07$, $p = 0.06$, $n = 40$). Even when the upstream riparian corridor length was shortened to 400 m and then to 200 m, and the definition of recently harvested was narrowed to <10 year, no significant relationships between temperature and the condition of the near upstream riparian forest was found. for these models, the percentage of basin area harvested was the best predictor of variation in mean maximum stream temperatures. The probability of stream temperatures increasing beyond DOE standards (16 °C for seven-day average of maximum temperatures) increased with percent harvest. Nine of the 18 sites with 50-75% harvest and seven of the nine sites with >75% harvest failed to meet these standards. The authors interpret these results as evidence that the total amount of forest harvested within a basin, and within a riparian stream network are the most important predictors of changes in summer stream temperatures. They conclude that watersheds with 25-100% of their total area harvested had higher stream temperatures than watersheds with little or no harvest.</p>
--	--	--	--	--	--

Reiter et al., 2020	Clearcut, no buffer (CC_NB), clearcut with 10-m no cut buffer (CC_B), thinning with 10 m no-cut buffer (TH_B), and unharvested reference (REF) streams.	Stream temperature	Temperature data was separated into 5 th , 25 th , 50 th , 75 th , and 95 th percentiles. the researchers also quantified the percentage of summer where temperatures were above 16 and 15 °C.	Sample sizes are relatively low for some treatments. (CC_NB; n = 4); (CC_B; n = 3); (TH_B; n =1); (REF; n = 7).	A 10 m buffer was sufficient in maintaining summer temperature changes compared to reference streams regardless of upland treatment (clear-cut, thinning). Unbuffered streams (Clear-cut to streams) showed significant increases in stream temperatures with an average of 3.6 °C (SE = 0.4) increase relative to reference streams. Unbuffered streams spent 1.3% and 4.7% of the recorded time above 16 °C and 15 °C respectively (habitat temperature thresholds for two local amphibian larvae, coastal tailed frog, coastal giant salamander). The authors conclude that while significant changes in mean and percentile changes in temperature were observed, the amount of time spent above critical temperature thresholds for important amphibian species was minimal.
Reiter et al., 2015	. Various buffer prescriptions changed over time. (mid1970s – 1980s = “nominal”; mid 1980s – mid 1990s = 23 m; 2001 – 2009 = 30 m buffers)	Stream temperature data from four permanent sampling stations in the Deschutes River Watershed from 1975- 2009. Results for this analysis are for 3 watersheds (1-large, 1-medium, 1-small)	Long term stream and air temperature collected from sampling stations. To detect correlations of stream and air temperature change with land management activity separately from climate changes the data was fit to a model that included the effects of climate.	Methods for stream temperature data collection varied at different periods resulting in a margin of error for monthly temperatures of 0.14°C for 1975 - 1983, 0.09°C for 1984 – 1999, and 0.02°C. for 2000 – 2009.	Results for trends in stream temperature over the 35-year study period without adjustment for climate change showed no statistically significant trend in water temperature changes for the large watershed, while the medium watershed (Thurston Creek) showed decreasing trends in TMAX_WAT for June, July, and August, ranging in magnitude from 0.05°C (August) to 0.08°C (July) per year. For the smaller watershed, Hard Creek (Ware Creek was not included in this analysis), had significant decreasing trends in TMAX_WAT for July, August, and September. The magnitude of these trends was yearly decreases of TMAX_WAT by 0.05, 0.08, and 0.05°C, for July, August, and September, respectively. Significant changes in trends for TMIN_WAT were only found for the large basin site with yearly increases of 0.04, 0.03, and 0.04°C for July, August, and September, respectively. Results for stream temperature trends after adjusting for changes in air temperature (climate) showed significant decreasing trends in TMAX_WAT for the large basin by 0.04, 0.03, and 0.04°C yearly, for July, August, and September, respectively. For the medium basin, trends showed yearly decreases in TMAX_WAT of 0.07, 0.08, 0.06, and 0.03 for June, July, August, and September, respectively. For the small basin, climate adjusted trends in TMAX_WAT showed significant decreases in yearly trends by 0.05, 0.08, and 0.05 for July, August, and September, respectively. When stream temperature was examined with its correlation with estimated annual

					shade recovery from initial harvest (indexed by ACD). Significant correlations were found for monthly temperature metrics that were adjusted for climate, for all basins. The authors conclude that the results of this study show evidence that implementation of protection buffers in this area were sufficient in maintaining stream temperatures. Conversely, this study also shows evidence that despite these protections from land management induced stream temperature changes, these protections have been somewhat offset by the warming climate conditions.
Roon et al., 2021a	Thinning treatments resulting in a mean shade reduction of <5% (-8.0 - -0.5) at one watershed and 23.0% at two watersheds (-25.8, -20.1)	Stream temperature, solar radiation, Shade	Stream temperature was collected using digital sensors; solar radiation was measured using silicon pyranometers; riparian shade was measured using hemispherical photography.	Only 1-year pre- and post-treatment data. Site selection and replication was not random and thus may not be applicable outside of the northern California redwood forests.	No significant changes in stream temperatures were detected in the low-intensity thinning treatment watersheds. For the higher intensity thinning treatments. Maximum weekly average of the maximum temperatures increased during spring by a mean of 1.7 °C (95% CI: 0.9, 2.5), summer by a mean of 2.8 °C (1.8, 3.8), and fall by a mean of 1.0 °C (0.5, 1.5) and increased in downstream reaches during spring by a mean of 1.0° C (0.0, 2.0) and summer by a mean of 1.4° C (0.3, 2.6). Thermal variability of streams were most pronounced during summer increasing the daily range by a mean of 2.5° C (95% CI: 1.6, 3.4) and variance by a mean of 1.6°C (0.7, 2.5), but also increased during spring (daily range: 0.5 °C; variance: 0.3° C) and fall (daily range: 0.4° C; variance: 0.1 °C). Increases in thermal variability in downstream reaches were limited to summer (daily range: 0.7° C; variance: 0.5 °C). The authors interpret their results as evidence that that changes in shade of 5% or less caused minimal changes in temperature while reductions in shade of 20–30% resulted in much larger increases in temperature.
Roon et al., 2021b	Effective shade reductions ranging between 19-30% along 200 m reach, or 4-5% along 100 m reach.	local and downstream temperature	Stream temperature collected with digital temperature sensors within harvest area and every 200 m downstream of stream network.	Stream temperature data was only collected for one-year pre- and one-year post-harvest.	In the reaches with higher reductions in shade (19-30%) there was accumulation of 45° to 115°C additional degree days from pre- to post treatment years, while the reaches with lower reductions in shade (4-5%) only accumulated 10° to 15°C additional degree days. Travel distance of increased stream temperatures also appeared to be dependent on thinning intensity. The lower shade reduction reaches had an increased temperature effect downstream with travel distance of 75-150 m, while the high shade reduction sites had a downstream travel

					distance of 300- ~1000 m. In the high shade reduction sites, treatment reaches that were further apart (> 400 m) showed dissipation in increased stream temperatures downstream, while in parts of the stream where treatments were <400 m apart, temperature increases did not always dissipate before entering another the next treatment reach.
Sugden et al., 2019	Montana state law : 15.2 m wide buffers no more than half the trees greater than 204 mm (8 in) diameter at breast height (DBH). In no case, however, can stocking levels of leave trees be reduced to less than 217 trees per hectare. .	Stream temperature, fish population, Canopy cover	Daily max, min, and average stream temperatures collected with data loggers during summer months. The fish community was inventoried 100 m reaches using an electro-fishing pass of capture method. Canopy cover was estimated using a combination of simulation modeling and using a concave spherical densiometer.	Data only collected for one year pre-harvest and one year post-harvest.	The mean basal area (BA) declined from 30.2 m ² /ha pre-harvest to 26.4 m ² /ha post-harvest (mean = -13%, range from -32% to 0%). Windthrow further reduced the mean BA to 25.9 m ² /ha (mean = -2%, range = -32% -0%). Change in mean canopy cover were not significant based on the simulation modeling (-3%), or densiometer readings (+1%). Results of the model for the effect of harvest on stream temperature showed no detectable increase in treatment streams relative to control streams. The estimated mean site level response in maximum weekly maximum temperatures (MWMWT) varied from - 2.1 °C to +3.3 °C. Overall, 20 of 30 sites had estimated site level response within ±0.5 °C. There were five sites that had an estimated site-level response greater than 0.5 °C (i.e. warming) and five sites that had an estimated site level response less than -0.5 °C (i.e. cooling). Results for the fish population showed approximately 7% increase in trout population from pre-harvest to post-harvest, but this difference was not significant.
Swartz et al., 2020	In the experimental reaches 30 m gaps were created, centered on a tree next to the stream and at least 30 m in from the beginning of the reach. Actual gap sizes varied across sites from approximately 514 m ² to 1,374 m ² with a mean of 962 m ² .	Stream temperature, Light reaching stream, canopy cover	Riparian shade-hemispherical photos. Light reaching the stream- photodegradation of fluorescent dyes. Stream temperature - HOBO sensors for seven-day moving average of mean and maximum temperatures.	Data was collected for one year pre-harvest, during harvest year (harvest took place in late fall 2017), and one-year post-harvest.	Results showed that after gaps were cut, the BACI analysis showed strong evidence for significant increase in mean reach light (p < 0.01) to a mean of 3.91 (SD ± 1.63) moles of photons m ⁻² day ⁻¹ , overall resulting in a mean change in light of 2.93 (SD ± 1.50) moles of photons m ⁻² day ⁻¹ . Through the entirety of the treatment reach mean shading declined by only 4% (SD ± 0.02%). Overall, the gap treatments did not change summer T 7DayMax or T 7DayMean significantly across the 6 study sites. However, reaches showed a statistically significant effect of the gap for average daily maximums (p < 0.01) and for average daily means (p = 0.02). The regression comparison reveals there will be on average an additional 0.12 °C/°C increase in daily maximum temperature in the reach with a gap.

					Likewise, for the daily mean, for every degree increase in the shaded reference reach, an average additional increase of 0.05 °C in a reach with a small gap is expected. The regression comparison reveals there will be on average an additional 0.12 °C/°C increase in daily maximum temperature in the reach with a gap. Likewise, for the daily mean, for every degree increase in the shaded reference reach, an average additional increase of 0.05 °C in a reach with a small gap is expected.
Warren et al., 2013	Old-growth forests were estimated to be over 500 years old, and mature second growth forests were estimated to be between 31 and 59 years old.	Light reaching bottom of stream, canopy cover	The percent of canopy cover was estimated using a densiometer, the amount of light reaching the bottom of the stream was estimated using a fluorescent dye that degrades overtime from light exposure	Relatively small sample sizes (n = 4). Significant differences were only found in 3 of the four paired reaches.	Results showed that the differences in stream light availability and percent forest cover between old-growth and second-growth reaches were significant in both south-facing watersheds in mid-summer at an alpha of 0.01 for the dye results and 0.10 for the cover results. For the north-facing watersheds differences in canopy cover and light availability (alpha = 0.01, and 0.10 respectively) were only significant at 1 of the two reaches. Overall, three of the four paired old-growth reaches had significantly lower mean percent canopy cover, and significantly higher mean decline in fluorescent dye concentrations The authors interpret these results as evidence that old-growth forest canopies were more complex and had more frequent gaps allowing for more light availability and lower mean canopy cover, on average, than in adjacent mature second- growth forests.

3531

3532 **Appendix II**

3533

3534 **Shade and LW**

3535

3536 Anderson & Meleason, 2009

3537

3538 Anderson, P.D., Meleason, M.A., 2009. Discerning responses of down wood and understory
3539 vegetation abundance to riparian buffer width and thinning treatments: an equivalence-
3540 inequivalence approach. *Can. J. For. Res.* 39, 2470–2485 <https://doi.org/10.1139/X09-151>

3541

3542 The purpose of this study was to determine the effect of buffer width on understory vegetation
3543 and down woody responses both within the unthinned buffer and in the adjacent thinned stand. A
3544 secondary objective of this study was to explore the ability of equivalence-nonequivalence
3545 statistical tests at assessing the degree of similarity between stands. The focus of this study was
3546 on second-growth stands dominated by Douglas-fir at multiple sites along the coast and Cascade
3547 Range in western Oregon. Six combinations of buffer width and upslope density management
3548 prescription were evaluated: one site potential tree height buffer averaging 69 m adjacent to
3549 thinning and a 0.4 patch opening; variable width buffer averaging 22 m adjacent to thinning and
3550 a 0.4 patch opening; streamside retention width averaging 9 m adjacent to thinning; and an
3551 unthinned stand serving as a reference. Pearson correlation and multivariate analysis of variation
3552 were used to examine data on percent cover of small and large down wood, and percent cover of
3553 shrubs, herbs, and moss. Inferences on buffer performance were generated using linear mixed
3554 model analysis, equivalence-inequivalence tests, and two post-hoc comparisons. The results from
3555 this study show upland thinning led only to subtle changes in understory vegetation cover and
3556 composition with vegetation responses most prevalent with narrow buffer widths and particularly
3557 when adjacent to patch openings. There was a lack of significant change in down wood response
3558 to treatments.

3559

3560 **Shade**

3561

3562 Anderson et al., 2007

3563

3564 Anderson, P.D., Larson, D.J., Chan, S.S., 2007. Riparian buffer and density management
3565 influences on microclimate of young headwater forests of western Oregon. *For. Sci.* 53, 254–
3566 269. <https://doi.org/10.1093/forestscience/53.2.254>

3567

3568 The purpose of this study was to characterize variation in overstory density, canopy closure, and
3569 microclimate as a function of distance from headwater streams, and (2) determine differences in the
3570 ability of thinned stands and unthinned stands to maintain understory microclimate above the stream
3571 channel and in the riparian zone. The focus of this study was on second-growth (30- to 80-year-old)
3572 Douglas-fir forests characteristic of western Oregon. The study was located at four sites along the
3573 Oregon coast and at one site on the western Oregon Cascade Range. Stands were either thinned to
3574 approximately 198 trees per acre (TPA) or were left unthinned and ranged from 500-865 TPA. Within
3575 thinned stands, 10% of the area was harvested to create patch openings and 10% was left as clusters of
3576 “leave islands”. Streams within treated stands were surrounded by buffers of either (1) one site-potential
3577 tree averaging 69 m (B1), (2) variable width buffer averaging 22 m (VB), or (3) streamside retention
3578 buffer averaging 9 m (SR-T). These six combinations of buffer width and adjacent density management
3579 were evaluated using univariate linear modeling and compared with untreated (UT) stands. Microsite
3580 and microclimate data were obtained through repeated transect measurements extending laterally from
3581 stream center and into the riparian zone and upland treated stand 2-5 years after treatment. The stand
3582 basal area was determined through variable radius plot sampling. Canopy cover was estimated through
3583 photographic techniques during the summer leaf-on period. The results from this study show that the
3584 ability of narrow streamside buffers (SR-T) at moderating stream microclimate in treated stands was
3585 questionable. Visible sky at stream center only differed significantly between SR-T (9.6%) and UT
3586 (4.2%) stands. The SR-T stands showed a +4.5°C difference in daily maximum temperatures just above
3587 stream center when compared to the UT stands. However, this difference was not statistically significant.
3588 The researchers report that SR-T had a weak temperature gradient (tested at 0-10 m and 10-30 m
3589 increments from stream center) indicating the stream center and buffer microclimates were nearly the
3590 same as upslope in the thinned stand. Within the riparian buffer zones daily maximum temperatures
3591 were higher in all treated stands when compared to UT stands. The differences in daily maximum
3592 temperatures ranged from 1.1°C (B1) to 4.0°C (SR-T), but the difference was only significant in one
3593 SR-T stand. The maximum air temperature within buffer zones adjacent to patch openings was 3.5°C
3594 higher than in UT stands. Soil temperature changes were only evident within patch openings ranging
3595 from 3.6 - 8.8°C higher than in UT stands. The researchers of this study conclude by saying that buffers
3596 with widths defined by the transition of riparian to upslope vegetation or significant topographic slope
3597 breaks appear sufficient at mitigating effects from upslope harvests on the above-stream microclimate.
3598 Their suggestions for further study center around cross-disciplinary research into the relationships
3599 between forest structure, microclimate, and habitat suitability on headwater riparian organisms.

3600

3601 **Stream Temperatures**

3602

3603 Cole & Newton, 2013

3604

3605 Cole, E., & Newton, M. (2013). Influence of streamside buffers on stream temperature response
3606 following clear-cut harvesting in western Oregon. *Canadian journal of forest research*, 43(11), 993-1005.
3607 <https://doi.org/10.1139/cjfr-2013-0138>

3608

3609 This study compares the changes in stream temperatures following a clearcut with three different buffer
3610 treatments – no tree buffer, predominantly sun-sided 12 m wide partial buffer, and a two-sided 15-30 m
3611 buffer (BMP for this area). The study was conducted on four small ~~fish-bearing~~ streams in
3612 the area surrounding Corvallis, Oregon. Streams were dominated by both hardwood and conifers and
3613 were located at low- and mid-elevations. Each treatment alternated with unharvested reference sections
3614 along study reaches spanning 1800-2600 meters. Stream temperature data adjacent to treatment and
3615 downstream of treatment were collected for 2 –years prior and 4 to 5 years following harvest. Time-
3616 series regression analysis was used to evaluate the change in temperatures between pre- and post-
3617 harvest. The researchers controlled for yearly fluctuations in temperatures by analyzing the difference in
3618 stream temperature entering and exiting the experimental reaches. Results showed significant increases
3619 in daily maximum, mean, and diel fluctuations in temperatures post-harvest for all no tree buffers (up to
3620 3.8 °C). The no tree buffers also showed small but significant changes below predicted summer minima
3621 by as much as 1.2°C. The partial buffer units varied in their response to treatment exhibiting increases,
3622 decreases, and no change from preharvest trends. For example, at one site, there were no detectable
3623 changes in means, minima, or diel fluctuations but significantly lower maximum temperatures post-
3624 harvest ($p = 0.0021$; actual temperatures not reported). Partial buffers at another site reported lower
3625 trends in mean, maxima, and diel fluctuations in temperature post-harvest, and no difference in minima.
3626 Only one partial buffer site showed increases in all recorded trends (mean, minima, maxima, diel
3627 fluctuations). The BMP buffered treatment sites also showed variation in results. One site showed no
3628 detectable changes, one site showed small but significant ($p < 0.0350$; actual temperatures not reported)
3629 decreases in downstream temperatures. Only two BMP buffered sites showed significant ($p < 0.0499$)
3630 increases in mean, maxima, and diel fluctuations in temperatures. The highest increase in maxima for
3631 any BMP buffered site was 5.3°C. Changes in temperature trends in uncut reference post-treatment were
3632 minimal and attributed to downstream effects from the treatment reaches. However, when post-harvest
3633 trends in upstream treated sites were higher than pre-harvest temperatures tended to fall below pre-
3634 harvest values when passing through the unharvested downstream units. For within-unit trends,
3635 unharvested units downstream from no tree and partial buffers showed trends of significantly decreasing
3636 daily maximum temperatures. When the data was analyzed by 7-day moving mean maximum
3637 temperatures, the no tree buffers showed significant increases after harvest. The authors report that most
3638 partial and BMP buffers resulted in minimal increases or negligible changes to the 7-day moving mean
3639 maximum temperatures (actual values not reported). Significant changes in one or more temperature
3640 trends (mean, minima, maxima, diel fluctuations) were detected in all treatment stream post-harvest with
3641 only one exception at a BMP buffered site This was a well planned and executed experimental design
3642 that shows how changes in stream temperatures post-harvest are directly related to residual buffer
3643 treatment while also showing evidence that many other factors such as stream features (orientation,

3644 topography, ground water source) can compound or ameliorate these effects (I.e., changes in temperature
3645 were highly affected by site factors).

3646

3647 **Stream Temperature**

3648

3649 Johnson & Jones, 2000

3650

3651 Johnson, S. L., & Jones, J. A. (2000). Stream temperature responses to forest harvest and debris flows in
3652 western Cascades, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences*, 57(S2), 30-39.
3653 <https://doi.org/10.1139/f00-109>

3654

3655 This paper is a study of the changes in mean stream temperature minimum, maximum, diurnal
3656 fluctuation, and interannual and seasonal variability following harvest in three small basins of the
3657 H.J. Andrews experimental watershed between 1962 and 1966. The experimental design used
3658 historic stream temperature data to examine changes in stream temperature following clear-cut
3659 (no buffer) and burning in one watershed; patch cutting and debris flows (resulted in the removal
3660 of all streamside vegetation 3 years after cut) treatments in another watershed; and one old-
3661 growth uncut reference watershed. All watersheds were dominated by 450-year-old Doug-fir
3662 forests prior to harvest. Data was analyzed for the period 1959-1997. Mean weekly temperature
3663 maximum, minimum, and annual fluctuations were compared between all three watersheds using
3664 a complete factor analysis of variance (ANOVA). The experiment also involved long-term
3665 monitoring to evaluate time until recovery of pre-treatment temperature fluctuations. Results
3666 showed a significant increase in stream temperatures in both treatment watersheds after treatment
3667 compared to the unharvested site. The unharvested watershed showed higher interannual
3668 variability in maximum stream temperatures ranging from 15 to 19°C. The two treatment
3669 watersheds, despite differences in disturbances, (clear-cut and burn vs. Patch cut and debris-
3670 flow) followed similar trajectories from 1966-1982. Stream temperature summer maximums
3671 reached 23.9°C and 21.7°C 1-2 years post-harvest (clear-cut/burn and patch-cut/debris flow
3672 respectively) and returned to pre-harvest summer temperatures by 1980 (~15 years post-harvest).
3673 Both treatment watersheds exhibited significant increases in mean weekly minimum and
3674 maximum stream temperatures in the summer months immediately following harvest and for at
3675 least 3 years compared to the unharvested reference. The clear-cut and burn watershed's
3676 weekly maximum summer temperatures ranged between 5.4 and 6.4°C higher, and mean weekly
3677 minimum ranged 1.6-2.0°C higher than the reference streams for 4 years post-harvest. The patch-
3678 cut and debris-flow watershed exhibited mean weekly maximum stream temperatures 3.5-5.2°C
3679 higher than in the reference stream for 3 years following harvest/disturbance. Prior to harvest and
3680 30 years post-harvest the mean weekly maximum and minimum stream temperatures for both
3681 treatment streams differed less than 1.1°C from the reference stream. These differences in stream
3682 temperatures from treated and untreated sites were amplified during periods of high solar inputs

3683 and reduced during periods of cloud cover. Differences in stream temperatures were greatest
3684 during the end of July and beginning of June. Diurnal fluctuations in stream temperatures were
3685 also significantly higher in both treatment watersheds (6-8 °C in the clearcut, and 5-6 °C in the
3686 patch-cut) relative to the reference stream (1-2 °C). Stream temperatures returned to pre-harvest
3687 levels after 15 years of growth.

3688

3689 **Large Wood (LW)**

3690

3691 Bahuguna et al., 2010

3692

3693 Bahuguna, D., Mitchell, S.J., Miquelajauregui, Y., 2010. Windthrow and recruitment of large woody
3694 debris in riparian stands. *Forest Ecology and Management* 259, 2048–2055.
3695 <https://doi.org/10.1016/j.foreco.2010.02.015>

3696

3697 The purpose of this paper was to evaluate the effect of riparian buffer width on windthrow and LW
3698 recruitment and to contrast data with unharvested controls. This paper also seeks to document the
3699 geometry of post-harvest windthrow from buffers of varying widths and to develop a model framework
3700 for incorporating supply of LW originating from windthrow to streams from riparian buffers. The focus
3701 of this paper is on dense young conifer-dominated forests originating from harvest followed by wildfire.
3702 This study is located in the Coast Mountains, approximately 60 km east of Vancouver, BC. Two buffer
3703 widths on each side of the stream (10 m and 30 m) along with an unharvested control were each
3704 replicated three times in the experiment. The researchers used a strip plot sampling method running
3705 parallel to the stream to collect data on species, diameter, height, and status (standing live/dead)
3706 beginning in the year prior to harvest and annually thereafter for seven years. A General Linear Model
3707 Procedure was used to determine the significance of variables. The Pearson correlation coefficient was
3708 used to assess correlations and potential predictor variables. Multiple linear regression was then used to
3709 determine the utility of the variables at determining LW height above the stream. Following harvest,
3710 11% of initially standing timber was blown down in the first and second years in the 10 m buffer,
3711 compared to 4% in the 30 m buffer, and 1% in the unharvested controls. Following 8 years post-harvest,
3712 a significant amount of annual mortality occurred in the unharvested control at 30%, compared to 15%
3713 in both 30 m and 10 m buffers. 77% of LW was in the 10 cm - 20 cm diameter class while the mean
3714 diameter of standing trees in riparian buffers was 30 cm indicating small diameter trees were
3715 significantly more represented in streams. Only 3% of windthrown logs fell perpendicular to the stream
3716 with the majority falling diagonal-perpendicular relative to the stream. The researchers of this study
3717 conclude that recruitment of logs into streams lags behind the post-harvest pulse of windthrow by
3718 several years. The lag depends on the size, species, and condition of logs, and their direction of fall
3719 relative to stream valley geometry.

3720

3721 **Species Richness**

3722

3723 Baldwin et al., 2012 (Removed from focal list)

3724

3725 Baldwin, L.K., Petersen, C.L., Bradfield, G.E., Jones, W.M., Black, S.T., Karakatsoulis, J., 2012.

3726 Bryophyte response to forest canopy treatments within the riparian zone of high-elevation small streams.

3727 Can. J. For. Res. 42, 141–156. <https://doi.org/10.1139/x11-165>

3728

3729 The purpose of this study was to examine the influence of forest harvesting practices and distance from
3730 the stream on riparian-bryophyte communities. The experiment was limited to the montane spruce forest
3731 type which is considered moderately open and dominated by lodgepole pine in the uplands and by
3732 hybrid spruce in well-developed riparian areas. The study took place at five different watersheds located
3733 approximately 70 km from Kamloops, BC. Three primary treatments: clear-cut (n=7), two-sided buffer
3734 averaging approximately 15 m on both sides (n=10), and a continuous forest (n=6) were used to sample
3735 numerous environmental variables including elevation, aspect, slope, buffer width, and CWD decay
3736 class. Bryophytes (classified into life history strategies), stand structure, and microhabitat were also
3737 measured 1, 5, and 10 m from the streams edge. Additionally, the DBH of all conifer stems as well as
3738 percent vegetation cover were measured along transects. All data were collected in July-August of 2007
3739 and 2008. Minimum time since disturbance for clearcut sites was 13 years versus a minimum of 5 years
3740 in buffered sites. An analysis of variance was used to compare environmental, stream, and stand
3741 structure characteristics among canopy treatments. Mean values were calculated for stand structure and
3742 substrate variables recording in transects. Bryophytes were analyzed within functional groups based on
3743 growth form, substrate affiliations, and life history. Linear models were used to evaluate the effects of
3744 distance to stream, forest canopy treatment, and their interaction on response variables. Overall CWD
3745 did not differ significantly among treatments, although buffer treatment sites had significantly higher
3746 volume of CWD in early decay classes compared to clearcut and continuous forests. The researchers
3747 suggest the early decay class CWD in buffer treated sites was likely the result of increased stem
3748 breakage. After accounting for distance from the stream, the richness and frequency of bryophyte
3749 functional communities was intermediate to continuous and clearcut sites. Compared to continuous sites,
3750 buffered sites featured significantly lower richness and frequency of many forest-associated groups.
3751 Furthermore, buffered sites also did not support increased richness or frequency of disturbance-
3752 associated species. Clearcut treatments featured higher levels of disturbance associated species including
3753 colonists, canopy species, and species typically found on mineral soil. Data from this study also showed
3754 bryophyte species richness and frequency decline with increasing distance from the stream. The authors
3755 conclude by noting that while bryophyte communities in buffered sites are significantly more diverse
3756 than communities in clearcut sites, reductions in forest-associated species as well as in the bryophyte
3757 mat as a result of large-scale forestry indicate that the ecological function of buffer-dwelling bryophyte
3758 communities may be hindered and could benefit alongside large uncut forest reserves.

3759

3760 **Sediment**

3761

3762 Mueller & Pitlick, 2013

3763

3764 Mueller, E. R., & Pitlick, J. (2013). Sediment supply and channel morphology in mountain river
3765 systems: 1. Relative importance of lithology, topography, and climate. *Journal of Geophysical*
3766 *Research: Earth Surface*, 118(4), 2325-2342. <https://doi.org/10.1002/2013JF002843>

3767

3768 This study used correlation analysis to assess the relative impact of lithology, basin relief, mean basin
3769 slope, and drainage density on in stream sediment supply defined by the bankfull sediment concentration
3770 (bedload and suspended load). The study used sediment concentration data from 83 drainage basins in
3771 Idaho and Wyoming. Lithologies of the study area were divided into four categories ranging from
3772 hardest to softest- granitic, metasedimentary, volcanic, and sedimentary. The results showed the
3773 strongest correlation of bankfull sediment concentration was with basin lithology, and showed little
3774 correlation strength with slope, relief and drainage density. As lithologies become dominated by softer
3775 parent materials (volcanic and sedimentary rocks), bankfull sediment concentrations increased by as
3776 much as 100-fold. These results suggest that lithology can be more important in estimating sediment
3777 supply than topography. The authors discuss using a correlative analysis but give little description of
3778 what that analysis was or how they compare the values of each correlation strength to see if the
3779 differences were significant.

3780

3781 **CWD Modeling**

3782

3783 Benda et al., 2016

3784

3785 Benda, L.E., Litschert, S.E., Reeves, G., Pabst, R., 2016. Thinning and in-stream wood
3786 recruitment in riparian second growth forests in coastal Oregon and the use of buffers and tree
3787 tipping as mitigation. *J. For. Res.* 27, 821–836. <https://doi.org/10.1007/s11676-015-0173-2>

3788

3789 The purpose of this study was to develop a model which examines the effects of riparian thinning
3790 on in-stream wood recruitment in second growth stands. A secondary objective of this study was
3791 to model how manual felling of trees in no-harvest buffer zones impacts the effects of thinning.
3792 The study site was located within the Alcea watershed in central coastal Oregon. Silvicultural
3793 simulation treatments used the reach scale wood model (RSWM) and included: (1) no harvest
3794 control; (2) single entry thinning from below (thinning from below removes the smallest trees to
3795 simulate suppression mortality) with and without a 10 m width no-cut buffers; (3) double entry

3796 thinning from below with the second thinning occurring 25 years after the first with and without
3797 10 m no-cut buffers (4) Each thinning treatment was also combined with some mechanical
3798 introduction of thinned trees into the stream encompassing a range between 5 and 20 % of the
3799 thinned trees. . The simulation model RSWM was run for 100 years in 5-year time steps. In the
3800 no-harvest control, the model output shows the density of live trees declines from 687 trees-per-
3801 hectare (tph) in 2015 to 266 tph in 2110 due to natural suppression mortality (-61 % from initial
3802 conditions). The single-entry thin reduces stand density to 225 tph in 2015 (-67 %) and declines
3803 further to 160 tph by 2110 (-77 %). The double entry thinning resulted in 123 tph after the
3804 second thinning in 2040 (-82%) and maintained that density until 2110. Both thinning treatments
3805 resulted in a substantial reduction of dead trees that could contribute to in-stream wood over
3806 time. The model output for single entry thinning treatments predicts a 33% or 66% reduction of
3807 in-stream wood over a century relative to the unharvested reference for harvest on one side or
3808 both sides of the stream, respectively. Adding the 10-m no cut buffer reduced total loss to 7 and
3809 14%. Including mechanical tipping of 5,10,15, and 20% of cut stems without a buffer in the
3810 single entry thinning treatment changes the relative in-stream percentages of wood relative to the
3811 reference stream to -15, -6, +1, and +6%, respectively. To completely offset the loss of in stream
3812 wood due to single entry thinning mechanical tipping of 14 and 12% were required without and
3813 with buffers. Double entry thinning treatments without a buffer predicted further reduction in
3814 wood recruitment over a century of simulation with 42 and 84% reduction of in stream wood
3815 relative to the reference stream when one side and both sides of the channel were harvested.
3816 Adding a 10 m buffer reduced total reduction of in stream wood to 11 and 22% for thinning on
3817 one and both sides of the channel. To offset the predicted changes of in stream wood volume
3818 following double entry harvest would require tipping of 10 and 7% of cut stems without and with
3819 the 10-m buffer. The authors conclude that thinning without some mitigation efforts resulted in
3820 large losses of in stream wood over a century. However, by including a 10-m no cut buffer or a
3821 practice of mechanical tipping can offset these losses Although predictions from this study
3822 contribute to the in-stream wood recruitment conversation moving forward, the model contained
3823 limitations such as utilizing data from FIA plots which only approximate riparian forest
3824 conditions.

3825

3826 **Modeling Stream Litter Delivery**

3827

3828 Bilby & Heffner, 2016

3829

3830 Bilby, R.E., Heffner, J.T., 2016. Factors influencing litter delivery to streams. *Forest Ecology and*
3831 *Management* 369, 29–37. <https://doi.org/10.1016/j.foreco.2016.03.031>

3832 The purpose of this study was to understand the relative influence of wind speed and direction,
3833 topography, litter type, species, and stand conditions on the distance from which litter is
3834 delivered to streams. This study utilized a combination of field experiments, literature, and

3835 simple models to estimate the width of a delivery areas. The effects of wind speed on litter
3836 delivery distance were measured on litter samples from two common species of the Pacific
3837 Northwest, Douglas-fir and red alder by releasing litter from a riparian tree canopy at various
3838 wind speeds and recording the distances traveled for each litter type at each wind speed. The
3839 relationship between distance of litter recruitment area and variables of interest (e.g., wind speed,
3840 topography, litter type...) were determined with a linear mixed effects model Data for wind speed
3841 and direction was recorded for one year in 30 min intervals along Humphrey Creek in the
3842 Cascade Mountains of western Washington. Results showed that under the wind conditions
3843 recorded at Humphrey Creek the majority of the litter recruited into the stream originated from
3844 within 10 m of the stream regardless of litter or stand type. No difference was found in delivery
3845 distance and litter type (needles or broadleaf) at young sites. However, needles released at mature
3846 sites had a higher proportion of cumulative input from greater distances than needles or alder
3847 leaves released at younger sites. This is likely due to the higher canopy and thus higher release
3848 position. Litter travel distance was linearly related to wind speed ($p < 0.0001$) Doubling wind
3849 speed at one site led to a 67-87% expansion of the riparian contribution zone in the study area.
3850 The results reveal a trend that suggests slope also contributes to the width of the litter
3851 contributing area. However, the authors did not apply statistical analysis to these values and only
3852 speculate that increasing the slope from 0-45% would increase the width of the litter contributing
3853 area by 70%. Overall, the results of this study show evidence that wind speed has a strong effect
3854 on the width of litter delivery areas within riparian areas, but that relationship is also affected
3855 stand age and litter type. Trends in the data also suggest that topography is an important factor,
3856 but it was not quantified.

3857

3858 **Stream Temperature**

3859

3860 Bladon et al., 2016

3861

3862 Bladon, K.D., Cook, N.A., Light, J.T., Segura, C., 2016. A catchment-scale assessment of stream
3863 temperature response to contemporary forest harvesting in the Oregon Coast Range. *Forest
3864 Ecology and Management* 379, 153–164. <http://dx.doi.org/10.1016/j.foreco.2016.08.021>

3865

3866 The purpose of this study was to compare the effects of contemporary riparian forest harvest
3867 treatments under the Oregon Forest Practices Act (15 m riparian management area with a
3868 minimum of ~3.7 m² conifer basal area retained for every 300 m length of stream) with historical
3869 riparian forest harvest practices (no maintenance of streamside vegetation) on stream
3870 temperatures. This study took place in the Siuslaw National Forest in the Oregon Coast Range
3871 as part of the Alsea Watershed Study Revisited. Historical records of stream temperatures were
3872 sourced from the original Alsea Watershed Study that monitored stream temperature changes
3873 from 1958-1973, before and after streamside timber harvesting in 1966. Stream temperature data

3874 was collected for contemporary forest practices over a 6-year period (3 years pre- and 3 years
3875 post-harvest; 2006-2012). Data for the contemporary harvest was also compared with stream
3876 temperature changes in unharvested reference streams to support a Before-After-Control Impact
3877 (BACI) design. Stream temperature thermistors were installed, and data was taken at 30-minute
3878 intervals at three sections of both the harvested (2 within harvest boundary and 1 downstream)
3879 and reference sites. Mean canopy closure, as measured with a densiometer, along the stream
3880 channel in the harvested portion of Needle Branch was reduced from ~96% in the pre-harvest
3881 period to ~89% in the post-harvest period. Comparatively, mean canopy closure along the stream
3882 channel in the reference sites were ~92% in the pre-harvest period and 91% in the post-harvest
3883 period. Data was analyzed to assess whether there were changes in the 7-day moving mean of
3884 daily maximum stream temperature, mean daily stream temperature, and diel stream temperature
3885 following harvest. The results showed no significant changes in any of the three parameters
3886 measured following contemporary forest harvesting practices when analyzed across all
3887 catchments for all summer months (July to September). When the mean 7-day moving maximum
3888 temperature was constrained to the summer period between July 15 – August 15 across all sites
3889 there was a significant increase in stream temperatures in the harvested sites by $0.6 + 0.2^{\circ}\text{C}$
3890 following harvest. However, when the data was arranged for individual pair-wise comparisons
3891 with the unharvested sites, and intrinsic annual and site variability was accounted for, the
3892 increases in stream temperature (ranging from $0.3 \pm 0.3^{\circ}\text{C}$ to $0.8 \pm 0.3^{\circ}\text{C}$) were not significant at
3893 any site. The only comparison made in the study to the original Alsea Watershed study was with
3894 the single day maximum stream temperatures for pre- and post-harvest. The contemporary
3895 practices showed a change of single day maximum stream temperatures from 15.7°C to 14.7°C
3896 (a reduction) from pre- to post-harvest. In contrast, the historical stream temperature data showed
3897 an increase in single day maximum stream temperatures from 13.9°C (pre-harvest) to as much
3898 as 29.4°C (2-years post-harvest). The authors caution that while these results support the
3899 conclusion that contemporary forest practices in Oregon are sufficient in maintaining stream
3900 temperatures after riparian forest harvest, and much more efficient than historical practices; these
3901 results should not be generalized to areas outside of coastal Oregon. The authors caution that the
3902 streams in this study have potential for a muted stream temperature response following harvest
3903 relative to other regions because of the (1) north-south stream orientation, which would
3904 maximize RMA effectiveness (2) steep catchment and channel slopes that can increase stream
3905 velocity and hyporheic exchange, (3) potential increases in groundwater contributions after
3906 harvest.

3907

3908 **Stream temperature**

3909

3910 Bladon et al., 2018

3911

3912 Bladon, K.D., Segura, C., Cook, N.A., Bywater-Reyes, S., Reiter, M., 2018. A multicatchment
3913 analysis of headwater and downstream temperature effects from contemporary forest harvesting.
3914 *Hydrological Processes* 32, 293–304. <https://doi.org/10.1002/hyp.11415>

3915

3916 The purpose of this study was to (1) examine the effects of contemporary forest harvesting
3917 practices on headwater stream temperature, (2) determine if increased temperatures from
3918 harvesting was detectable in downstream fish-bearing streams, and (3) examine the relative role
3919 of geology and forest management on influencing the differential stream temperature responses
3920 in both headwater and downstream reaches. This study took place at three paired watershed
3921 studies, of which two (Alesa, Trask) were located in the Oregon coast range, and one (Hinkle)
3922 was located in the western Cascades of Oregon. This study featured pre- and post-harvest
3923 measurements, as well as measurements within and downstream from harvested and reference
3924 sites. Buffer widths at harvested sites varied but averaged 20 m on either side of streams.
3925 Statistical models were generated which analyzed whether (a) the 7-day moving average of daily
3926 maximum stream temperature (7daymax) changed between pre- and post-harvest sites, and (b)
3927 whether post-harvest changes in 7daymax were detectable downstream. A regression analysis
3928 was also performed to assess the relative relationship between catchment lithology and percent
3929 catchment harvested on temperature at all sites. Statistical models were generated for each
3930 harvest site and reference pair. The pre-harvest relationship in stream temperatures for paired
3931 sites were used to create predicted changes in stream temperatures post-harvest. The post-harvest
3932 stream temperatures were then compared to the predicted values and the 95% prediction
3933 intervals. If post-harvest values of the 7daymax were outside the prediction interval the authors
3934 referred to these observations as statistical “exceedances”. Results showed that the 7daymax
3935 exceeded the predictive interval at 7 of the 8 harvested headwater sites (within the harvested
3936 boundary) when analyzed across all harvest years. The exceedances were largest in the first year
3937 after harvest but diminished in the second and third year at two treatment sites. However, at one
3938 site, the elevated 7daymax continued for three years post-harvest. In 4 of the 7 harvested sites
3939 with exceedances, the exceedances were recorded between 22 and 100% of the time. Smaller
3940 increases in stream temperatures were detected in the other 3 streams with exceedances, the
3941 exceedances occurred < 15% of the time. There was no evidence of elevated stream temperatures
3942 beyond the predicted intervals in any of the downstream sites following harvesting. The
3943 magnitude of change in stream temperature and transmission of warmer water downstream were
3944 a function of percentage of catchment harvested and the underlying geology. Although, these
3945 relationships were scale dependent. At the upstream, harvested sites there was a strong
3946 relationship between stream temperature increases and catchment lithologies, but no statistically
3947 significant relationship between stream temperature changes and percent of catchment harvested.
3948 Sites downstream from harvested areas showed a strong relationship with the interaction of
3949 percentage of catchment harvested and the underlying lithologies. The greatest temperature
3950 increases at downstream sites were in areas with a higher percentage of catchment harvested and
3951 were underlain by more resistant lithologies. There was no evidence for increases in stream
3952 temperatures in catchments with a high percentage of harvest that were underlain by permeable
3953 geology. The authors suggest that this relationship may be due to the buffering effect of increases

3954 in summer low flows and greater groundwater or hyporheic exchange. They conclude that the
3955 variability of rock permeability and the relative contribution of groundwater during summer
3956 months, and their effect on stream temperatures following harvest should be investigated further.

3957

3958 **Wood Loading**

3959

3960 Burton et al., 2016

3961

3962 Burton, J.I., Olson, D.H., Puettmann, K.J., 2016. Effects of riparian buffer width on wood
3963 loading in headwater streams after repeated forest thinning. *Forest Ecology and Management*
3964 372, 247–257. <https://doi.org/10.1016/j.foreco.2016.03.053>

3965

3966 The purpose of this study was to examine the relationship between in-stream wood loading and
3967 riparian buffer width in thinned stands in conjunction with several stand, site, and stream
3968 variables. This study is a part of a larger density management study which covered 6 sites along
3969 the coastal and western Cascade Range of Oregon. The sites used for this study were dominated
3970 by Douglas-fir and ranged in age from 30-70 years old. Two consecutive thinning treatments
3971 took place on a portion of each site, while the other portions were designated as an unthinned
3972 control. Treated sites featured one of four buffer width prescriptions: (1) ~ 70-m buffer
3973 representative of one site potential tree, (2) ~15-m buffer, (3) a 6-m buffer representative of trees
3974 immediately adjacent to the stream. Wood surveys were carried out at four times during the
3975 study: (1) prior to the first thinning, (2) five years after the first thinning, (3) 9-13 years after the
3976 first thinning and just prior to the second thinning, and (4) one year after the second thinning. At
3977 each site, the first thinning was to 200 trees per ha (tph), the second thinning (~10 years later)
3978 was to ~85 tph, alongside an unthinned reference stand ~400 tph. Spatial and geomorphic
3979 characterization were measured using a combination of field and geospatial data. Hierarchical
3980 linear mixed models were developed with repeated measures using a multi-step process to
3981 examine relationships between large wood volume in headwater streams over time and in-stream
3982 wood characteristics (decay stage, zone), buffer width, time since thinning, and reach and
3983 geomorphology (drainage basin area, width:depth ratio, gradient). Wood volume was found to
3984 increase exponentially with drainage basin area; for every 1-ha increase in drainage basin area,
3985 wood volume increased by 0.63%. Slightly higher volumes of wood were found in sites with a
3986 narrow 6-m buffer, as compared with the 15-m and 70-m buffer sites in the beginning 5 years
3987 after the first harvest and maintained through year 1 of the second harvest (end of study). The
3988 authors attributed this difference to a higher likelihood of logging debris and/or windthrow but
3989 was not analyzed. Low volumes of wood from stands in the stem-exclusion phase were found to
3990 contribute to overall in-stream wood. The results showed that between 82-85% of the wood with
3991 discernable sources (90% for wood in early stages of decay; 45% of wood in late stages of
3992 decay) came from within 15 m of the stream, and the relative contribution of wood to streams

3993 declined rapidly with increasing distance. The authors hypothesize that this finding in
3994 conjunction with their results, which show a positive relationship between basin area and wood
3995 volume suggests a greater role for other large wood recruitment processes such as creep,
3996 landslides, and debris flow.

3997

3998 **Sediment**

3999

4000 Bywater-Reyes et al., 2018

4001

4002 Bywater-Reyes, S., Bladon, K.D., Segura, C., 2018. Relative Influence of Landscape Variables
4003 and Discharge on Suspended Sediment Yields in Temperate Mountain Catchments. *Water*
4004 *Resources Research* 54, 5126–5142. 10.1029/2017WR021728

4005

4006 The purpose of this paper was to improve our ability to predict suspended sediment yields by
4007 quantifying how sediment yields vary with catchment lithography and physiography, discharge,
4008 and disturbance history. This study took place at the HJ. Andrews Experimental Site in the
4009 Western Cascade Range of Oregon. The questions this paper sought to answer were (1) What is
4010 the relative association between discharge and catchment setting (i.e., lithology and
4011 physiography) and suspended sediment yields over an ~60-year period (2) Is there an
4012 association between historical forest management activities (i.e., forest harvesting and road
4013 building) or extreme hydrologic events and the spatial and temporal trends in suspended
4014 sediment yield Data was collected from 10 catchments, 8 within the Lookout Creek Watershed, 1
4015 just below the Lookout Creek Watershed, and 1 that drains to the adjacent Blue River. The data
4016 set spanned a 60-year period from 1955-2015 Methods for determining suspended sediment
4017 concentration involved using either vertically integrated storm-based grab samples, or discharge-
4018 proportional composite samples where composite samples were collected every three weeks at
4019 the outlet of each catchment. A linear mixed effects model (log transformed to meet the
4020 normality assumption) was used to predict annual sediment yield. In this model, site was treated
4021 as a random effect while discharge and physiographic variables were treated as fixed variables.
4022 This allowed for the evaluation of the relationships between sediment yield and physiographic
4023 features (slope, elevation, roughness, and index of sediment connectivity) while accounting for
4024 site. To account for the effect of disturbance history a variable was added to the model when the
4025 watershed had a history of management or natural disturbances. If the models for the disturbed
4026 watersheds significantly underpredicted the sediment discharge, the timing of the sudden
4027 increases were further examined to assess whether it correlated with a disturbance event. Last,
4028 the authors considered changes in stage derived from comparing measured historic stage values
4029 to those predicted from current rating curves. Changes in stage were interpreted as a relative bed-
4030 elevation change resulting from changes in scour and deposition of material likely moved as
4031 bedload. The results of this study show that sediment yield varied greatly across space and time

4032 with the lowest annual yield occurring in 2001 (~0.2 t/km²) at one catchment, and the highest
4033 annual yield (~953 t/km²) occurring in 1969 at another catchment. Annual suspended sediment
4034 yield was most strongly correlated with the standard deviation of watershed slope (r=0.72), Only
4035 moderately correlated with slope (r = 0.32), and with drainage area (r = 0.38). Standard deviation
4036 of slope was also strongly correlated with TPI (a surface roughness index), and standard
4037 deviation of index of connectivity. When considering disturbance, the largest magnitude changes
4038 in bed-elevation (I.e., sediment movement), were after floods with a ≥ 30-year return interval.
4039 The authors conclude that variability in watershed slope was the best predictor of annual
4040 suspended sediment yield relative to other physiographic variables. The authors report that the
4041 variability in watershed slope combined with cumulative annual discharge explained 67% of the
4042 variation in annual sediment yield across the 60-year data set. The results, however, show that
4043 annual sediment yields also moderately correlated with many other physiographic variables and
4044 caution that the strong relationship with watershed slope variability is likely a proxy for many
4045 processes, encompassing multiple catchment characteristics. For example, the strong relationship
4046 between watershed slope standard deviation and surface roughness. For the relationships
4047 between disturbance and sediment yield the authors conclude that the few anomalous years of
4048 high sediment yield occurred in watersheds with high slope variability and within a decade of
4049 forest management and a large flood event. The authors further caution that the high variability
4050 of sediment yield over space and time indicate that the factors tested in this study should be
4051 tested more broadly to investigate their utility to forest managers.

4052

4053

4054 **LW, Wildfire**

4055

4056 Chen et al., 2005

4057

4058 Chen, X., Wei, X., Scherer, R., 2005. Influence of wildfire and harvest on biomass, carbon pool,
4059 and decomposition of large woody debris in forested streams of southern interior British
4060 Columbia. *Forest Ecology and Management* 208, 101–114. doi:10.1016/j.foreco.2004.11.018

4061

4062 The purpose of this study was to compare the components of in-stream LW features between
4063 wildfire and forest harvesting disturbances. This study focuses particularly on the change in
4064 biomass and carbon pool among LW under different disturbances. This study was located in the
4065 central Okanagan Valley, Kelowna, British Columbia. A total of 19 forest streams, first and
4066 second order, within the study area were divided into four categories based on disturbance
4067 history of the adjacent upland forest and included: (1) riparian forest harvested 10 years ago; (2)
4068 riparian forest harvested 30 years ago; (3) riparian forest burnt ~ 40 years ago; and (4)
4069 undisturbed old-growth riparian forests that had a mean forest age of 163 years.. All harvested
4070 streams were clear-cut to the stream edge. New trees had established on these sites within 1-3

4071 years of harvest (planted or natural growth) and resulted in lodgepole pine being the dominant
4072 species. The wildfire streams included those that had been burnt ~40 years ago with no post-fire
4073 harvest or salvage logging. In stream LW was recorded for analysis if it had a minimum diameter
4074 of 10 cm and length of 1.0 m and were situated within the bankfull width. LW biomass was
4075 determined through the conversion of wood density and wood volume. LW was also categorized
4076 by decay class (3 classes), species, orientation submergence, and distance from the beginning of
4077 the study reach. Sampling took place during the period between July and October 2003 along a
4078 150 m study reach for each stream. An analysis of variance was used to determine the
4079 relationships between the chosen variables. When significant differences were found, the data
4080 was further analyzed with the data was fitted with a linear regression model to obtain
4081 correlations between the three variables (volume, biomass, and carbon). Results from this study
4082 show that on average the riparian sites disturbed by wildfire had the highest biomass, volume,
4083 and carbon content for individual LW pieces, followed by the 10-year harvest, then the old-
4084 growth forest; the 30-year harvest had the lowest of all streams for all parameters. Mean LW
4085 biomass of each individual piece of wood was significantly higher in sites which had been
4086 burned than in harvested sites. Biomass values were, on average, 31 kg in the wildfire sites,
4087 compared to 21 kg and 19 kg for sites harvested 10 years ago and 30 years ago, respectively. The
4088 volume of individual pieces in wildfire sites was significantly higher than in old-growth sites,
4089 and nearly significantly higher than in sites harvested 30 years ago. No statistical significance
4090 was found comparing piece volume in wildfire sites to sites harvested 10 years ago. The average
4091 carbon content of individual pieces of wood was also highest in the wildfire sites but the
4092 differences were not significant. The authors present data that the LW found in the wildfire and
4093 30-year harvest sites was mostly in the third decay class (most decayed), with less than 1% of
4094 LW in the class 1 decay class. Statistical significance was not discussed in the results for
4095 differences in decay class. The authors conclude that streams adjacent to wildfire disturbed and
4096 recently harvested (10-years post-harvest) forests contained significantly higher LW individual
4097 pieces and total volume than old-growth and 30-year post-harvest sites. Further because biomass,
4098 volume, and carbon were significantly higher in the 10-year post harvest sites, but there was no
4099 difference in the 30-year post-harvest sites and the old-growth sites; the authors speculate that
4100 harvest can increase the abundance of LW in the short-term from leaving harvest residues but
4101 reduces the abundance of LW over the long-term (~30 years post) due to a lack of recruitment
4102 from the young forests, and loss of in-stream LW from decomposition. The three main takeaways
4103 presented by the authors for this paper were (1) LWD input in old growth forested streams was
4104 relatively stable, (2) timber harvesting activities would cause a short-term increase of LWD
4105 stocks and might greatly reduce LWD loadings over a long-term, and (3) wildfire disturbance
4106 would delay LWD recruitment because not all burnt trees would fall in the stream immediately
4107 after the wildfire.

4108

4109

4110 **LW**

4111

4112 Chen et al., 2006

4113

4114 Chen, X., Wei, X., Scherer, R., Luider, C., Darlington, W., 2006. A watershed scale assessment of
4115 in-stream large woody debris patterns in the southern interior of British Columbia. *Forest
4116 Ecology and Management* 229, 50–62. <https://doi.org/10.1016/j.foreco.2006.03.010>

4117

4118 The purpose of this study was to (1) determine the spatial distribution and variation of LW
4119 characteristics (size, amount, volume, mass, orientation, position) within different order streams
4120 of forested watersheds; (2) to examine the relationship between LW characteristics and stream
4121 features through channel networks; and (3) to estimate the total density, volume and mass of LW
4122 at the watershed scale using a combination of field surveys and GIS data. This study took place
4123 at three different watersheds located in the south-central interior of British Columbia near
4124 Kelowna. A total of 35 study reaches with stream orders ranging from first- through fifth-order
4125 were selected to measure spatial distribution and variability of LW characteristics. Data collected
4126 for each reach was binned into 4 stream size categories (I = first order; II = second to third order;
4127 III = third to fourth order; IV = fourth to fifth order). Study sites were selected based on the
4128 following criteria. (1) the streams were in areas of intact mature riparian forests (>80 years); (2)
4129 the stream side forests were not disturbed by human activities, such as harvesting, road building;
4130 (3) the streams were not salvaged. Therefore, the results from this study provide a baseline of
4131 LWD characteristics in intact mature riparian forests in the southern interior of British Columbia.
4132 LW in this study is defined as having a diameter of > 0.1 m and a length > 1.0 m. LW
4133 characteristics (decay class, orientation, position within channel, distance from downstream end
4134 of channel) were recorded for any piece of LW that was within or above the bankfull width of the
4135 channel. Watershed features and the distribution of stream orders were derived from remotely
4136 sensed data. Mean values of LW density, volume, and biomass were compared between stream
4137 size classes with an analysis of variance (ANOVA). Results from this study show that LW size,
4138 volume, and biomass generally increased with increasing stream size. For example, the mean
4139 LWD diameter in stream size I (16.4 cm) was lower than that in stream size III (20.6 cm) and
4140 size IV (20.5 cm), respectively. Mean LW length also increases with stream size from 2.3 m in
4141 size I, 2.9 m in size II, 3.1 m in size III, and 3.9 m in size IV. Stream IV had the highest mean
4142 volume (0.18 m³), significantly higher than stream size I (0.06 m³). LW volume was also
4143 significantly lower than in stream sizes II, and III. LW density (pieces per 100 m² of stream
4144 area), however, decreased as stream size increased. For example, LW density (defined as piece
4145 numbers per 100 m²) numbers were 19, 17, 12, and 4 for stream size I, II, III, and IV
4146 respectively. Increases in channel bankfull width ($R^2 = 0.52$) and stream area ($R^2 = 0.58$) was
4147 found to be strongly inversely correlated with LW density. Taken together, this study shows that
4148 spatial variation and distribution of LW characteristics vary as a function of stream size. From
4149 their results the authors conclude that in small sized streams, LW exhibit high density (number of
4150 pieces per 100 m²), low volume and biomass per unit area of stream. While in large sized
4151 streams, LW number, volume and biomass per unit of stream area are low but mean individual
4152 LW size was high.

4153

4154 **Stream Temperature Response to Harvesting**

4155

4156 Gravelle & Link, 2007

4157

4158 Gravelle, J.A., Link, T., 2007. Influence of Timber Harvesting on Headwater Peak Stream
4159 Temperatures in a Northern Idaho Watershed. *Forest Science* 53, 189–205.

4160

4161 The purpose of this study was to examine the effects of clearcutting and partial cutting on
4162 summer peak water temperatures in downstream fish-bearing streams, and to measure direct
4163 harvesting impacts on peak water temperature within headwater catchments. This study took
4164 place at the Mica Creek Experimental Watershed in Northern Idaho. Three headwater drainages
4165 were used to assess harvesting impacts on stream temperatures: (1) Watershed 1 which had 50%
4166 of the drainage area clearcut in 2001; (2) Watershed 2 which was thinned to a 50% target shade
4167 removal in Fall 2001; (3) and an unimpacted control. Riparian buffers were applied adjacent to
4168 the streams under the Idaho Forest Practices Act. This means, for fish-bearing streams the
4169 riparian management area must be at least 75 ft (22.9 m) wide on each side of the ordinary high-
4170 water mark (definable bank). Harvesting is still permitted, but there is a restriction where 75% of
4171 existing shade must be left. There are also leave tree requirements, which is a target number of
4172 trees per 1,000 linear feet (305 m), depending on stream width. For non-fish-bearing streams
4173 there is a 30 ft (9.1 m) equipment exclusion zone on each side of the ordinary high-water mark
4174 (definable bank). There are no shade requirements and no leave tree requirements, but skidding
4175 logs in or through streams is prohibited. Stream temperature data and canopy cover percentage
4176 data were collected at multiple sites within and downstream of treatment areas between 1992-
4177 2005. However, for the non-fish-bearing, headwater sites pre-treatment data was only collected
4178 one season prior to treatment. Temperature data was summarized as maximum daily temperature
4179 and was analyzed using simple linear regression to estimate changes in stream temperature
4180 following harvest during the summer months (July 1 – September 1). Results from this study
4181 show that there is no strong evidence of a posttreatment increase in stream temperature at long-
4182 term downstream sampling points for each harvest treatment. In general, the downstream sites
4183 showed a cooling effect between -0.2 and -0.3°C . The estimated cooling effect could not be
4184 attributed to any cause (e.g., increase in water yield), but the authors conclude that there was no
4185 post-harvest increase in peak summer temperatures at the downstream sites. For streams
4186 immediately adjacent to the clearcut treatment (headwater streams) a significant increase in
4187 temperature was detected at 2 sites ranging between 0.4 and 1.9°C , while a marginally
4188 significant decrease in temperature was detected at the third site (-0.1°C , $p = 0.06$). At the sites
4189 located immediately adjacent to partial cuts, results showed mixed results with decreases in
4190 temperature (-0.1°C ; non-significant) at one site and significant but minimal changes at another
4191 site (0.0 - 3.0°C) across the individual post-harvest years. Overall, there were minimal to no
4192 changes in stream peak temperatures following treatment in the partial-cut riparian areas. The

4193 authors go on to point out that headwater stream temperatures were highly variable, and that the
4194 shade value of understory vegetation may be an important factor contributing to results.

4195

4196 **SED**

4197

4198 Bywater-Reyes et al., 2017

4199

4200 Bywater-Reyes, S., Segura, C., Bladon, K.D., 2017. Geology and geomorphology control
4201 suspended sediment yield and modulate increases following timber harvest in temperate
4202 headwater streams. *Journal of Hydrology* 548, 754–769.
4203 <https://doi.org/10.1016/j.jhydrol.2017.03.048>

4204

4205 The purpose of this study was to assess the influence of natural controls (basin lithology and
4206 physiography) and forest management on suspended sediment yields in temperate headwater
4207 catchments. The study sought to achieve three objectives: (1) Quantify how suspended sediment
4208 yield varies by catchment setting in forested headwater catchments, (2) Determine whether
4209 contemporary forest management practices impact annual suspended sediment yield (SSY) in
4210 forested headwater catchments (3) Determine whether there are natural catchment settings that
4211 result in different levels of vulnerability or resilience to increases in suspended sediment yield
4212 associated with disturbances (e.g., harvest activities). This study analyzed 6 years of data from
4213 the Trask River Watershed in Northeastern Oregon and included data from harvested and
4214 unharvested sub-catchments underlain by heterogeneous lithologies. Baseline SSY data collection
4215 began in water year 2010 and continued through water year 2015, with road upgrades (July–
4216 August 2011) and harvest (May–November 2012) occurring in the middle of the study period.
4217 Generalized least square candidate models quantifying the parameters from each site were used
4218 to test differences in the relationship between suspended sediment yield and catchment setting.
4219 Results from this study indicate that site lithology was a first order control over SSY with SSY
4220 varying by an order of magnitude across lithologies observed. Specifically, SSY was greater in
4221 catchments underlain by Siletz Volcanics ($r = 0.6$), the Trask River Formation ($r = 0.4$), and
4222 landslide deposits ($r = 0.9$) and displayed an exponential relationship when plotted against
4223 percent watershed area underlain by these lithologies, combined. In contrast, the site effect had a
4224 strong negative correlation with percent area underlain by diabase ($r = 0.7$), with the lowest SSY
4225 associated with 100% diabase independent of whether or not earthflow terrain was present.
4226 Following timber harvest (water year 2013), increases in SSY occurred in all harvested
4227 catchments. The SSY in both PH4 (clearcut with buffers) and GC3 (clearcut without buffers)
4228 declined to pre-harvest levels by water year 2014. Interestingly, the SSY in UM2 (clearcut
4229 without buffers) increased annually throughout the post-harvest period, ultimately resulting in
4230 the highest SSY of all catchments during the final two years of the study after producing the
4231 lowest SSY in the pre-harvest period. Catchment physiographic variables (hypsoetry, slope,

4232 standardized topographic position index (SD TPI), and sediment connectivity (IC)) appeared to
4233 be good indicators of the underlying lithology of each site. Principle component analysis
4234 constructed from physiographic variables separated sites underlain by resistant diabase from
4235 those underlain by mixed lithologies along the PC1 axis. While sites along the second axis (PC2)
4236 were separated by relative values of earthflow terrain (high proportion vs. Little to none). Sites
4237 with low SSY and underlain by more resistant lithologies were also resistant to harvest-related
4238 increases in SSY. The authors conclude that sites underlain with a friable lithology (e.g.,
4239 sedimentary formations) had SSYs an order of magnitude higher, on average, following harvest
4240 than those on more resistant lithologies (intrusive rocks). In general, sites with higher SSY also
4241 had 1) lower mean elevation and slope, 2) greater landscape roughness, and 3) lower sediment
4242 connectivity (potential for sediment transport based on physiography). The authors suggest that
4243 their research be undertaken in different regions with different disturbance types to broadly apply
4244 their findings.

4245

4246 **Plant Communities**

4247

4248 D'Souza et al., 2012

4249

4250 D'Souza, L.E., Six, L.J., Bakker, J.D., Bilby, R.E., 2012. Spatial and temporal patterns of plant
4251 communities near small mountain streams in managed forests. *Can. J. For. Res.* 42, 260–271.
4252 <https://doi.org/10.1139/x11-17>

4253

4254 The purpose of this study was to examine spatial and temporal patterns in plant communities
4255 along fish-bearing streams in western Washington. The focus of this study is on areas which were
4256 harvested to the streambank within the last 100 years. The study took place in the western
4257 Cascade Mountains of Washington. Sites were randomly selected using a geographic information
4258 system. Stands that had been impacted by road development were excluded. Stands were
4259 stratified into a chronosequence of age classes: young (31-51 years), mature (52-70 years), old
4260 (>100 years). Due to availability, the sample sizes included 11 young stands, 10 mature stands,
4261 but only 4 old stands. Vegetation characteristics were captured in each stand using 0.16 ha plots
4262 located 30 m from stand edges to limit the influence of adjacent stands. Transects perpendicular
4263 to the stream were used 10 m apart and extended 80 m upslope. Vegetation and physical features
4264 along each transect were sampled using a series of subplots at 10 m intervals from the channel.
4265 The authors found little variation in riparian landform type and or canopy cover and were not
4266 included in the analysis for their effect on vegetation. Plant communities were examined
4267 spatially as a function of distance to stream and temporally by using the chronosequence of stand
4268 ages. Three distinct plant communities were observed in the shrub and herb layer (riparian: 0-9
4269 m; transitional: 10-29 m; and upslope: 30-80 m) and their composition differed significantly
4270 between communities. A total of 12 species were identified as indicators of these communities.

4271 For the shrub layer, community composition differed between old stands and young and mature
4272 stands. In the herb layer, community composition differed between all age classes. The results
4273 from this study suggest that plant communities along small fish-bearing streams have distinct
4274 changes in community with distance to stream, but also reflect successional status in nearby
4275 forests. The authors conclude by suggesting increased research in understanding the effects of
4276 forest management on streamside vegetation.

4277

4278 **LW Residence Time**

4279

4280 Hyatt & Naiman, 2001

4281

4282 Hyatt, T.L., Naiman, R.J., 2001. The Residence Time of Large Woody Debris in the Queets
4283 River, Washington, Usa. *Ecological Applications* 11, 191–202. [https://doi.org/10.1890/1051-
4284 0761\(2001\)011\[0191:TRTOLW\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[0191:TRTOLW]2.0.CO;2)

4285

4286 The purpose of this study was to determine the depletion rate of LW by examining differences in
4287 size and species composition in the Queets River compared to the adjacent forest. This study
4288 took place in the Queets River Watershed located on the west slope of the Olympic Mountains in
4289 Washington. Field sampling was carried out at 25 transects and four different sites. Increment
4290 cores from in-stream LW were cross-dated against cores from riparian conifers to estimate the
4291 time which LW was recruited into the channel. LW pieces which were in a heightened state of
4292 decay were dated using carbon-dating techniques. the most common tree species (> 30 cm
4293 diameter) in the riparian zone is red alder, followed by Sitka spruce and western hemlock,
4294 whereas the most common species of LWD (> 30 cm diameter) is Sitka spruce, followed by red
4295 alder and western hemlock. Each of the hardwood species is better represented among standing
4296 trees than among LWD, and each of the conifers are better represented as LW than among trees
4297 in the riparian zone. The depletion curve developed in the results was based only on conifer LW
4298 because hardwood LW was either too small or too young to provide accurate estimates of
4299 residence time in the stream. Based on the depletion curve developed for all available LW
4300 showed that wood typically disappears from the active channel within the first 50 years, while
4301 some pieces may remain for several hundred years. By cross-referencing the LW depletion
4302 curves with field notes the authors suggest that the longer residence time, beyond 50 years, was
4303 dependent on more than one process such as burial. Decay class was not an accurate predictor of
4304 LW age. Also, Dependent vegetation on or around LWD was a poor and often misleading
4305 indicator of residence time. Many LWD pieces that had 1–5 year old vegetation growing on

4306 or around them were discovered to have died and presumably recruited to the channel 20 years
4307 previous. The authors conclude that LW originating from hardwoods is depleted faster than
4308 conifers. Considering the depletion rate curve, the authors speculate that the majority of LW is
4309 transported out of the system within 50 years, while pieces of LW that are buried or jammed in

4310 the river floodplain may remain for hundreds of years. Overall, ~80% of LW residing in the
4311 active channel were living within 50 years of the study. The authors explain there are several
4312 caveats to the depletion curve created for this study (1) the depletion constant was developed for
4313 a large, mostly alluvial river and should probably not be applied to smaller streams (mean
4314 bankfull width at study transects on the Queets is 165 m and the range is 51–398 m; mean key
4315 LWD length is 23.4 m, and the range is 5.3–69.0 m). Also, from the data the authors infer that
4316 alluvial channel trap wood from upstream, and constrained channels export LWD downstream,
4317 so it is not to be expected that the LWD resident in a channel was recruited from the riparian
4318 zone in that reach. In general, the authors conclude that for this study the depletion curve shows
4319 that the half-life of LW is ~20 years and thus all resident LW will be exported, buried, or broken
4320 down within 3-5 decades. Also, hardwood LW will be depleted from the channel more rapidly
4321 than conifers.

4322

4323 **Litter Input**

4324

4325 Hart et al., 2013

4326

4327 Hart, S.K., Hibbs, D.E., Perakis, S.S., 2013. Riparian litter inputs to streams in the central
4328 Oregon Coast Range. *Freshwater Science* 32, 343–358. <https://doi.org/10.1899/12-074.1>

4329

4330 The purpose of this study was to understand how riparian vegetation composition, understory
4331 density, and topography affect the quantity and quality of litter input to streams throughout the
4332 annual cycle. This study took place within 5 contiguous watersheds located in the central Coast
4333 Range of Oregon. At each of the study sites uniform areas along a ≤ 300 m stream reach, 3 plots
4334 were delineated on 1 side of the stream, each 8x 25 m along the stream. Three treatments were
4335 applied: (1) a no cut or fence control; (2) cut and remove a 5 x 8 m section adjacent to stream
4336 plants < 10 cm DBH and >12 cm height every 2 months; and (3) 5 m fence extending
4337 underground and parallel to the stream to block litter moving downslope from reaching stream.
4338 Vertical and lateral litter traps were installed at each site and collected monthly between August
4339 2003-August 2004. Variation of riparian vegetation and woody debris characteristics were
4340 analyzed with a 3-way ANOVA using overstory, treatments, and sections and their interactions.
4341 Two-way ANOVA with repeated measures was used to compare seasonal and monthly control
4342 and treatment inputs for different overstory and litter types. 1-way ANOVA was used to test for
4343 differences in nutrient concentration flux between overstory type. Results from this study show
4344 that deciduous forests dominated by red alder delivered significantly greater vertical and lateral
4345 inputs to stream than did coniferous forests dominated by Douglas-fir. Deciduous-site vertical
4346 litter input (mean, 95% CI; 504 g m⁻¹ y⁻¹, 446.6–561.9) exceeded that from coniferous sites
4347 (394 g m⁻¹ y⁻¹, 336.4–451.7) by 110 g/m² (28.6–191.6) over the full year. Annual lateral inputs
4348 at deciduous sites (109 g m⁻¹ y⁻¹, 75.6–143.3) were 46 g/m (1.2– 94.5) more than at coniferous

4349 sites (63 g m⁻¹ y⁻¹, 28.9–96.6). Lateral inputs calculated for a 3-m-wide stream accounted for
4350 9.6% (5.4–12.5) of total annual inputs at coniferous sites and 12.7% (10.2–14.5) of total inputs at
4351 deciduous sites. Composition of litter also differed significantly by overstory type. Annual lateral
4352 inputs at coniferous sites were dominated by deciduous leaves (.33%), twigs (.23%), and leftover
4353 (.18%) litter types, whereas annual lateral inputs at deciduous sites were deciduous leaves (.61%)
4354 and leftover (.15%) litter types. Leftover litter types were defined as those that were too small or
4355 decayed to identify, bark, moss, or lichens. Vertical litter inputs at deciduous sites were
4356 dominated by deciduous leaves (.65%) and deciduous-other (.15%) litter types. While deciduous
4357 leaves (.33%), coniferous needles (.24%), and twigs (.21%) composed the annual vertical litter
4358 inputs at coniferous sites. The strongest deciduous inputs to streams occurred in November.
4359 Annual lateral litter input increased with slope at deciduous sites ($R^2 = 0.4073$, $p = 0.0771$), but
4360 showed no strong relationship at coniferous sites ($R^2 = 0.1863$, $p = 0.2855$). Total nitrogen flux
4361 to streams at deciduous sites was twice as much as recorded at coniferous sites. However, there
4362 was seasonal effect where the N fluxes in deciduous sites was only higher in autumn. The
4363 authors of this study conclude by suggesting management in riparian areas consider utilizing
4364 deciduous species such as red alder for greater total N input to aquatic and terrestrial ecosystems
4365 along with the increased shade and large woody debris provided by coniferous species.

4366

4367 **Effect of Contemporary Management on Nutrient Concentration and Cycling**

4368

4369 Gravelle et al., 2009

4370

4371 Gravelle, J.A., Ice, G., Link, T.E., Cook, D.L., 2009. Nutrient concentration dynamics in an
4372 inland Pacific Northwest watershed before and after timber harvest. *Forest Ecology and*
4373 *Management* 257, 1663–1675. <https://doi.org/10.1016/j.foreco.2009.01.017>

4374

4375 The purpose of this study was to assess the effects of contemporary forest harvesting practices on
4376 nutrient cycling and concentrations. This study took place at the Mica Creek Experimental
4377 Watershed in Northern Idaho. Seven steel Parshall flumes were installed at select locations
4378 within the watershed to assess the effects of clearcut to stream and partial cut (50% shade
4379 retention) harvesting practices. All harvesting was conducted in compliance with the Idaho
4380 Forest Practices Act. Within fish-bearing streams (Class I) Harvesting is permitted, but 75% of
4381 existing shade must be retained. There are also leave tree requirements for a target number of
4382 trees per 1000 linear feet (305 m), depending on stream width. In Mica Creek, this was roughly
4383 200 trees in the 3–12 in. (8–30 cm) diameter class per 305 m of the riparian management zone
4384 (RMZ). Along non-fish-bearing streams (Class II) the RMZ is 30 feet (9.1 m) of equipment
4385 exclusion zone on each side of the ordinary high-water mark (definable bank); skidding logs in
4386 or through streams is prohibited. There are no shade requirements and no requirements to leave
4387 merchantable trees. Two-sided riparian buffers were left on all Class I streams during harvest

4388 operations. Timber was removed from both sides of the Class II streams. In the post-harvest and
4389 post-burn conditions, Class II streams in clearcut treatments had only a small amount of green
4390 tree retention within the riparian zone, while in partial cut treatments equal amounts of canopy
4391 cover (approximately 50%) were removed from both sides of the stream. This study followed the
4392 BACI design and featured a pre-treatment measurement phase (1992-1997), a post-road
4393 construction phase (1997-2001), and a post-harvest phase (2001-2006). A student's t-test was
4394 used to analyze the data between the observed and predicted values of post-treatment sites for
4395 several nitrogen and phosphorus compound concentrations (Kjeldahl nitrogen (TKN), nitrate +
4396 nitrite (NO₃ + NO₂), TP, total ammonia nitrogen (TAN) consisting of unionized (NH₃) and
4397 ionized (NH₄⁺) ammonia, and unfiltered orthophosphate (OP) samples). Results from the post-
4398 road construction period showed no significant changes in concentrations of any nutrients
4399 analyzed. Results from this study show statistically significant increases in NO₃ and NO₂
4400 concentrations following clearcut and partial harvest cuts in headwater streams. Increases at the
4401 clearcut treatment site were greatest, where mean monthly concentrations increased from 0.06
4402 mg-N L⁻¹ during the calibration and post-road periods to 0.35 mg-N L⁻¹. There was also an
4403 observable seasonal effect on NO₃ + NO₂ concentrations with the peak concentration of 0.89
4404 mg-N L⁻¹ occurred at F1 in April 2004, with mean monthly concentrations of 0.43 mg-N L⁻¹
4405 and 0.59 mg-N L⁻¹ in water years (October–September) 2004 and 2005, respectively. Similar
4406 results were also observed at sites further downstream although changes were smaller which, the
4407 authors point out this may be due to in-stream uptake and/or dilution. No significant changes of
4408 in-stream concentration of any other nutrient recorded were found between time periods and
4409 treatments except for one downstream site that showed a small increase in orthophosphate by
4410 0.01 mg P L⁻¹. In general, the results of this study show that forest management influences in-
4411 stream NO₃ + NO₂ immediately adjacent to treatment and downstream of treatment. The authors
4412 conclude by suggesting future research in understanding variability in nutrient concentrations
4413 and cycling as affected by seasons and storm runoff events.

4414

4415 **Organic Matter Inputs**

4416

4417 Kiffney & Richardson, 2010

4418

4419 Kiffney, P.M., Richardson, J.S., 2010. Organic matter inputs into headwater streams of
4420 southwestern British Columbia as a function of riparian reserves and time since harvesting.
4421 *Forest Ecology and Management* 260, 1931–1942. <https://doi.org/10.1016/j.foreco.2010.08.016>

4422

4423 The purpose of this paper was to assess how differences in riparian buffer width and timing since
4424 harvest affect terrestrial particulate organic matter flux into streams. The focus of this paper was
4425 on 1st and 2nd order headwater streams located approximately 45 km east of Vancouver in
4426 British Columbia, Canada. Sites were measured over an 8-year period and included clear-cut

4427 (n=3), 10-m buffered reserve (n=3), 30-m buffered reserve (n=2), and uncut control (n=2)
4428 treatments. For streams receiving a 10 or 30-m reserve, there was no logging on either side of the
4429 stream within these reserves. Study reaches were approximately 200m long. Vertical litter inputs
4430 were collected monthly and at approximately 6–8-week intervals during each season for years
4431 1,2,6,7, and 8 years after harvest. Litter was separated into broadleaf deciduous, twig, needles,
4432 and other (seeds, cones, and moss) categories following collection and subsequently dried and
4433 weighed using a microbalance. A mixed-model analysis of covariance was used for Fall data
4434 with riparian treatment as a fixed effect and year as a covariate. Secondly, ordinary least
4435 squares regression was used to quantify the functional relationship between reserve width and
4436 litter flux within each year. Results show riparian treatments having significant effects on the
4437 quantity and composition of litter input into streams. Inputs consisting of needles and twigs were
4438 significantly lower while deciduous inputs were higher in clearcuts compared to other
4439 treatments. Differences in litter flux relative to riparian treatment persisted through year 7, while
4440 a positive trend between reserve width and litter flux remained through year 8. For example, one-
4441 year post-treatment, needle inputs were 56x higher during the Fall into control and buffered
4442 treatments than into the clearcut. Needle inputs remained 6x higher in the buffer and control sites
4443 through year 7, and 3-6x higher in year 8 than in the clearcut sites. Twig inputs into the control
4444 and buffered sites were ~25x higher than in the clearcut sites in the first year after treatment.
4445 There was no significant difference in treatment for deciduous litter but a trend of increasing
4446 deciduous litter input in the clear cut was observed in the data. For example, one-year post-
4447 treatment deciduous litter was lowest in the clearcut, but by year 8 deciduous litter was highest in
4448 the clearcut sites relative to control and buffered sites. The linear relationship between reserve
4449 width and litter inputs was strongest in the first year after treatment, explaining ~57% of the
4450 variation, but the relationship could only explain ~17% of the variation in litter input by buffer
4451 width by year 8 (i.e., the relationship degraded over time). The authors interpret these results as
4452 evidence that riparian reserves showed a similar litter flux to streams when compared to uncut
4453 controls. They also conclude that litter flux from riparian plants to streams, was affected by
4454 riparian reserve width, time since logging, and potentially channel geomorphology.

4455

4456 **In-stream Wood Loads**

4457

4458 Jackson & Wohl, 2015

4459

4460 Jackson, K.J., Wohl, E., 2015. Instream wood loads in montane forest streams of the Colorado
4461 Front Range, USA. *Geomorphology* 234, 161–170.
4462 <http://dx.doi.org/10.1016/j.geomorph.2015.01.022>

4463

4464 The purpose of this study was to examine in-stream wood loads and geomorphic effects between
4465 stands of different ages and stands with different disturbance histories The first objective of this

4466 study was to determine whether instream wood and geomorphic effects differ significantly
4467 among old-growth, younger, healthy, and beetle-infested forest stands. The second objective of
4468 this study was to determine whether instream wood loads correlate with valley and channel
4469 characteristics. The authors hypothesized that streams in old-growth montane forests have (1)
4470 significantly larger in stream and floodplain wood loads than those in younger stands, (2) greater
4471 frequency of volume of jams than those in younger forests, and (3) more wood created
4472 geomorphic effects. They also hypothesized that instream wood loads in healthy montane forests
4473 are significantly smaller than in beetle-infested forests. Last, they hypothesized that instream
4474 wood load correlates with lateral valley confinement, with unconfined valleys having the greatest
4475 in-stream and total wood loads. This study took place within the Arapaho and Roosevelt National
4476 Forests in Colorado. Sediment storage, channel geometry, in-stream wood load, and forest stand
4477 characteristics were measured along 33 pool-riffle or plane-bed stream reaches (10 located in
4478 old-growth (> 200 years); 23 located in younger forests (age range not reported)). LW
4479 characteristics were recorded for all in-stream wood ≥ 10 cm diameter and ≥ 1 m in length. Pair-
4480 wise t-test or Kruskal-Wallis tests were used to check for significant differences in wood load,
4481 logjam volume, and logjam frequencies. To test for significant differences in wood created
4482 geomorphic effects a principal component analysis was used. Results indicated that channel
4483 wood load (OG = 304.4 ± 161.1 ; Y = 197.8 ± 245.5 m³/ha), floodplain wood load (OG = 109.4
4484 ± 80 ; Y = 47.1 ± 52.8 m³/ha), and total wood load (OG = 154.7 ± 64.1 ; Y = 87.8 ± 100.6 m³
4485 /ha) per 100 m length of stream and per unit surface area were significantly larger in streams of
4486 old-growth forests than in young forests. Streams in old-growth forests also had significantly
4487 more wood in jams, and more total wood jams per unit length of channel than in younger forests
4488 (jam wood volume: OG = 7.10 ± 6.9 m³; Y = 1.71 ± 2.81 m³). When standardized to stream
4489 gradient, old-growth streams had significantly greater pool volume and significantly greater
4490 sediment volume than younger stands. No significant difference was detected in in-stream wood
4491 loads between healthy and beetle-infested stands. Although wood load in streams draining from
4492 pine beetle infested forests did not differ significantly from healthy forests, best subset regression
4493 (following principal component analysis) indicated that elevation, stand age, and pine beetle
4494 infestation were the best predictors of wood load in channels and on floodplains. The authors
4495 speculate that beetle infestation is affecting in-stream wood, but perhaps not enough time has
4496 passed since the infestation for the affected trees to fall into the stream. Time since beetle-
4497 infestation was not reported.

4498

4499 **LW Recruitment**

4500

4501 May & Gresswell, 2003

4502

4503 May, C.L., Gresswell, R.E., 2003. Large wood recruitment and redistribution in headwater
4504 streams in the southern Oregon Coast Range, U.S.A. *Can. J. For. Res.* 33, 1352–1362.

4505 <https://doi.org/10.1139/x03-023>

4506

4507 The purpose of this study was to understand the relative influence of processes that recruit and
4508 redistribute wood into channels and to understand how these processes vary spatially. Specific
4509 research questions included the following:(i) Do processes that deliver and redistribute wood
4510 differ in small colluvial channels compared with larger alluvial channels? (ii) Do proximal and
4511 distal controls on wood delivery differ for colluvial and alluvial channels? (iii) How do input and
4512 redistribution processes influence the functional role of wood in the channel? The focus of this
4513 research is specifically on differences between small colluvial channels and large alluvial
4514 channels in the southern Oregon Coast Range. All downed wood exceeding 20 cm mean
4515 diameter and 2 m in length, and in contact with the bank-full channel were measured in three
4516 second order and one third-order stream. Large wood was categorized based on the various
4517 mechanisms delivering it to the stream channel. Categories included (i) direct delivery from local
4518 hillslopes and riparian areas, (ii) fluvial redistribution, (iii) debris flow transported, or (iv) an
4519 unidentified source. Results from this study show that stream size and topographic position
4520 strongly influence processes that recruit and redistribute wood in channels. Processes of slope
4521 instability were shown to be important conveyors of wood from upland forests to small colluvial
4522 channels. In the larger alluvial channels, windthrow was found to be the dominant recruitment
4523 process from adjacent riparian area. Results showed that Wood derived from local hillslopes and
4524 riparian areas accounted for the majority of pieces (63%) in small colluvial channels. The larger
4525 alluvial channel received wood from a greater variety of sources, including recruitment from
4526 local hillslopes and riparian areas (36%), fluvial redistribution (9%), and debris flow transported
4527 wood (33%). However, because pieces recruited from local sources (hillslope and riparian area)
4528 were larger, these sources of wood had a disproportionately large contribution to volume of wood
4529 in the stream. For example, wood recruited from the local hillslopes and riparian areas accounted
4530 for 36% of wood pieces in the alluvial stream, which accounted for 74% of the total volume of
4531 wood. Slope instability and windthrow were the dominant mechanisms for wood recruitment into
4532 small colluvial channels. Windthrow was the dominant recruitment mechanism for wood
4533 recruitment into larger alluvial channels. Distributions of the source distance of wood pieces
4534 were significantly different between colluvial and alluvial channels. In colluvial streams, 80% of
4535 total wood and 80% of total wood volume recruited originated from trees rooted within 50 m of
4536 the channel. In the alluvial channel, 80% of the pieces of wood and 50% of the total volume
4537 originated from trees which came from 30 m of the channel. The primary function of wood in
4538 smaller colluvial channels was sediment storage (40%) and small wood storage (20%). The
4539 primary function of wood in larger alluvial channels is bank scour (26%), stream bed scour
4540 (26%), and sediment storage (14%). Recruitment and redistribution processes were shown to
4541 affect the location of the piece relative to the channel/flow direction, thus influencing its
4542 functional role. The authors conclude that wood recruited from local sources is variable by
4543 position in the stream network because of differences in recruitment processes, degree of
4544 hillslope constriction, and slope steepness.

4545

4546 **Sediment**

4547
4548 Macdonald et al., 2003a
4549
4550 Macdonald, J. S., Beaudry, P. G., MacIsaac, E. A., & Herunter, H. E. (2003). The effects of forest
4551 harvesting and best management practices on streamflow and suspended sediment concentrations
4552 during snowmelt in headwater streams in sub-boreal forests of British Columbia, Canada.
4553 Canadian Journal of Forest Research, 33(8), 1397-1407. <https://doi.org/10.1139/x03-110>
4554
4555 (BACI, only single year pre-harvest)
4556
4557 This study investigates the changes in suspended sediment concentration and stream discharge
4558 during freshet (spring snowmelt) at two harvest intensities relative to each other and an
4559 unharvested control watershed, pre- and post-harvest. The design included three small sub-
4560 boreal, first order, forest streams (<1.5 m width) in the central interior of British Columbia
4561 (Baptiste watershed). Both treatment streams received a 55% harvest treatment; one (low-
4562 retention) removed all merchantable timber >15 cm DBH for pine and > 20 cm DBH for spruce
4563 within 20 m of the stream; the other treatment (high-retention) removed all merchantable timber
4564 > 30 cm within 20 m of the stream; and an un-harvested control. Data for stream flow and total
4565 suspended sediments (TSS) was collected using Parshall flumes downstream from the treatment
4566 and control sites for one-year pre- and four-years post-harvest during snowmelt periods.
4567 Regression analysis was used to analyze relationships between treatment and control reaches pre-
4568 and post-treatment to estimate and compare predicted changes in TSS. The results showed an
4569 increase in freshet discharge for both treatments above predicted values for the entirety of the
4570 study. During the year prior to treatment, TSS relationships of both treatment watersheds during
4571 freshet closely matched those of the control. Immediately following harvest TSS concentrations
4572 increased above predicted values for both treatment streams. Increased TSS persisted for two-
4573 years post-harvest in the high-retention treatment, and for 3-years in the low-retention. The
4574 authors speculate that the treatment areas may have accumulated more snow (e.g., more exposed
4575 area below canopy) than in the control reaches leading to the increase in discharge. This study
4576 shows evidence that harvest intensity (low vs. high retention) is proportional to the increase in
4577 stream discharge, TSS, and recovery time to pre-harvest levels.
4578
4579 **LW**
4580
4581 Fox & Bolton, 2007
4582

4583 Fox, M., & Bolton, S. (2007). A regional and geomorphic reference for quantities and volumes of
4584 instream wood in unmanaged forested basins of Washington State. *North American Journal of*
4585 *Fisheries Management*, 27(1), 342-359. <https://doi.org/10.1577/M05-024.1>

4586

4587 This study uses in-stream LW values from 150 stream segments located in unmanaged
4588 watersheds, across all of Washington State, to investigate the relationships between
4589 geomorphology, forest zone, and disturbance regimes with LW recruitment. The purpose of this
4590 study was to create a base-line value of central tendency for in-stream LW values in “natural”
4591 streams for which salmonids are theoretically adapted. The authors define natural and
4592 unmanaged as streams that (1) had no part of the basin upstream of the survey site ever logged
4593 using forest practices common after European settlement and (2) the basin upstream of the
4594 survey site contains no roads or human modifications to the landscape that could affect the
4595 hydrology, slope stability, or other natural processes of wood recruitment and transport in
4596 streams. Sites were stratified to capture the variations in forest types, channel morphologies, and
4597 hydrological origins. The authors used descriptive statistics to establish and evaluate correlations
4598 between wood loading and watershed characteristics to reveal the highest valued variables
4599 influencing wood loading. Following this analysis, the variables with the highest mechanistic
4600 values in determining wood loading were evaluated and compared using simulation modeling.
4601 Results showed that in-stream wood volume increased with drainage area and as streams became
4602 less confined. However, bank full width (BFW) was a significantly better predictor of wood
4603 parameters than basin size. There was observational evidence that alluvial channels contained
4604 more wood volume on average than bedrock channels. However, due to limits in sample size
4605 following stratification, statistical analysis could not be completed. Sample sizes for isolating
4606 gradient and confinement were also too small to apply statistical analyses. Fire was found to
4607 influence in-stream wood quantities and volumes west of the Cascade crest; In-stream wood
4608 volume increased with adjacent riparian timber age as determined by the last stand replacing fire.
4609 Other disturbances such as debris flow, snow avalanche, and flooding were too few in frequency
4610 in the study area to be analyzed statistically. From these results the authors developed thresholds
4611 for expected “key piece volume (m^3) (pieces with independent stability) of wood for three BFW
4612 classes (20-30 m, >30 – 50 m, > 50 m width) per 100 m stream length for streams with BFW
4613 greater than 20 m. From percentile distributions the authors recommend minimum volumes,
4614 defined by the 25th percentiles, of approximately 9.7 m³ for the 20- to 30-m BFW class, 10.5 m³
4615 for the 30- to 50-m BFW class, and 10.7 m³ for channels greater than 50 m BFW per 100 m
4616 length of stream. The results of this study suggest that BFW is the single greatest predictor of in-
4617 stream wood quantity and volume relative to other predictor variables. However, this result
4618 comes with the caveat that other processes and geomorphologies (e.g., channel bed form,
4619 gradient, confinement) are also important in the mechanisms for wood recruitment, modeling in
4620 this study showed too much inconsistency with these predictor variables to draw strong
4621 conclusions. Further the authors warn that these values for reference conditions are only
4622 applicable to streams with bank-full widths between 1 and 100 m, gradients between 0.1% and
4623 47%, elevations between 91 and 1,906 m, drainage areas between 0.4 and 325 km², glacial and
4624 rain- or snow-dominated origins, forest types common to the Pacific Northwest.

4625

4626 **LW and sediment**

4627

4628 Gomi et al., 2001

4629

4630 Gomi, T., Sidle, R. C., Bryant, M. D., & Woodsmith, R. D. (2001). The characteristics of woody
4631 debris and sediment distribution in headwater streams, southeastern Alaska. *Canadian Journal of*
4632 *Forest Research*, 31(8), 1386-1399. <https://doi.org/10.1139/x01-070>

4633

4634 This study investigated different riparian conditions related to harvest and disturbance
4635 (landslides), their influence on woody debris and sediment distributions, and their related
4636 functions in headwater streams. This study examined the effects of recent and past timber
4637 harvests on woody debris abundance and distribution, landslides and debris flow on woody
4638 debris abundance and sediment accumulations, and the function of in-stream woody debris on
4639 sediment storage. The researchers examined 15 steep headwater streams in the Maybeso
4640 Experimental Forest and Harris River basin in the Tongass National Forest, Prince of Wales
4641 Island, southeastern Alaska. Treatments of headwater streams included five management or
4642 disturbance regimes: old growth (OG), recent clear-cut (CC; 3 years), young growth conifer
4643 forest (YC; 37 years after clear-cut), young growth alder (YA; 30 years after clear-cut), and
4644 recent landslide and debris flow channels (LS). Three headwater streams were sampled for each
4645 of the 5 treatments, 15 streams total. Analysis of covariance (ANCOVA) was used to compare
4646 LW quantity and distribution, and sediment quantity and distribution, across plots nested within
4647 each treatment site. Results showed in-channel numbers of LW pieces were significantly higher
4648 in YC and CC sites when compared to OG, YA, and LS sites. The number of LW pieces was
4649 highest in YC streams even though logging concluded 3 decades prior to sampling. No
4650 significant differences in LW volume were found among OG, CC, and YC streams. However,
4651 LW volume per 100 m of stream length in YC was twice that in OG. The total volume of LW per
4652 100 m associated with CC channels was half that in OG channels. However, the majority of the
4653 LW volume in OG systems was outside of the bank-full area. When the data was stratified by
4654 channels that experienced landslides (LS and YA), the number of LW pieces among OG, YA, and
4655 LS was not statistically significant. However, the in-channel volumes of LW in LS and YA
4656 channels were significantly lower than in OG sites because individual LW pieces in the OG sites
4657 were relatively larger than in the LS and YA sites. There was high variability among sites in the
4658 amount of sediment stored within streams. The authors conclude that timber harvesting and
4659 related landslides and debris flows affect the distribution and accumulation of LW and related
4660 sediment accumulation in headwater streams. These effects are summarized as (i) inputs of
4661 logging slash and unmerchantable logs significantly increase the abundance of in-channel woody
4662 debris; (ii) in the absence of landslides or debris flows, these woody materials remain in the
4663 channel 50–100 years after logging; (iii) relatively smaller woody debris initially stores
4664 sediment; (iv) when landslides and debris flows occur 3–15 years after logging because of

4665 intensive rain and weakening of root strength (Sidle et al. 1985), woody debris is evacuated from
4666 headwater streams and deposited in downstream reaches; (v) although less woody debris remains
4667 in the scour zone, woody debris pieces and jams contribute to sediment storage in both the scour
4668 and deposition zones of landslide and debris flow channels; (vi) red alder stands actively
4669 recolonize riparian zones of headwater streams for 20–50 years after mass movement and recruit
4670 woody debris and organic materials, which in turn provide sediment storage sites; and (vii)
4671 subsequent sediment movement after landslides and debris flows are affected by residual woody
4672 debris and newly introduced debris.

4673

4674 **LW and sediment**

4675

4676 Johnson et al., 2000 (removed from focal list)

4677

4678 Johnson, S. L., Swanson, F. J., Grant, G. E., & Wondzell, S. M. (2000). Riparian forest
4679 disturbances by a mountain flood—the influence of floated wood. *Hydrological processes*,
4680 14(16-17), 3031-3050. [https://doi.org/10.1002/1099-1085\(200011/12\)14:16/17<3031::AID-](https://doi.org/10.1002/1099-1085(200011/12)14:16/17<3031::AID-HYP133>3.0.CO;2-6)
4681 [HYP133>3.0.CO;2-6](https://doi.org/10.1002/1099-1085(200011/12)14:16/17<3031::AID-HYP133>3.0.CO;2-6)

4682

4683 This study examined the differences in riparian forest responses to a 100-year flood event along
4684 eight third- to fifth-order streams in the Cascade Mountain Range of Oregon. Disturbance
4685 intensities were grouped into three categories: purely fluvial (high water flow only), fluvial with
4686 uncongested wood transport, and fluvial with congested wood transport. Riparian forest
4687 responses were heavily influenced by pre-flood forest structure and disturbance/harvest history,
4688 especially the characteristics of LW presence within streams and along channels. The quantity
4689 and severity of toppled trees (fully uprooted vs. partially uprooted) during the flood event was
4690 proportional to the quantity and congestion of LW already present (i.e., higher volumes of LW
4691 already present during the flood event increased the frequency of toppled trees and newly
4692 deposited LW in streams). Further, stands that experienced higher frequencies of toppled trees
4693 also showed higher frequencies and magnitudes of debris flow. The authors concluded that the
4694 land use practices, and disturbance histories influenced the age and structure of the riparian
4695 forests, but also the availability of the agents of disturbance (presence of LW) during the 100-
4696 year flood event. This paper is a good discussion of how pre-disturbance structure affects the
4697 response of riparian forests to disturbances (in this case, flood), however, there is no statistical
4698 analysis discussed in the methods. This is purely descriptive science that involves an intensive
4699 survey of before and after riparian forest structures.

4700

4701 **Sediment**

4702
4703 Yang et al., 2022 (removed from focal list)
4704
4705 Yang, Y., Safeeq, M., Wagenbrenner, J. W., Asefaw Berhe, A., & Hart, S. C. (2022). Impacts of
4706 climate and forest management on suspended sediment source and transport in montane
4707 headwater catchments. *Hydrological Processes*, 36(9), e14684.
4708 <https://doi.org/10.1002/hyp.14684>
4709
4710 This paper investigates the changes in annual hysteresis patterns for in-stream suspended
4711 sediment in 10 headwater streams at 2 sites, Providence Creek (rain-snow-dominated,
4712 transitional), and Kings River Experimental Watershed (snow-dominated). Aside from
4713 precipitation pattern differences in the two catchments, the researchers also compared differences
4714 in hysteresis patterns for forested riparian control, burn-only, thin-only, and thin-and-burn
4715 combined areas. The differences in the proportion of clockwise-loop hysteresis patterns for
4716 suspended sediments in the warmer rain-snow-transition sites compared to the colder snow-
4717 dominated sites suggests that warming temperatures may cause the snow-dominated basins to
4718 receive sediment from extended source areas and for longer periods if they transition to rain
4719 dominated catchments. The results found no discernable difference in hysteresis loops between
4720 the control, burn-only, thin-only, and thin-and-burn combined areas. Further, there seemed to be
4721 little change in the hysteresis loops during drought, average, and excessively wet years. The
4722 authors speculate that local conditions will be more important in understanding the impacts of
4723 climate change than changes in precipitation patterns or average annual temperatures alone.
4724 Mainly, there is evidence that if snow-dominated watersheds become warm enough to transition
4725 to rain-dominated, there is potential for disruption to sediment discharge frequency, rates, and
4726 source distance. The indiscernible difference in hysteresis loops for the different treatments also
4727 suggests that management practices imposed to ameliorate these changes may not be completely
4728 effective.
4729
4730 **Nutrients**
4731
4732 Vanderbilt et al., 2003
4733
4734 Vanderbilt, K. L., Lajtha, K., & Swanson, F. J. (2003). Biogeochemistry of unpolluted forested
4735 watersheds in the Oregon Cascades: temporal patterns of precipitation and stream nitrogen
4736 fluxes. *Biogeochemistry*, 62(1), 87-117. DOI:10.1023/A:1021171016945
4737

4738 This study uses long-term datasets (ranging from 20-30 years) from six watersheds in the H.J.
4739 Andrews Experimental Watershed (HJA) in the west-central Cascade Mountains of Oregon to
4740 investigate patterns in dissolved organic nitrogen (DON) and dissolved inorganic nitrogen (DIN)
4741 export with watershed hydrology. The objectives of this study were to 1) characterize long-term
4742 patterns of N dynamics in precipitation and stream water at the HJA, 2) analyze relationships
4743 between annual output of N solutes and annual stream discharge, 3) analyze relationships
4744 between seasonal stream water N solute concentrations and precipitation and stream discharge,
4745 and 4) compare results with those from other forested watersheds. Precipitation data were
4746 collected at three-week intervals from 10/1/1968 until 5/24/1988 and at one-week intervals
4747 thereafter. Stream chemistry samples were collected weekly for the entirety of the study. Stream
4748 discharge was measured continuously throughout the study. The researchers used regression
4749 analysis of annual N inputs and outputs with annual precipitation and stream discharge to
4750 analyze patterns. The results showed DON was the largest component of N input at the low-
4751 elevation collector, followed by PON (particulate organic N), NO₃-N, and NH₄-N. At the high-
4752 elevation collector, NO₃-N input was higher than at low elevation and was the largest component
4753 of N in bulk and wet-only inputs, followed by NH₄-N, DON, and PON. For annual stream
4754 outputs, DON was the largest fraction of annual N output, followed by PON, NH₄-N and then
4755 NO₃-N. Total annual discharge was a positive predictor of annual DON export in all watersheds
4756 with r² values ranging from 0.42 to 0.79. In contrast, significant relationships between total
4757 annual discharge and annual export of NO₃-N, NH₄-N, and PON were not found in all
4758 watersheds. No systematic long-term average seasonal trends were observed for NO₃-N or PON
4759 concentrations. Elevated concentrations of NH₄-N occurred in spring and early summer in all
4760 three watersheds, although they are not convincingly synchronous. DON concentrations
4761 increased in the fall in every watershed. The increase in concentration began in July or August
4762 with the earliest rain events, and peak DON concentrations occurred in October through
4763 December before the peak in the hydrograph. DON concentrations then declined during the
4764 winter months. The authors conclude that total annual stream discharge was a positive
4765 predictor of DON output suggesting a relationship to precipitation. Also, DON had a consistent
4766 seasonal concentration pattern. All other forms of N observed showed variability and
4767 inconsistencies with annual and seasonal stream discharge. The authors speculate that different
4768 factors may control organic vs. Inorganic N export. Also, DIN may be strongly influenced by
4769 terrestrial or in-stream biotic controls, while DON is more strongly influenced by climate. Last,
4770 the authors suggest that DON in streams may be recalcitrant, and largely unavailable to stream
4771 organisms. The authors emphasize the importance of analyzing data from multiple watersheds in
4772 a single climactic zone to make inferences about stream chemistry.

4773

4774 **Stream temperature**

4775

4776 Roon et al., 2021b

4777

4778 Roon, D. A., Dunham, J. B., & Torgersen, C. E. (2021). A riverscape approach reveals
4779 downstream propagation of stream thermal responses to riparian thinning at multiple scales.
4780 *Ecosphere*, 12(10), e03775. <https://doi.org/10.1002/ecs2.3775>

4781

4782 This study uses a riverscape approach to evaluate the effects of streamside forest thinning on
4783 stream temperatures at multiple spatiotemporal scales. This study addresses the question of how
4784 thinning second-growth riparian forests influences local and downstream temperatures at
4785 watershed extents. This study attempts to answer this question by addressing four objectives: (1)
4786 quantify pretreatment spatial and temporal variability in stream temperature conditions; (2)
4787 evaluate local responses in stream temperature to riparian thinning; (3) assess the spatial extent
4788 and temporal duration of downstream effects to local responses in temperature; and (4)
4789 characterize local and downstream responses to thinning with a conceptual framework based on
4790 waveforms. The researchers compared upstream, local, and downstream, stream temperature
4791 fluctuations following different intensities of streamside forest thinning at 10 treatment reaches
4792 across three watersheds in the redwood forests of northern California. Treatments varied by
4793 landowners. In two watersheds thinning treatments were intended to reduce 50% of canopy
4794 closure within the riparian zone along a 200 m reach on both sides of the active channel. This
4795 treatment resulted in a reduction in effective shade over the stream between 19-30%. In the other
4796 treatment watershed, thinning treatments reduced basal area by as much as 40% on both sides of
4797 the active channel along a 100 m long reach. Reductions in effective shade over the stream in
4798 these sites ranged from 4-5%. The analysis considered each reach both individually and
4799 collectively to understand how site and treatment heterogeneity may affect thermal responses at
4800 local and watershed extents. Temperature data were collected before, during, and after treatment
4801 and in the thinned experimental reaches and in adjacent unthinned control reaches with digital
4802 temperature sensors. Temperature data was collected for only 1-year pre-treatment and 1-year
4803 post-treatment. For data analysis, semivariograms of summer degree days were used to
4804 determine the presence of spatial autocorrelation. To control temporal variations in local and
4805 downstream responses summer cumulative degree-days were plotted for pre- and post- treatment
4806 temperatures and along a longitudinal gradient. A Lagrangian framework was used to track
4807 changes in temperature through space and time. Results showed that increases in thermal
4808 heterogeneity occurred in the treatment reaches, in the year following treatment (20° to 139°C),
4809 compared to the pre-treatment year (66° to 112°C). Local changes in stream temperature were
4810 dependent on thinning intensity, with higher levels of canopy cover reduction leading to higher
4811 increases in local stream temperatures. In the reaches with higher reductions in shade (19-30%)
4812 there was accumulation of 45° to 115°C additional degree days from pre- to post treatment years,
4813 while the reaches with lower reductions in shade (4-5%) only accumulated 10° to 15°C
4814 additional degree days. Travel distance of increased stream temperatures also appeared to be
4815 dependent on thinning intensity. The lower shade reduction reaches had an increased temperature
4816 effect downstream with travel distance of 75-150 m, while the high shade reduction sites had a
4817 downstream travel distance of 300- ~1000 m. In the high shade reduction sites, treatment reaches
4818 that were further apart (> 400 m) showed dissipation in increased stream temperatures
4819 downstream, while in parts of the stream where treatments were <400 m apart, temperature

4820 increases did not always dissipate before entering another the next treatment reach. The analyses
4821 with the conceptual framework based on waveforms showed there was no evidence of
4822 cumulative watershed effects at the downstream extent. The authors conclude that their results
4823 show evidence that riparian forest management impacts may extend beyond local stream
4824 environments. Further, the authors propose that riparian forest management that uses a holistic
4825 approach may be more effective in preserving some functions (e.g., shade).

4826

4827 **Sediment**

4828

4829 Wissmar et al., 2004

4830

4831 Wissmar, R.C., Beer, W.N. & Timm, R.K. (2004) Spatially explicit estimates of erosion-risk
4832 indices and variable riparian buffer widths in watersheds. *Aquat. Sci.* 66, 446–455 . DOI:
4833 10.1007/s00027-004-0714-9

4834

4835 The purpose of this study is to use management records, the spatial distribution, and the
4836 variability of different landcover types that can contribute to unstable conditions to develop
4837 erosion-risk indices and variable riparian buffer widths in watersheds of different drainages in
4838 the State of Washington. The objectives of this study were to 1) define erosion risk indices based
4839 on “different land cover types,” 2) evaluate erosion risk indices with sediment inputs into
4840 streams, 3) use erosion risk categories to define locations of stream reaches that are susceptible
4841 to different levels of erosion 4) use categories to identify distribution of channels requiring
4842 variable width buffers for protection 5) Test procedure by applying ground-truthed data from the
4843 upper Cedar River drainage near Seattle, Washington. The land cover types used to assess risk
4844 included unstable soils, immature forests, roads, critical slopes for land failure, and rain-on-snow
4845 events. Based on available data, the researchers developed a map of these land cover features
4846 with sediment input values to define erosion risk indices. The indices were used to categorize the
4847 landscape into 6 levels of erosion risk. Results of the mapped erosion risk categories explained
4848 65% of the variation associated with sediment inputs. The highest-risk areas contained a
4849 combination of all landscape cover factor combinations (rain-on-snow zone, critical failure
4850 slope, unstable soil, immature forests, and roaded areas). The lowest risk categories contained
4851 only rain-on-snow zones, and critical failure slopes. Roaded areas and unstable soils were only
4852 present in risk categories 3-6. This paper shows the importance of investigating multiple factors
4853 when evaluating the controls on sediment discharge and stream inputs. Further, when factors
4854 influencing erosion combine in an area, their effects are compounded.

4855

4856 **Nutrient and forest structure**

4857
4858 Devotta et al., 2021 (removed)
4859
4860 Devotta, D. A., Fraterrigo, J. M., Walsh, P. B., Lowe, S., Sewell, D. K., Schindler, D. E., & Hu,
4861 F. S. (2021). Watershed Alnus cover alters N: P stoichiometry and intensifies P limitation in
4862 subarctic streams. *Biogeochemistry*, 153(2), 155-176. DOI:10.1007/s10533-021-00776-w
4863
4864 This study investigates how coverage of alder species affects the aquatic N and P availability
4865 across a natural alder coverage gradient in 26 streams of southwestern Alaska. Alder coverage in
4866 the Alaskan streams was inversely related to elevation (i.e., lower coverage at higher elevations).
4867 To identify the presence of alder as the N and p contributing factor, the researchers analyzed
4868 resin lysimeter samples from select watershed soils supporting variable percent coverages of
4869 alder. Soils supporting alders leached, on average, three times more N and two times more P than
4870 soils not containing alders. The relationship between alder coverage and N and P values was not
4871 linear. Still, the authors identified 30% alder coverage as a transitional threshold from low to
4872 markedly higher soil N and p availability. The higher soil N and P resulted in higher dissolved N
4873 in streams, but the higher soil P under alder coverage did not translate to higher stream P
4874 availability. The authors speculate that soil chemistry or local soil biota may be immobilizing the
4875 soil P from transport into the streams. This led to a high N:P ratio in the spring and summer
4876 stream chemistry of reaches supporting >30% alder coverage. As climate change causes
4877 increasing temperatures, alder may begin to expand its range into higher elevations. This, in turn,
4878 may lead to increased N availability, but higher P limitations in high-elevation montane streams.

4879

4880 **Sediment and lithology**

4881

4882 Fratkin et al., 2020 (removed from focal, scope and results not relevant to review)

4883

4884 Fratkin, M. M., Segura, C., & Bywater-Reyes, S. (2020). The influence of lithology on channel
4885 geometry and bed sediment organization in mountainous hillslope-coupled streams. *Earth
4886 Surface Processes and Landforms*, 45(10), 2365-2379. <https://doi.org/10.1002/esp.4885>

4887

4888 This study compares the differences in channel form patterns, sediment flow, grain size, and
4889 sheer stress thresholds between two gravel-bed streams, one on basalt and one on sandstone
4890 parent material in the Oregon Coast Range. Study sites were in a region where widespread
4891 landslides and debris flows occurred in 1996. The researchers compared channel
4892 geomorphologies (e.g., slope, valley width, channel geometry, etc.) to evaluate thresholds and

4893 channel bed adjustments since the 1996 events. The results showed similar sediment coarsening
4894 patterns in the first several kilometers indicating hillslope influence, but downstream fining was
4895 lithology dependent. The authors hypothesized threshold channel conditions in the basalt basin,
4896 and non-threshold conditions in the sandstone basin with a tendency to expose bedrock, based on
4897 the relative competencies (i.e., basalt = high-competency, sandstone = low-competency).
4898 However, results showed evidence of threshold conditions for over 60% of the streams in both
4899 basins. The authors inferred a cycle adjustment to correct the assumed sediment delivery from
4900 the 1996 flood season. The authors speculate that the basalt basins would act as threshold
4901 channels over longer time periods despite a higher debris flow frequency. This paper provides
4902 some evidence that lithologies impose control on channel adjustments driven by different rock
4903 competencies. This difference in rock competency ultimately controls the grain size fining rates
4904 and bed load transport (sediment availability).

4905

4906 **Nutrient and species composition**

4907

4908 Whigham et al., 2017 (removed from focal)

4909

4910 Whigham, D. F., Walker, C. M., Maurer, J., King, R. S., Hauser, W., Baird, S., ... & Neale, P. J.
4911 (2017). Watershed influences the structure and function of riparian wetlands associated with
4912 headwater streams—Kenai Peninsula, Alaska. *Science of the Total Environment*, 599, 124-134.
4913 <https://doi.org/10.1016/j.scitotenv.2017.03.290>

4914

4915 This field study was designed to test the hypothesis that alder cover in watersheds influences the
4916 structure and function of riparian wetlands adjacent to headwater streams. The researchers
4917 compared biomass production, biomass distribution (aboveground vs. belowground),
4918 decomposition rates, and chemical characteristics of interstitial groundwater, between watersheds
4919 with and without alder coverage. Study sites were located on two headwater streams located in
4920 the Kenai Peninsula in south-central Alaska. The results showed that aboveground biomass was
4921 higher in watersheds with alder cover, but the largest differences were in the litter layer and the
4922 belowground biomass. Watersheds without alder had significantly higher belowground root
4923 biomass. The litter overhanging the stream was higher in N content at the alder sites than in the
4924 no-alder sites. The quantity of litter overhanging the stream was higher in the no-alder sites.
4925 Interstitial groundwater was significantly higher in dissolved N at the alder sites. The results of
4926 this study show that species composition within the riparian area can have a considerable effect
4927 on nutrient concentrations which consequently affect stream chemistry, biomass production,
4928 vegetation structure, and decomposition rates.

4929

4930 **LW**

4931

4932 Wing & Skaugset, 2002

4933

4934 Wing, M. G., & Skaugset, A. (2002). Relationships of channel characteristics, land ownership,
4935 and land use patterns to large woody debris in western Oregon streams. *Canadian Journal of*
4936 *Fisheries and Aquatic Sciences*, 59(5), 796-807. <https://doi.org/10.1139/f02-052>

4937

4938 This study investigated the relationships of land use, land ownership, and channel and habitat
4939 characteristics with LW quantity and volume in 3793 stream reaches in western Oregon State
4940 (west of Cascade crest). This study analyzed an extensive spatial database of aquatic habitat
4941 conditions created for western Oregon using stream habitat classification techniques and a
4942 geographic information system (GIS). The overall objectives of this study were to identify the
4943 database factors most strongly related to LWD abundance and to determine whether ownership
4944 and land use patterns are related to LWD abundance. Regression tree analysis is an exploratory
4945 regression analysis that allows for the inclusion of multiple explanatory variables. LW counts (by
4946 piece, and by key pieces (logs at least 0.60 m in diameter and 10 m long)) and volume were used
4947 as the response variables and explanatory variables included morphology of active channel
4948 (hillslope, terrace, terrace hillslope, unconstrained), lithology (e.g., alluvium, basalt, etc.), Land
4949 use and land cover (e.g., young timber, old timber, rural resident, agriculture, etc.), ownership
4950 (private industrial (PI), private non-industrial (PNI), state, federal (BLM, USFS)), vegetation
4951 type, and other channel characteristics. The analysis was run at the reach scale. Results showed
4952 that the most important predictor for LW volume was land ownership with PNI split from all
4953 other ownership types. Mean LW volumes in stream reaches with PNI ownership were 3.1 m³
4954 while mean volume of LW in reaches in all other ownerships (PI, state, BLM, USFS) were 17.9
4955 m³. However, this was likely because the PNI lands held a disproportionately higher percentage of
4956 unforested lands compared to all other ownership types. When the ownership and land use
4957 variables were removed, stream gradient became the most important explanatory variable for LW
4958 volume. The split for stream gradient occurred for reaches with < 2.3% gradient averaged 5.8 m³
4959 while higher gradient streams averaged 17.9 m³ per reach. When ownership and land use were
4960 included but non-forested lands were removed, stream gradient again was the most important
4961 predictor with the split occurring for stream reaches with gradients less than 4.7% averaging 11.5
4962 m³, which was less than half of the average found at higher gradient reaches (25.2 m³); in this
4963 model the stream gradient split explained 11% of the variation observed of instream LW volume.
4964 For LW pieces in forested stream reaches bankfull channel width was the most important
4965 explanatory variable with the split occurring for streams channels less than 12.2 m wide. LW
4966 pieces for streams <12.2 m wide averaged 11.1 LW pieces per reach while larger channels
4967 averaged 4.9 pieces per reach; in this model the BFW split explained 7% of the variation in LW
4968 pieces found in forested streams. For key LW pieces (logs at least 0.60 m in diameter and 10 m
4969 long) in forested reaches, stream gradient was again the most important explanatory variable
4970 with the split occurring at a slope of 4.9%. The streams with a gradient < 4.9% averaged 0.5 key
4971 LW pieces per reach while streams with higher gradients averaged 0.9 key LW pieces per reach;

4972 in this model stream gradient explained 8% of the variation in key LW pieces found in streams.
4973 For forested streams, lithology caused second, third or fourth level splits after stream gradient or
4974 BFW. In three of these four splits, Mesozoic sedimentary and metamorphic geologies, located in
4975 southern Oregon stream reaches, were grouped and split from basalt, cascade, and marine
4976 sedimentary geologies. In stream reaches in Mesozoic sedimentary and metamorphic geologies,
4977 the quantity of LWD was roughly half the amount found in other geologies. The only exception
4978 to this grouping was for LW volume in larger stream reaches, where basalt and marine
4979 sedimentary geologies were grouped separately from all other geologies in a fourth-level split
4980 and contained more LW volume. The authors conclude that the geomorphic characteristics of
4981 stream reaches, in particular stream gradient and bankfull width, in forested areas correlated best
4982 with LW presence.

4983

4984

4985 **LW and plant communities**

4986

4987 Rot et al., 2000 (removed from focal list)

4988

4989 Rot, B. W., Naiman, R. J., & Bilby, R. E. (2000). Stream channel configuration, landform, and
4990 riparian forest structure in the Cascade Mountains, Washington. *Canadian Journal of Fisheries
4991 and Aquatic Sciences*, 57(4), 699-707. <https://doi.org/10.1139/f00-002>

4992

4993 This study investigates the hierarchical relationships between the “five key elements”, valley
4994 constraint, riparian landform, riparian plant community, channel type, and channel configuration.
4995 for 21 sites in mature old-growth riparian forests of the western Cascade Mountains in
4996 Washington State. The objective of this article is to expand this perspective over several spatial
4997 scales and the temporal life span of a conifer by examining how channel configuration interacts
4998 with valley constraint, streamside landform, channel bedform, and successional processes within
4999 the riparian forest. Stepwise regression was used to examine the relationship between physical
5000 and biological characteristics and the individual elements of channel configuration. Channel
5001 configuration is the channel elements at the habitat unit scale, including channel units (total
5002 number of pool–riffle habitat units per 100 m of channel length), LW pieces (per 100 m of
5003 channel length), LW volume (cubic meters per 100 m of channel length), pool spacing, percent
5004 pools, and percent LW-formed pools. Results showed that significantly more total LW pieces
5005 were found in forced pool–riffle channels than in the bedrock and plane-bed channels (Kruskal–
5006 Wallis, $p < 0.05$). Forced pool–riffle channels averaged 16.4 pieces per 100 m, bedrock 10.8
5007 pieces, and plane-bed 10.1 pieces. The volume of LW (cubic meters per 100 m) followed a
5008 similar trend. The percentage of deep pools (>0.5 m) formed by LW increased with stand age (r^2
5009 = 0.36). LW diameters were significantly smaller for ages 55–220 than for ages 333–727
5010 (Kruskal–Wallis, $p = 0.01$). The authors conclude that scale is an important consideration for

5011 management of aquatic habitat. At the largest spatial scale, results showed valley constraint
5012 significantly influenced off-channel habitat (plant communities associations and landform
5013 categories) and in-stream LW volume within forced pool-riffle channels. At the smallest scale,
5014 channel type (bedrock, plane-bed, and forced pool-riffle) was most closely related to LW
5015 volume, density, and the number of LW-formed pools. The diameter of the in-channel LW
5016 increased with riparian forest stand age. Streams adjacent to old-growth forests in-channel LW
5017 diameter were equivalent to or greater than the average standing riparian tree diameter at all
5018 sites. In younger stands, the relationship of in-stream LW diameter had a mixed relationship with
5019 riparian tree average diameters. The authors speculate this may be due to many in-stream LW
5020 pieces being relics from previous old-growth communities. In this area, four landform classes
5021 differentiated the riparian communities (floodplain, low terrace, high terrace, slope). Most were
5022 dominated by conifers, except the floodplain landforms, which supported a higher density of
5023 deciduous species, but a higher basal area of conifer species. The results of this study provide
5024 more evidence, similar to other studies, that channel geomorphology and valley constraint are
5025 important predictors of LW abundance (quantity and volume) in streams. The novelty in this
5026 study is how the riparian area landforms lead to different riparian plant communities, which
5027 consequently affect the input of LW.

5028

5029 **Nutrients**

5030

5031 Yang et al., 2021

5032

5033 Yang, Y., Hart, S. C., McCorkle, E. P., Stacy, E. M., Barnes, M. E., Hunsaker, C. T., ... & Berhe,
5034 A. A. (2021). Stream water chemistry in mixed-conifer headwater basins: role of water sources,
5035 seasonality, watershed characteristics, and disturbances. *Ecosystems*, 24(8), 1853-1874.
5036 DOI:10.1007/s10021-021-00620-0

5037

5038 This study investigated the effects of drought and forest thinning operations (independently and
5039 combined) on water chemistry from multiple basin water sources (snowmelt, soil solution,
5040 stream water) in the Mediterranean climate headwater basins of the Sierra National Forest. Data
5041 on water chemistry was taken 2 years prior and 3 years following drought and thinning
5042 operations in two watersheds, each with thinned and control stands. This data was analyzed to
5043 answer 3 questions: 1. How does the chemistry of different water sources (that is, snowmelt, soil
5044 solution at two depths, stream water) vary monthly and interannually prior to drought and
5045 thinning? 2. How does drought alone and drought combined with thinning impact water
5046 chemistry? 3. Can watershed characteristics predict stream water chemistry over contrasting
5047 water years? The authors used general linear models to analyze differences in chemistry by water
5048 source, repeated measures analysis of variance for effects of drought and thinning on water
5049 chemistry, and linear regression to predict water chemistry based on watershed characteristics.

5050 Results showed that monthly concentrations of dissolved C and N varied among different water
5051 sources prior to drought and thinning. For dissolved organic carbon (DOC) soil solution at 13 cm
5052 depth (mean \pm SE of $25.97 \pm 2.75 \text{ mg l}^{-1}$, across months for 2 years) had higher monthly
5053 concentrations than soil solution collected at 26 cm depth ($16.93 \pm 1.55 \text{ mg l}^{-1}$). Snowmelt (9.67
5054 $\pm 0.89 \text{ mg l}^{-1}$) and stream water ($5.33 \pm 0.52 \text{ mg l}^{-1}$) had the lowest concentrations. For total
5055 dissolved Nitrogen (TDN) and dissolved organic nitrogen (DON), soil solution at 13 cm depth
5056 (1.72 ± 0.57 and $1.66 \pm 0.57 \text{ mg l}^{-1}$, respectively), soil solution at 26 cm depth (0.94 ± 0.32 and
5057 $0.92 \pm 0.32 \text{ mg l}^{-1}$), and snowmelt (0.94 ± 0.17 and $0.73 \pm 0.18 \text{ mg l}^{-1}$) had higher
5058 concentrations than stream water (0.11 ± 0.02 and $0.08 \pm 0.01 \text{ mg l}^{-1}$). For dissolved inorganic
5059 nitrogen (DIN), snowmelt ($0.25 \pm 0.05 \text{ mg l}^{-1}$) had the highest concentration followed by the soil
5060 solution at 13 cm depth ($0.06 \pm 0.01 \text{ mg l}^{-1}$). Soil solution at 26 cm depth ($0.03 \pm 0.01 \text{ mg l}^{-1}$)
5061 and stream water had the lowest values ($0.04 \pm 0.01 \text{ mg l}^{-1}$). For pH, snowmelt (pH 6.09 ± 0.06)
5062 was more acidic than soil solutions at both depths (7.52 ± 0.23 at 13 cm depth and 7.79 ± 0.11 at
5063 26 cm depth) and stream water (7.37 ± 0.07). Drought alone altered DOC in stream water, and
5064 DOC:DON in soil solution in unthinned (control) watersheds. Volume-weighted concentration of
5065 DOC was 62% lower ($p < 0.01$) and DOC:DON was 82% lower ($p = 0.004$) in stream water in
5066 years during drought (WY 2013–2015) than in years prior to drought (WY 2009 and 2010).
5067 Drought combined with thinning altered DOC and DIN in stream water, and DON and TDN in
5068 soil solution. For stream water, volume-weighted concentrations of DOC were 66–94% higher in
5069 thinned watersheds than in control watersheds for all three consecutive drought years following
5070 thinning. No differences in DOC concentrations were found between thinned and control
5071 watersheds before thinning. Watershed characteristics explained inconsistently the variation in
5072 volume-weighted mean annual values of stream water chemistry among different watersheds.
5073 The authors conclude that their results showed evidence that the influences of drought and
5074 thinning are more pronounced for DOC than for N in streams.

5075

5076 **Geology**

5077

5078 Kusnierz and Sivers, 2018 (removed from focal)

5079

5080 Kusnierz, P.C., Sivers, E., 2018. How important is geology in evaluating stream habitat? *J Soils*
5081 *Sediments* 18, 1176–1184. DOI:10.1007/s11368-017-1885-z

5082

5083 The purpose of this study was to assess the importance of considering geology when evaluating
5084 stream habitat conditions. Stream habitat data were collected from 424 sites on federally
5085 managed lands in western Montana, USA. These sites represented a variety of ecoregions, stream
5086 types, management practices, and geologies. The importance of accounting for geology in data
5087 analysis was evaluated using five sediment-related habitat variables and three analyses that
5088 examined (1) differences across geology for the entire dataset and for sites in reference and

5089 managed watersheds; (2) differences between reference and managed sites within geologies; and
5090 (3) the relative strength of geology as a factor when accounting for the effects of management,
5091 stream type, and ecoregion. This objective was pursued by using five sediment-related habitat
5092 variables (Log instability index, Log roughness-corrected index of relative bed stability, Median
5093 substrate size, Percent pool tail fines < 6 mm, Percent stable banks). Five sediment-related
5094 habitat variables were collected from 424 sites on federally managed lands between 2009-
5095 2012. Factorial ANOVA on ranks was performed to evaluate the relative importance of geology
5096 when other factors were taken into account. Results from this study show that differences in
5097 sediment-related habitat variables did not differ significantly according to geology; however,
5098 observed differences were typically drawn from managed sites. The authors conclude by
5099 advising against using geology as the sole means of stratifying habitat data when attempting to
5100 account for between-site variability.

5101

5102 **Stream Temperatures**

5103

5104 Leach et al., 2017 (removed from focal list)

5105

5106 Leach, J.A., Olson, D.H., Anderson, P.D., Eskelson, B.N.I., 2017. Spatial and seasonal variability
5107 of forested headwater stream temperatures in western Oregon, USA. *Aquat Sci* 79, 291–307.
5108 DOI:10.1007/s00027-016-0497-9

5109

5110 This study is a case study of thermal regimes for headwater streams in the Keel Mountain Study
5111 area. This study examined (1) forested headwater stream temperature variability in space and
5112 time; (2) relationships between stream temperature patterns and weather, above-stream canopy
5113 cover, and geomorphic attributes; and (3) the predictive ability of a regional stream temperature
5114 model to account for headwater stream temperature heterogeneity. Stream temperature data was
5115 collected at 48 sites within a 128-ha watershed in western Oregon between 2012 and 2013.
5116 Spatial statistical modeling was used to relate stream temperature patterns to site characteristics
5117 (elevation, stream width, catchment area, slope, aspect, channel substrate, and terrain shading), a
5118 cluster analysis was used to capture the full variability in annual stream temperatures. Results
5119 from this study show considerable variability in stream temperature over relatively small areas,
5120 and between seasons. The greatest spatial variability existed during summer (up to 10 Celsius)
5121 and during cold and dry winter periods (up to 7.5 Celsius). Geomorphic attributes typically used
5122 in stream temperature models were not good predictors of variability at headwater scales.

5123

5124 **Stream Temperatures**

5125

5126 Groom et al., 2011b

5127

5128 Groom, J.D., Dent, L., Madsen, L.J., Fleuret, J.(2011b). Response of western Oregon (USA)
5129 stream temperatures to contemporary forest management. *Forest Ecology and Management* 262,
5130 1618–1629. <https://doi.org/10.1016/j.foreco.2011.07.012>

5131

5132 The objective of this paper was to assess the riparian characteristics that best predict shade, and
5133 to determine the stream temperature changes that result following harvest. This study took place
5134 in the Oregon Coastal Range at 33 sites (15 state-owned and 18 private-owned). The 33 sites
5135 studied were approximately 50-70 years old and predominately composed of Douglas-fir and red
5136 alder. Private sites (n = 18) followed FPA rules whereby the riparian management area (RMA)s
5137 are 15 and 21 m wide on small and medium fish-bearing streams, with a 6 m no-cut zone
5138 immediately adjacent to the stream. Harvesting is allowed in the remaining RMA to a minimum
5139 basal area of 10.0 (small streams) and 22.9 (medium streams) m²/ha. State sites (N = 15)
5140 followed the state management plan whereby a 52 m wide buffer is required for all fish-bearing
5141 streams, with an 8 m no cut buffer immediately adjacent to the stream. Limited harvest is
5142 allowed within 30 m of the stream only to create mature forest conditions. Harvest operations
5143 within this zone must maintain 124 trees per hectare and a 25% Stand Density Index. Additional
5144 tree retentions of 25–111 conifer trees and snags/hectare are required between 30 and 52 m. A
5145 site’s control reach was located immediately upstream of its treatment reach. The control reaches
5146 were continuously forested to a perpendicular slope distance of at least 60 m from the average
5147 annual high-water level. Reach lengths varied from 137 m to 1,829 m with means of 276 m and
5148 684 m for the control and treatment reaches, respectively. Temperature recording stations were
5149 located upstream and downstream of both control and treatment sites. Stream temperature data
5150 was summarized to provide daily minimum, maximum, mean, and fluctuation for analysis. The
5151 temperature data was modeled using mixed-effects linear regression. Shade analysis included
5152 trees per hectare, basal area per hectare, vegetation plot blowdown, and tree height. A linear
5153 regression analysis of shade data (n = 33) was performed and compared small-sample AIC values
5154 to determine relative model performance among 8 a priori models. Results showed that average,
5155 minimum, and diel stream temperatures increased on private sites following harvest, suggesting a
5156 relationship between decreased shade derived from buffer width and an increase in stream
5157 temperature. Outputs from the model predicted an increase of ~2 °C for minimum shade
5158 conditions and a decrease of ~ -1 °C for maximum shade conditions. For sites that exhibited an
5159 absolute change of shade > 6% from pre-harvest to post-harvest experienced an increase in
5160 maximum temperatures. Further, the model predicted an increase in stream temperature
5161 proportional to treatment reach length. The authors estimate an increase in maximum and
5162 minimum temperatures of 0.73 and 0.59 °C per km, respectively. Following harvest, maximum
5163 temperatures at private sites increased relative to state sites on average by 0.71 °C. Similarly,
5164 mean temperatures increased by 0.37 °C (0.24 - 0.50), minimum temperatures by 0.13 °C (0.03 -
5165 0.23), and diel fluctuation increased by 0.58 °C (0.41 - 0.75) relative to state sites. A comparison
5166 of within site changes in maximum temperatures pre-harvest to post-harvest showed an overall

5167 increase at private sites, but not all sites behaved the same and some had decreases in maximum
5168 temperatures. The average of maximum state site temperature changes = 0.0 °C (range = -0.89 to
5169 2.27 °C). Observed maximum temperature changes at private sites averaged 0.73 °C (range = -
5170 0.87 to 2.50 °C) and exhibit a greater frequency of post-harvest increases from 0.5 to 2.5 °C
5171 compared to state sites. Private site shade values also appeared to decrease pre-harvest to post-
5172 harvest. Private post-harvest shade values differed from pre-harvest values (mean change in
5173 Shade from 85% to 78%); however, no difference was found for state site shade values pre-
5174 harvest to post-harvest (mean change in Shade from 90% to 89%). They did not find evidence
5175 that shade differed if one or both banks were harvested for private sites although the sample size
5176 for single sided harvests was low. Similarly, private site shade values did not appear to differ
5177 between medium or small streams. Results from this study also show that between 68% and
5178 75% of variability in post-harvest shade may be accounted for by basal area within 30 m of the
5179 stream, tree height, and potentially blowdown. The authors speculate that their results suggest
5180 sites with shorter trees have higher post-harvest shade and this may be due to the negative
5181 correlation between crown ratios and tree heights. Overall, this study shows that buffers managed
5182 by state sites were sufficient at mitigating the effects of upland harvesting on stream temperature.
5183 Increases in stream temperature on private sites were related to decreases in shade, which were
5184 related to decreases in basal area on sites with greater tree heights. The authors suggest that their
5185 results are likely relevant to other high-rainfall low-order Douglas-fir dominated streams in the
5186 Pacific Northwest that are subject to similar harvest practices.

5187

5188 **Litter**

5189

5190 Yeung et al., 2019

5191

5192 Yeung, A. C., Stenroth, K., & Richardson, J. S. (2019). Modelling biophysical controls on stream
5193 organic matter standing stocks under a range of forest harvesting impacts. *Limnologia*, 78,
5194 125714. <https://doi.org/10.1016/j.limno.2019.125714>

5195

5196 This study investigates the relative impact of major biophysical controls (stream temperature,
5197 riparian litterfall, and stream discharge) on in-stream CPOM (coarse particulate organic matter)
5198 quantity across a variety of streamside timber harvest intensities using simulation modeling. The
5199 CPOM model used was developed by Stenroth et al., 2014, for similar stream types and
5200 conditions of coastal rainforest streams of British Columbia. The model was calibrated using
5201 data from multiple published studies from, primarily the Pacific Northwest region, and several
5202 other North American regions, that quantified stream flow, temperature, and CPOM following
5203 different timber harvest intensities within 4 years of harvest. The model used an estimated
5204 response of low, moderate, high, and very high severity timber harvest for litterfall (-10%, -30%,
5205 -50%, -90%), peak flows (+20%, +40%, +100%, +300%), and stream temperature (+1°C, +2°C,

5206 +4°C, +6 °C). These changes in litterfall, peak flow, and stream temperature were modeled and
5207 analyzed individually and cumulatively to estimate their relative and combined effects on in
5208 stream CPOM standing stocks. Results of the model showed that in general the standing stocks
5209 of CPOM decreased under the independent effects of reduced litterfall and elevated peak flows
5210 and increased with higher stream temperatures. Along the gradient of harvest severities, litterfall
5211 reductions on depleting CPOM standing stocks were at least an order of magnitude greater than
5212 those of elevated peak flows. At low severity, litterfall reductions led to a 13.5% reduction of
5213 CPOM stocks while peak flow increases at high severity harvest only led to a 5% reduction in
5214 CPOM stocks. The magnitude of CPOM changes induced by litterfall reductions was
5215 consistently greater than stream temperature increases, but their differences in magnitude became
5216 smaller at higher levels of disturbance severity. For example, at low severity, stream
5217 temperatures only led to an increase on CPOM stocks by 1.1% while litter fall reductions led to a
5218 reduction of CPOM by 13.5%. However, at the high intensity treatment CPOM stocks changed
5219 by -90.24%, and +72.07% for litterfall, and stream temperature respectively. For scenarios
5220 involving perturbations of multiple model drivers (combined effects), the effect size of
5221 disturbance was significantly negative (indicating significantly lower CPOM standing stocks
5222 than in undisturbed conditions) whenever litterfall reductions reached 50% or above (i.e., high
5223 severity). When litterfall reductions were 30% or below, the effect size of disturbance varied with
5224 the relative changes in peak flows and stream temperature. Only the effects of litterfall-
5225 temperature interactions on CPOM standing stocks were significant ($p < 0.001$). The authors
5226 interpret these results as evidence that litterfall reduction from timber harvest was the strongest
5227 control on in-stream CPOM quantity for 4 years post-harvest. Further, the authors propose that
5228 the decreased activity of CPOM consumers caused by increasing stream temperatures may be
5229 enough to offset the loss of litterfall inputs on CPOM stocks. The caveat of this study is that it
5230 did not include LW dynamics in preserving CPOM post-harvest. As other studies have shown,
5231 harvest can increase in-stream LW, and in-stream LW can act as a catchment for CPOM.

5232

5233 **Drought Frequency**

5234

5235 Wise, 2010

5236

5237 Wise, E. K. (2010). Tree ring record of streamflow and drought in the upper Snake River. *Water*
5238 *Resources Research*, 46(11). <https://doi.org/10.1029/2010WR009282>

5239

5240 This study used newly collected tree-ring data augmented with existing chronologies from sites
5241 at three headwater streams in the Snake River Basin to estimate streamflow patterns for the
5242 1600-2005 time-period. The reconstructed chronologies were tested for significant correlations
5243 with streamflow patterns during the 1911-2005 time period prior to extrapolation. Streamflow
5244 patterns derived from instrumental data and from reconstructed chronologies were compared

5245 with other streamflow reconstructions of three other western rivers in similar climates to
5246 examine synchronicity among the rivers and gain insight into possible climatic controls on
5247 drought episodes. The reconstruction model developed for the analysis explained 62% of the
5248 variance in the instrumental record after adjustment for degrees of freedom. Results showed
5249 evidence that droughts of the recent past are not yet as severe, in terms of overall magnitude, as a
5250 30-year extended period of drought discovered in the mid-1600s. However, in terms of number
5251 of individual years of < 60% mean-flow (i.e., low-flow years), the period from 1977-2001 were
5252 the most severe. Considering the frequency of consecutive drought years, the longest (7-year-
5253 droughts), occurred in the early 17th and 18th centuries. However, the 5-year drought period from
5254 2000-2004 was the second driest period over the 415-year period examined. The author explains
5255 that the area has continued to experience a drought period, but its severity could not be
5256 calculated as it hadn't ended by the time of the study (2010). The correlative analysis of the
5257 chronologies developed for the upper Snake River with other rivers of the West (the upper
5258 Colorado, the Sacramento, and the Verde Rivers) showed mixed results with periods of positive
5259 and negative correlations. The author interprets these results as evidence that drought frequency
5260 in general, in this area appears to be increasing in severity and that mean annual flow appears to
5261 be reducing in the latter half of the 20th and the beginning of the 21st century. The exceptions
5262 being the 1930's dustbowl, and an unusually long dry period in the early 1600s.

5263

5264 **Shade and structure**

5265

5266 Warren et al., 2013

5267

5268 Warren, D. R., Keeton, W. S., Bechtold, H. A., & Rosi-Marshall, E. J. (2013). Comparing
5269 streambed light availability and canopy cover in streams with old-growth versus early-mature
5270 riparian forests in western Oregon. *Aquatic sciences*, 75(4), 547-558. DOI:10.1007/s00027-013-
5271 0299-2

5272

5273 This study investigates the differences in canopy cover and streambed light availability between
5274 paired reaches in old-growth (> 500 years old) and secondary-growth (~40-60 years old) riparian
5275 forests on canopy cover and streambed light exposure in four second order fish-bearing streams
5276 in the H.J. Andrews Experimental Forest. Streams were paired based on reach length and
5277 bankfull width and north (n=2), and south (n=2) facing watersheds. The overall mean percentage
5278 of canopy cover was estimated using a convex spherical densiometer every five meters along the
5279 thalweg of each stream reach. At each point densiometer readings were taken from four
5280 directions (upstream, downstream, left bank, right bank) The amount of light reaching the bottom
5281 of the stream was estimated every five meters using fluorescent dye that degrades overtime from
5282 light exposure. Differences in light availability and canopy cover were analyzed separately for
5283 each of the four reaches using a single factor ANOVA. To avoid the inclusion of overlapping

5284 canopy images from adjacent densiometer sampling locations, the canopy cover data from sites
5285 every 15 m (rather than every 5 m) were used in the comparison of canopy cover between the
5286 two age classes along each reach. Linear regression was used to compare values from mean
5287 densiometer readings with mean dye photodegradation site (every 5 meters). To evaluate the
5288 hypothesis that light availability in old-growth forested streams would be more variable than in
5289 second-growth forested streams, the standard deviations of the mean densiometer readings and
5290 mean photodegradation values were compared between old-growth and second-growth forested
5291 streams with an ANOVA. Results showed that the differences in stream light availability and
5292 percent forest cover between old-growth and second-growth reaches were significant in both of
5293 the south-facing watersheds in mid-summer at an alpha of 0.01 for the dye results and 0.10 for
5294 the cover results. For the north-facing watersheds differences in canopy cover and light
5295 availability (alpha = 0.01, and 0.10 respectively) were only significant at 1 of the two reaches.
5296 Overall, three of the four paired old-growth reaches had significantly lower mean percent canopy
5297 cover, and significantly higher mean decline in fluorescent dye concentrations. The authors
5298 interpret these results as evidence that old-growth forest canopies were more complex and had
5299 more frequent gaps allowing for more light availability and lower mean canopy cover, on
5300 average, than in adjacent mature second-growth forests.

5301

5302 **LW**

5303

5304 Teply et al., 2007

5305

5306 Teply, M., McGreer, D., Schult, D., & Seymour, P. (2007). Simulating the effects of forest
5307 management on large woody debris in streams in northern Idaho. *Western Journal of Applied*
5308 *Forestry*, 22(2), 81–87. <https://doi.org/10.1093/wjaf/22.2.81>

5309

5310 This paper uses simulation modeling to estimate the effects of timber harvest, under the Idaho
5311 Forest Plan (IFP), on in-stream LW loading for Class I streams (fish-bearing streams) of the
5312 Priest Lake Watershed in northern Idaho relative to unharvested riparian forest streams. Under
5313 the IFP, class one streams have a 25-foot no-cut-buffer that extends out from the high-watermark,
5314 and an additional 50 feet beyond the edge of the no-cut-buffer where harvest requires retention of
5315 88-trees-per-acre that are greater than 8-in diameter at breast height (DBH). This study used the
5316 Riparian Aquatic Interaction Simulator (RAIS) to estimate the potential wood loading for 58
5317 randomly selected north Idaho stream segments with and without harvest. Stream segments were
5318 measured in the field along the stream centerline from the upstream starting point (0 ft) to a
5319 downstream ending point (200 ft). Riparian stand conditions were measured within 75 ft-long by
5320 10-ft-wide strips oriented perpendicular to the stream at 25, 75, 125, and 175 ft downstream of
5321 the upstream starting point on each side of the stream segment to provide a total of eight strips
5322 for each stream segment. Along each strip, live trees and snags greater than 8 in dbh within the

5323 strip were located and measured. Three circular subplots, each 10 ft in diameter, were located
5324 along each 75-foot strip plot at 12.5, 37.5, and 62.5 ft from the stream edge. Within the subplots,
5325 smaller live trees (less than 8-in. dbh) were tallied by 1-in. dbh classes. Instream LW loads were
5326 surveyed along the same 200-ft stream segments located for measuring riparian stand conditions.
5327 Qualifying LW (greater than 4-in diameter and longer than 6.6 ft) occurring within the high-
5328 water mark along the entire extent of the segment was tallied. Observed instream LW loads
5329 ranged from 10 to 710 pieces per 1,000 ft of stream. Stream size measured by bank full width
5330 covered a wide range (1 ft to 190 ft), averaging 32.5 ft (SD = 28.1). The authors determined that
5331 active streambank erosion was uncommon in the study area and did not include it as a LW
5332 recruitment mechanism in their analysis. Simulation was based on a four-step process applied to
5333 each riparian stand: 1) Harvest the stand according to riparian management prescriptions, 2)
5334 Predict stand characteristics using growth and yield simulators, 3) Estimate the number of trees
5335 that fall due to mortality in each time step, 4) Calculate the probability that a tree would deliver
5336 LWD to the stream. The simulation evaluated both a harvest and a no-harvest scenario to predict
5337 mean in-stream LW loads after 30, 60, and 100 years. The results predicted mean LW loads at 30
5338 years for the 58 segments studied were 151.1 pieces per 1,000 ft for the no-harvest scenario (SD
5339 = 76.2) and 145.1 pieces per 1,000 ft for the harvest scenario (SD = 75.6), which were not
5340 significantly different ($P = 0.67$). However, on a pairwise basis, loads predicted for these
5341 segments using the harvest scenario were significantly lower by an average of about 6.0 pieces
5342 per 1,000 ft than those predicted via the no-harvest scenario ($P < 0.001$). Compared to the initial
5343 surveyed LW loads, LW loads at 30 years predicted in the no-harvest scenario decreased by an
5344 average of 19.5 pieces per 1,000 ft, representing a significant ($P < 0.007$) downward shift in the
5345 distribution. Predicted mean LW loads at 60 years were 136.1 pieces per 1,000 ft in the no-
5346 harvest scenario (SD = 49.2) and 128.3 pieces per 1,000 ft under the harvest scenario (SD =
5347 48.3). At 100 years, predicted mean LW loads were 122.5 (SD = 35.4) and 116.7 (SD = 35.8),
5348 respectively. Based on 20-piece LW classes, the frequency distributions of predicted loads
5349 between the scenarios were not significantly different at either time step. However, on a pairwise
5350 basis, predicted loads for the harvest scenario were significantly lower than the no-harvest
5351 scenario by an average of 7.8 ($P < 0.001$) and 5.8 ($P < 0.001$) pieces per 1,000 ft at 60 years and
5352 100 years, respectively. Compared to LW loads predicted at 30 years and 60 years, LWD loads
5353 decreased significantly on a pairwise basis by an average of 15.1 ($P < 0.001$) and 13.6 ($P <$
5354 0.001) at 60 and 100 years, respectively. The authors note that the collective effect of the
5355 assumptions made for the simulation is likely to underestimate the number and variability of LW
5356 pieces recruited and retained in the streams sampled. The authors interpreted these results as
5357 evidence that the IFP prescriptions for class I Idaho streams were sufficient in maintaining LW
5358 recruitment potential.

5359

5360 **Shade**

5361

5362 Swartz et al., 2020

5363

5364 Swartz, A., Roon, D., Reiter, M., & Warren, D. (2020). Stream temperature responses to
5365 experimental riparian canopy gaps along forested headwaters in western Oregon. *Forest Ecology*
5366 *and Management*, 474, 118354. <https://doi.org/10.1016/j.foreco.2020.118354>

5367

5368 This study tested the effects of adding canopy gaps within young, regenerating forests of western
5369 Oregon on stream light availability and stream temperatures. The addition of gaps in the young
5370 regenerating forests were used to theoretically mimic the natural disturbance regimes and the
5371 higher canopy complexity of late-successional forests. The researchers used a before-after-
5372 control-impact design on six replicated streams within the Mckenzie River Basin. In the
5373 experimental reaches 30 m gaps were created, centered on a tree next to the stream and at least
5374 30 m in from the beginning of the reach. The study reaches were located on second- and third-
5375 order fish-bearing steep step-pool and cascade dominated headwater streams with boulder
5376 substrate that ranged from 2.2 to 6.4 m in bankfull width and were lined by 40- to 60-year-old
5377 riparian forests. Study sites in each stream encompassed two 120 m reaches with no large
5378 tributary inputs within or between the study reaches, and reference and treatment reaches were
5379 separated by a buffer section of 30–150 m. In each treatment reach, gaps were designed to create
5380 openings in the canopy that were approximately 20 m in diameter. Gaps were centered on a tree
5381 next to the stream at approximately meter 30 along each reach. The gaps sizes were intended to
5382 mimic naturally occurring gaps from an individual large tree mortality or small-scale disturbance
5383 events found in these systems which range from 0.05 to 1.0 gap diameter to tree height ratio with
5384 smaller gaps occurring more frequently. Using the Douglas-fir canopy height of 50 m, gaps were
5385 created in the 0.4–1.0 gap diameter to tree height ratio range (approximately 314 m² – 1,963
5386 m²). Actual gap sizes varied across sites from approximately 514 m² to 1,374 m² (0.45 – 0.74
5387 gap ratios) with a mean of 962 m² (mean gap ratio 0.61). Riparian shade was quantified with
5388 hemispherical photos. Light reaching the stream was quantified using photodegradation of
5389 fluorescent dyes placed at 5 m intervals, over a 24 -hour period. Stream temperature was
5390 recorded continuously, at 15-minute intervals, using HOBO sensors to quantify the seven-day
5391 moving average of mean and maximum temperatures. Data was collected for one year pre-
5392 harvest, during harvest year (harvest took place in late fall 2017), and one-year post-harvest. To
5393 determine the effects of experimental canopy gaps on stream light as well as reach responses a
5394 linear mixed-effects model was fit to the data. The results showed that after gaps were cut, the
5395 BACI analysis showed strong evidence for significant increase in mean reach light ($p < 0.01$) to
5396 a mean of 3.91 (SD \pm 1.63) moles of photons m⁻² day⁻¹. overall resulting in a mean change in
5397 light of 2.93 (SD \pm 1.50) moles of photons m⁻² day⁻¹. Mean stream shading could not be
5398 evaluated in the full BACI analysis because post-treatment hemispherical photographs could not
5399 be taken at all sites due to fire impeding access in 2018. For the remaining sites, the areas
5400 beneath each gap had notable localized declines in shade, through the entirety of the treatment
5401 reach mean shading declined by only 4% (SD \pm 0.02%). Overall, the gap treatments did not
5402 change summer T 7DayMax or T 7DayMean significantly across the 6 study sites. The mean
5403 response (change in reach difference before and after the cut) indicated an increase on average
5404 across the six sites in T7DayMax of 0.21 °C (\pm 0.12 °C) and in the T7DayMean of 0.15 °C (\pm 0.14
5405 °C); however, there was not statistical support of the BACI effect for either metric. The light

5406 response was not correlated with T 7DayMax responses ($r^2 < 0.01$, $p = 0.69$), nor was gap area
5407 ($r^2 = 0.01$, $p = 0.63$), but there was a significant relationship between discharge ($r^2 = 0.73$, $p =$
5408 0.03), and bankfull width ($r^2 = 0.93$, $p < 0.01$) and the T7DayMax response. Wetted width was
5409 also highly correlated with T 7DayMax responses, but the relationship was not as strong with
5410 this stream size metric as with discharge or bankfull width ($r^2 = 0.65$, $p = 0.05$). In contrast to the
5411 summary values, results from the analysis of individual days throughout the full 40-day summer
5412 period identifying differences in the relationships of daily maximums and daily means between
5413 reaches showed a statistically significant effect of the gap for average daily maximums ($p < 0.01$)
5414 and for average daily means ($p = 0.02$). The regression comparison reveals there will be on
5415 average an additional $0.12\text{ }^{\circ}\text{C}/^{\circ}\text{C}$ increase in daily maximum temperature in the reach with a gap.
5416 Likewise, for the daily mean, for every degree increase in the shaded reference reach, an average
5417 additional increase of $0.05\text{ }^{\circ}\text{C}$ in a reach with a small gap is expected. The authors conclude that
5418 adding gaps to young regenerating forests only minimally increases temperatures, dependent on
5419 stream size, and that riparian canopy gaps may be a viable management strategy that can be
5420 implemented with minimal effects on stream temperatures. This paper does not quantify changes
5421 in stream productivity, also expected from the increase in available light.

5422

5423 **Shade**

5424

5425 Sugden et al., 2019

5426

5427 Sugden, B. D., Steiner, R., & Jones, J. E. (2019). Streamside management zone effectiveness for
5428 water temperature control in Western Montana. *International Journal of Forest Engineering*,
5429 *30*(2), 87-98. <https://doi.org/10.1080/14942119.2019.1571472>

5430

5431 This study investigates the effects of riparian forest timber harvest, under the Montana
5432 Streamside Management Zone (SMZ) laws, on stream temperature in Class 1 streams (fish-
5433 bearing, or flow more than 6 months per year and are connected to downstream waters).
5434 Montana state law requires timber be retained within a minimum of 15.2 m of the class 1
5435 streams, with equipment exclusion zones extended on steep slopes for up to 30.5 m. Within the
5436 SMZ no more than half the trees greater than 204 mm (8 in) diameter at breast height (DBH) can
5437 be removed, and trees retained must be representative of the pre-harvest stand. In no case,
5438 however, can stocking levels of leave trees be reduced to less than 217 trees per hectare. The
5439 objectives of this study were to fill the information gap in this region by: (1) evaluating the
5440 performance of 15.2 m SMZs retained during harvest activities for protecting against adverse
5441 changes in summer maximum stream temperatures, (2) quantifying the level of timber removal
5442 occurring within operational SMZs that may help explain any observed changes, and (3)
5443 Evaluating fish response that may be associated with a stream temperature change. Data for
5444 stream temperature and fish population response was collected for 30 harvest reaches in western

5445 Montana (northern Rocky Mountain Region), for a minimum of one-year pre- and one-year post-
5446 harvest. Data for stream temperatures and fish populations were also collected from unharvested
5447 reference reaches upstream from the harvest sites as a control. Temperature data was collected
5448 with Optic StowAway™ and StowAway TidBit™ digital temperature loggers manufactured by
5449 Onset Computer Corporation. Shade over the stream surface was not directly measured in this
5450 study. Canopy cover was estimated using a combination of simulation modeling and using a
5451 concave spherical densiometer. Fish populations were estimated for 100 m reaches at study sites
5452 using an electro-fishing pass of capture method. Linear mixed effects models were used to
5453 analyze the relationship between year, stream position, harvest, fish populations and stream
5454 temperatures. The results showed that within harvest areas, the mean basal area (BA) declined
5455 from 30.2 m²/ha pre-harvest to 26.4 m²/ha post-harvest (mean = -13%, range from -32% to
5456 0%). Windthrow further reduced the mean BA to 25.9 m²/ha (mean = -2%, range = -32% -0%).
5457 Changes in mean canopy cover were not significant based on the simulation modeling (-3%), or
5458 densiometer readings (+1%). Results of the model for the effect of harvest on stream
5459 temperature showed no detectable increase in treatment streams relative to control streams. The
5460 estimated mean site level response in maximum weekly maximum temperatures (MWMT)
5461 varied from -2.1 °C to +3.3 °C. Overall, 20 of 30 sites had estimated site level response within
5462 ±0.5 °C. There were five sites that had an estimated site-level response greater than 0.5 °C (i.e.
5463 warming) and five sites that had an estimated site level response less than -0.5 °C (i.e. cooling).
5464 Results for the fish population showed approximately 7% increase in trout population from pre-
5465 harvest to post-harvest, but this difference was not significant. The authors conclude that the
5466 results suggest that Montana's 15.2 m SMZs retained during timber harvest activities are highly
5467 protective (change <0.5°C) of stream temperatures.

5468

5469 **LW**

5470

5471 Sobota et al., 2006

5472

5473 Sobota, D. J., Gregory, S. V., & Sickle, J. V. (2006). Riparian tree fall directionality and
5474 modeling large wood recruitment to streams. *Canadian Journal of Forest Research*, 36(5), 1243–
5475 1254. <https://doi.org/10.1139/x06-022>

5476

5477 The objectives of this study were to evaluate patterns of riparian tree fall directions in diverse
5478 environmental conditions and evaluate correlations with tree characteristics, forest structural
5479 variables, and topographic features. Specifically, the authors were interested in correlations
5480 between fall directionality and tree species type, tree size, riparian forest structure, and valley
5481 topography (side slope). Data was collected from 21 field sites located west of the Cascade
5482 Mountains crest (11 sites: Coast Range and west slopes of the Cascades), and in the interior
5483 Columbia Basin (10 sites: east slopes of the Cascades, Blue Mountains, and Northern Rockies)

5484 of Oregon, Washington, Idaho, and Montana, USA. Streams were second- to fourth-order
5485 channels and had riparian forests that were approximately 40 to >200 years old. The location of
5486 specific study reaches (200–300 m stream length) on each stream were selected randomly.
5487 Minimum size criteria for a fallen tree in this study were diameter at breast height (DBH) of 0.1
5488 m and height of 5 m. All fallen trees up to 50 m slope distance from stream or the first 100 trees
5489 were measured at all sites. Tree fall direction was standardized among sites by streamside
5490 location (upstream = 0° and 360°; toward stream = 90°; downstream = 180°; away from stream =
5491 –90° and 270°). Spearman rank correlations were used to compare site level statistics of tree fall
5492 directions with physical and riparian forest characteristics. Then trees were pooled among sites
5493 and classified by species for analysis of species, tree size, and valley side slope effects. To avoid
5494 small sample sizes species were grouped by side slope categories (< 40%, >40%). Average
5495 direction of tree fall by site was significantly correlated with valley constraint (Spearman $r = -$
5496 0.53; $P = 0.02$). Average direction of tree fall by site was weakly correlated with active channel
5497 width, tree stem density, and basal area ($P > 0.05$), with Spearman r coefficients of 0.22, –0.21,
5498 and 0.39, respectively. Trees on valley side slopes >40% for each species had a 95% CI that only
5499 included falls directly towards the stream channel; trees on side slopes <40% had a 95% CI for
5500 mean fall direction that included directly upstream, downstream, away from the stream, towards
5501 the stream, or all four directions simultaneously (consistent with random fall directions),
5502 depending on species. Tree size was only different between side slope categories for coastal
5503 Douglas fir on >40% side slopes which had a median DBH 1.2 to 1.9 times greater than trees on
5504 <40% side slopes. Also, red alder trees on side slopes > 40% had a median DBH 1.1 to 1.6 times
5505 greater than on side slopes < 40%. Model projections of LW recruitment calibrated with the
5506 results of the spearman rank correlations estimated that sites with uniform steep side slopes
5507 (>40%) produced between 1.5 (first resolution) to 2.4 (second resolution) times more in stream
5508 LW by number of tree boles than sites with uniform moderate side slopes (< 40%). The authors
5509 interpret their results as evidence that edaphic, topographic, and hydrologic characteristics are
5510 related to greater variability of tree fall directions on moderate slopes than on steep slopes. The
5511 authors conclude that models that use tree fall directions in predictions of LW recruitment should
5512 consider stream valley topography. The authors warn that while side slope categories (>40%,
5513 <40%) was the strongest predictor of tree fall direction in this study, they believe the differences
5514 in tree fall direction between these categories mainly characterized differences between fluvial
5515 (88% of moderate slope sites) and hillslope landforms (71% of steep slope sites). They suggest
5516 that the Implications from this study are most applicable to small- to medium-size streams
5517 (second- to fourth-order) in mountainous regions where sustained large wood recruitment from
5518 riparian forest mortality is the significant management concern.

5519

5520 **LW**

5521

5522 Schuett-Hames & Stewart, 2019a

5523

5524 Schuett-Hames, D., & Stewart, G. (2019a). Post-Harvest Change in Stand Structure, Tree
5525 Mortality and Tree Fall in Eastern Washington Riparian Buffers: Comparison of the Standard
5526 and All Available Shade Rules for the Fish-Bearing Streams in the Mixed Conifer Timber Habitat
5527 Type Under Washington's Forest Practices Habitat Conservation Plan. Cooperative Monitoring
5528 Evaluation and Research Report CMER. Washington State Forest Practices Adaptive
5529 Management Program. Washington Department of Natural Resources, Olympia, WA.

5530

5531 This report is a comparative analysis of the differences in stand structure, tree fall, and LW
5532 recruitment between riparian sites of eastern Washington harvested under the current Standard
5533 Shade Rule (SR), under the All-Available Shade rule (AAS), and unharvested reference sites
5534 (REF). Both shade rules have a 30-ft no-cut buffer (core zone) immediately adjacent to the
5535 stream. The SR prescription allows thinning in the buffer zone 30-75 feet (inner zone) from the
5536 stream while the AAS prescription requires retention of all shade providing trees in this area.
5537 Post-harvest surveys were completed at each site one–two years and five years post-harvest. A
5538 census was done of all standing trees ≥ 4 inches diameter at breast height (DBH) within 75 feet
5539 (horizontal distance) of the channel on both sides of the stream in each treatment and reference
5540 reach. The condition (live or dead), species, canopy class, and DBH were recorded for each tree.
5541 Dead or fallen trees with a decay class of 1 or 2 were classified as post-harvest mortality and a
5542 mortality agent was recorded (e.g. wind, erosion, suppression, fire, insects, disease, and physical
5543 damage). Metrics were calculated separately for regulatory zones defined by horizontal distance
5544 from the channel, including the core zone (0–30 feet) and inner zone (30–75 feet) and the
5545 combined core and inner zone (the full RMZ). Mixed model analysis was used to evaluate
5546 differences in treatment response. Results showed Cumulative wood recruitment from tree fall
5547 over the five-year post-harvest interval was highest in the SR group, lower in the AAS group and
5548 lowest in the REF group. The SR and AAS rates by volume were nearly 300% and 50% higher
5549 than the REF rates, respectively. The mixed model comparisons indicated that the frequency of
5550 wood input from fallen trees was significantly greater in SR group compared to both the REF
5551 and AAS groups ($p < 0.001$), while the difference between REF and AAS groups was not
5552 significant. Over 60% of pieces recruited from AAS and SR fallen trees consisted of stems with
5553 attached rootwads (SWAR), double the proportion in the REF sites. The REF-AAS and REF-SR
5554 differences in recruitment of SWAR pieces were significant ($p < 0.001$). Most recruiting fallen
5555 trees originated in the core zone (76%, 72%, and 64% for the REF, AAS and SR groups,
5556 respectively), while the proportion from the inner zone (30–75 feet from the stream) was ~10%
5557 greater for the SR group compared to the AAS and REF groups. The authors interpret the results
5558 and conclude that harvest of the adjacent stand outside the RMZ appeared to alter the spatial
5559 pattern of wood recruitment from fallen trees, increasing recruitment from trees located farther
5560 from the stream. Recruitment of fallen trees from the inner zone of the AAS and SR sites were
5561 two and four times the rate for the inner zones of the unharvested reference sites due to increased
5562 tree fall from wind disturbance in the buffers after harvest of the adjacent stand, as reported in
5563 other studies. It is important to note that this was a short-term study (5 years). The authors note
5564 that LW recruitment is a process that can change over decadal time scales. Adding that thinning

5565 and post-harvest mortality also reduced the standing stock of trees available for wood
5566 recruitment in the SR and AAS RMZs compared to unharvested REF RMZs.

5567

5568 **Litter and LW**

5569

5570 Six et al., 2022

5571

5572 Six, L. J., Bilby, R. E., Reiter, M., James, P., & Villarin, L. (2022). Effects of current forest
5573 practices on organic matter dynamics in headwater streams at the Trask River watershed,
5574 Oregon. *Trees, Forests and People*, 8, 100233. <https://doi.org/10.1016/j.tfp.2022.100233>

5575

5576 This study investigates the effects of different riparian timber harvest intensities on changes in
5577 canopy cover, and litter input into streams and litter transport downstream. The objective of this
5578 study was to investigate whether differing levels of tree retention adjacent to the channel altered
5579 coarse particulate organic matter (CPOM) delivery, retention, and transport. The authors
5580 hypothesized an inverse relationship between tree removal and litter delivery (i.e., increase in
5581 tree removal adjacent to the channel would result in a reduction of litter delivery). Data was
5582 collected for leaf litter in streamside litter traps, canopy cover percentage using hemispherical
5583 photos in-stream LW, and litter retention in stream flume litter traps pre- and post-treatment at
5584 five watersheds of the Trask River in the northern Oregon Coast range. The experimental design
5585 included three treatment watersheds: clearcut with no leave trees or retention buffer (CC),
5586 clearcut with leave trees (CC w/LT; retention of 5 trees per hectare/2 trees per acre), and clearcut
5587 with 15 m wide retention buffer (CC c/B) and two uncut references (REF 1, and 2) along
5588 headwater streams. Because there were no replication sites for treatments, data was analyzed
5589 using descriptive and graphical summaries of the data (i.e., no quantitative statistical analysis).
5590 Results showed a reduction of canopy cover from 91.4% to 34.4% in the clearcut treatment with
5591 no leave trees, from 89.8% to 76.1% in the clearcut treatment with leave trees, and from 89.5%
5592 to 86.9% in the clearcut treatment with the 15 m retention buffer. Change in canopy cover in the
5593 reference streams was < 1% for both reaches. Post harvest litter delivery decreased for the
5594 clearcut with no leave trees but increased for both the clearcut with leave tree and clear cut with
5595 retention buffer. The number of logjams, the total weight of logjams, and the volume of LW in
5596 streams increased for all treatment sites. The results of this study were consistent with similar
5597 studies and provide supporting evidence that riparian timber harvest can affect litter and LW
5598 delivery into and retention in streams.

5599

5600 **Shade and LW**

5601

5602 Schuett-Hames et al., 2011

5603

5604 Dave Schuett-Hames, Ashley Roorbach, Robert Conrad. 2011. Results of the Westside Type N
5605 Buffer Characteristics, Integrity and Function Study Final Report. Cooperative Monitoring
5606 Evaluation and Research Report, CMER 12-1201. Washington Department of Natural Resources,
5607 Olympia, WA.

5608

5609 This report presents the results from the Washington State Westside Type N Buffer
5610 Characteristics, Integrity and Function (BCIF) study. The purpose of the study was to evaluate
5611 the effects of westside riparian timber harvest prescriptions for Type Np (perennial non-fish-
5612 bearing) streams on resource objectives (riparian stand tree mortality, wood recruitment, channel
5613 debris, shade, and soil disturbance) described in the Forest and Fish Report of 1999. Three
5614 treatment prescriptions were evaluated, 1) clearcut harvest to the edge of the stream (CC) at eight
5615 sites, 50-foot-wide no-cut-buffers (50-ft) at 13 sites, and 56-foot radius circular no-cut-buffer at
5616 the perennial initiation point (PIP) at three sites (not used in statistical analysis due to small
5617 sample sizes). Each treatment site was paired with an uncut reference site as a control. The CC
5618 and 50-ft treatments were compared with treatment sites at three time periods (the first 1-3 years,
5619 years 4-5, and the whole 5-year period). Differences in variable mean values were checked for
5620 statistical significance between treatment and reference streams using non-parametric Mann-
5621 Whitney U tests. Tree fall rates (annual fall rates of live and dead standing stems combined) was
5622 over 8 times and 5 times higher in the 50-foot buffers than in the reference buffers 3 years after
5623 treatment when compared as a percentage of standing trees and as trees/acre/yr, respectively.
5624 These differences were significant for both metrics ($p \leq 0.001$). In the period 4-5 years post
5625 treatment rate of tree uprooting decreased but rate of stem breakage increased in the 50-foot
5626 buffer. For this period only the percentage of broken trees were significantly different (higher)
5627 than what was observed in the reference buffers. Over the entire five-year period, the percentages
5628 of standing trees that were uprooted and broken (as well as the combined total) were
5629 significantly greater in the 50-foot buffer. Wind was the dominant tree fall process, accounting
5630 for nearly 75% of combined fallen trees, 11% fell from other trees falling against them and 1.8%
5631 of fallen trees fell from bank erosion. Differences in mortality followed a similar pattern to tree
5632 fall rates. In the 50-foot buffer sites mortality rates were significantly higher (3.5 times higher)
5633 than in the reference sites for the first three years following harvest. However, in years 4-5
5634 mortality rates increased in the reference buffers after high-intensity storms resulting in non-
5635 significant differences in mortality during this period. The cumulative percentage of live trees
5636 that died over the entire five-year period was 27.3% in the 50-ft buffers compared to 13.6% in
5637 the reference reaches, but the difference was not statistically significant. This was likely because
5638 of the high variability in mortality between sites in the 50-foot buffers. LW recruitment into the
5639 channel after treatment was higher in the 50-ft buffers than in the reference patches during the
5640 first three years after harvest, over 8 times higher in pieces/acre/yr and over 14 times higher in
5641 volume/acre/yr. In years 4-5 after harvest LW recruitment decreased in the 50-ft buffers and
5642 increased in the reference patches, and the number of recruited LW pieces/acre/yr was greater in

5643 the reference patches, although the volume of LW recruited was greater in the 50-ft buffers. For
5644 the entire first 5 years after harvest, the 50-ft buffers recruited about twice the number of LW
5645 pieces recruited in the reference patches, and over 3 times the volume. The CC treatment,
5646 unsurprisingly, had significantly lower LW recruitment following harvest relative to the reference
5647 streams. Mean overhead shade (from trees and tall shrubs) was 13% lower in the 50-ft treatment,
5648 and 77% lower in the CC treatment relative to reference streams. The CC treatment, however,
5649 increased by 25% five years after harvest relative to values recorded 1-year following harvest.
5650 The implications of these results suggest that immediate and direct changes in stand structure,
5651 canopy cover, and LW are most severe for clear-cut treatments, but that the 50-foot buffer
5652 treatment showed an increase in LW and stand mortality, and a decrease in shade over the five-
5653 year period. Limitations of this study were the lack of pre-harvest data and the relatively short
5654 time-period (5-years) in evaluating impacts that may last for several decades.

5655

5656 Schuett-Hames & Stewart, 2019b (BCIF)

5657

5658 Schuett-Hames, D & Stewart, G. (BCIF), (2019). Changes in stand structure, buffer tree
5659 mortality and riparian-associated functions 10 years after timber harvest adjacent to non-fish-
5660 bearing perennial streams in western Washington. Cooperative Monitoring Evaluation and
5661 Research Report. Washington State Forest Practices Adaptive Management Program. Washington
5662 Department of Natural Resources, Olympia, WA.

5663

5664 This paper presents a 10 -year follow-up to the results of the BCIF report (Schuett-Hames et al.,
5665 2012) that originally presented 5-year post-treatment results. Over the 10-year period stand
5666 mortality in the 50-ft buffer treatment stabilized and showed a cumulative 14.1% reduction in
5667 live basal, while the reference stands showed a 2.7% increase in live basal area. The differences
5668 in these values were not significant. Cumulative LW recruited into stream channel over the 10-
5669 period was double in the 50-ft treatment streams than in the reference streams. However, the
5670 majority of the LW recruited in the 50-ft treatment streams came to rest above the streams,
5671 providing shade but not affecting streamflow, pool formation, or sediment storage. Further, while
5672 the 50-ft buffer treatment provided more LW recruitment in the short-term (10-years), the authors
5673 speculate there is a reduction in future LW recruitment potential given the removal of trees
5674 outside the 50-ft buffer. Canopy cover in the 50-ft treatment streams recovered to similar
5675 percentages as the reference's streams by the end of the 10-year period. The authors speculate
5676 that the 50-ft buffer was better at maintaining resource objectives than the clearcut but propose
5677 that the narrow buffers presented variable increases in mortality (specifically increased
5678 susceptibility to windthrow) and recommend further research before drawing definitive
5679 conclusions.

5680

5681 **Riparian thinning effects on shade, light, and temperature**

5682
5683 Roon et al., 2021a
5684
5685 Roon, D.A., Dunham, J.B., Groom, J.D., 2021. Shade, light, and stream temperature responses to
5686 riparian thinning in second-growth redwood forests of northern California. PLOS ONE 16,
5687 e0246822. <https://doi.org/10.1371/journal.pone.0246822>

5688
5689 The purpose of this study was to evaluate the effects of riparian thinning on shade, light, and
5690 temperature in three watersheds located in second-growth redwood stands in northern California.
5691 The objectives of this study were to evaluate: 1) the effects of experimental riparian thinning
5692 treatments on shade and light conditions; 2) how changes in shade and light associated with
5693 thinning affected stream temperatures at a reach-scale both locally and downstream; 3) how
5694 thermal responses varied seasonally; and 4) how these thermal responses were expressed across
5695 the broader thermal regime to gain a more complete understanding of thinning on stream
5696 temperatures in these watersheds. This study took place between 2016 and 2018 with thinning
5697 treatments applied during 2017 giving 1-year pre-treatment and 1-year of post-treatment data.
5698 Two study sites prescribed treatment on one side of the stream of a 45 m buffer width with a 22.5
5699 m inner zone with 85% canopy retention and a 22.5 m outer zone that retained 70% canopy
5700 cover (Green Diamond Resource Company, Tectah watershed). At the third treatment site
5701 thinning prescriptions included removal of up to 40% of the basal area within the riparian zone
5702 on slopes less than 20% on both sides of the channel along a ~100–150 m reach (Lost Man
5703 watershed, Redwood national park). Control reaches were located upstream from treatment
5704 reaches. Data analysis was conducted separately for each experimental watershed (i.e., 1 Lost
5705 man site, 2 Tectah sites). Stream temperature was collected using digital sensors; solar radiation
5706 was measured using silicon pyranometers; riparian shade was measured using hemispherical
5707 photography. A classical BACI analysis was performed to test the effects of riparian thinning on
5708 shade, light, and stream temperature using linear-effects models. Results for the Tectah
5709 watershed showed a significant reduction in canopy closure by a mean of 18.7%, (95% CI: -21.0,
5710 -16.3) and a significant reduction of effective shade by a mean of 23.0% (-25.8, -20.1) one-year
5711 post treatment. In the Lost man watershed, a non-significant reduction of mean shade by 4.1% (-
5712 8.0, -0.5), and mean canopy closure by 1.9% was observed in 2018. Results for below canopy
5713 light availability showed significant increases by a mean of 33% (27.3, 38.5) in the Tectah
5714 watershed, and non-significant increases in Lost man watershed of 2.5% (-1.6, 5.6) by 2018.
5715 Results for stream temperature changes showed variation seasonally and between watersheds.
5716 The Lost Man watershed showed no significant changes in average daily maximum, maximum
5717 weekly average of the maximum (MWMT), average daily mean, or maximum weekly average of
5718 the mean (MWAT). In the Tectah watershed, MWMT increased during spring by a mean of 1.7°C
5719 (95% CI: 0.9, 2.5), summer by a mean of 2.8°C (1.8, 3.8), and fall by a mean of 1.0°C (0.5, 1.5)
5720 and increased in downstream reaches during spring by a mean of 1.0°C (0.0, 2.0) and summer by
5721 a mean of 1.4°C (0.3, 2.6). Thermal variability of streams in the Tectah watershed were most
5722 pronounced during summer increasing the daily range by a mean of 2.5°C (95% CI:

5723 1.6, 3.4) and variance by a mean of 1.6°C (0.7, 2.5), but also increased during spring (daily
5724 range: 0.5°C; variance: 0.3°C) and fall (daily range: 0.4°C; variance: 0.1°C). Increases in thermal
5725 variability in downstream reaches were limited to summer (daily range: 0.7°C; variance: 0.5°C).
5726 Again, no significant changes in stream and downstream temperature variability were detected in
5727 the Lost Man watershed. In the Techtah watersheds the frequency of days with temperatures
5728 greater than 16°C increased in summer by a mean of 42.9 more days (95% CI: 31.5, 53.8) in
5729 thinned reaches and a mean of 16.3 more days (6.1, 27.4) in downstream reaches. Temperatures
5730 greater than 16°C persisted for a mean duration of 31.1 more consecutive days (21.0, 41.1) in
5731 thinned reaches and 11.6 more consecutive days (3.9, 20.0) in downstream reaches under the
5732 BACI analysis. The authors conclude that responses to the experimental riparian thinning
5733 treatments we evaluated differed greatly depending on treatment intensity. For example, they
5734 interpret their results as evidence that that changes in shade of 5% or less caused minimal
5735 changes in temperature while reductions in shade of 20–30% resulted in much larger increases in
5736 temperature. However, the authors warn that their data only evaluated immediate (1-year-post-
5737 treatment) changes in stream shade and temperatures. Also, the study was conducted in relatively
5738 small (< 10 km²) coastal watersheds and may not apply to larger watersheds of different regions.

5739

5740 **Sediment**

5741

5742 Safeeq et al., 2020

5743

5744 Safeeq, M., Grant, G.E., Lewis, S.L., Hayes, S.K., 2020. Disentangling effects of forest harvest
5745 on long-term hydrologic and sediment dynamics, western Cascades, Oregon. *Journal of*
5746 *Hydrology* 580, 124259. <https://doi.org/10.1016/j.jhydrol.2019.124259>

5747

5748 The purpose of this study was to separate and investigate the effects of changes in streamflow
5749 and sediment supply due to disturbances (specifically timber harvest), on sediment transport into
5750 streams. Timber harvest affects both streamflow and sediment supply simultaneously. The
5751 researchers used a reverse regression technique to evaluate the relative and absolute importance
5752 of changes in streamflow versus changes in sediment supply on sediment transport. The
5753 technique was applied to long-term data collected from two paired experimental watersheds in
5754 the H.J. Andrews Experimental Forest, Oregon. The two watersheds were paired by size, aspect,
5755 and topography. The treatment watershed was 100% clearcut during the period from 1962-1966,
5756 broadcast burned in 1966, and re-seeded in 1968. Streamflow, and sediment data were taken
5757 intermittently, and after large storm events from 1952 (pre-harvest) through 1988 for suspended
5758 sediment data, and 2016 for sediment bedload. The control watershed was forested, and had no
5759 treatments (e.g., harvest) during the study period. The results that considered the effects of
5760 harvest on streamflow alone showed an increase in annual water yield in the treatment watershed

5761 by 10% (136 mm/year) over the 51-year post-treatment period. There were no significant
5762 changes in precipitation patterns before or after harvest. Further, the patterns of streamflow in the
5763 control watershed showed diverging patterns in streamflow after the harvest period. The authors
5764 state that these patterns strongly suggest that the increase in streamflow in the treatment
5765 watershed was caused by timber harvest. The results for post-treatment sediment yields showed
5766 suspended load declined to pre-treatment levels in the first two decades following treatment,
5767 bedload remained elevated, causing the bedload proportion of the total load to increase through
5768 time. Changes in streamflow alone account for 477 Mg/km² (10%) of the suspended load and
5769 113 Mg/km² (5%) of the bedload over the post-treatment period. Increase in suspended sediment
5770 yield due to increase in sediment supply is 84% of the measured post-treatment total suspended
5771 sediment yield. In terms of bedload, 93% of the total measured bedload yield during the
5772 posttreatment period can be attributed to an increase in sediment supply. The authors interpret
5773 these results as evidence that while streamflow alone can cause a modest increase in sediment
5774 transport, it is negligible compared to the increases in sediment transport following harvest.
5775 Following harvest, changes on streamflow alone was estimated in being responsible for < 10% of
5776 the resulting suspended sediment transported into streams, while the increase in sediment supply
5777 due to harvest disturbance was responsible for >90%. The authors suggest these results provide
5778 evidence for a need to investigate thresholds for specific watershed management regimes to
5779 ameliorate these impacts following harvest, or thinning treatments. Also, the sharp increases in
5780 sediment transport following logging can be confidently attributed to the increase in sediment
5781 supply and delivery to streams due to the ground disturbances associated with logging rather than
5782 increased streamflow.

5783

5784 **Stream Temperature**

5785

5786 Reiter et al., 2020

5787

5788 Reiter, M., Johnson, S. L., Homyack, J., Jones, J. E., & James, P. L. (2020). Summer stream
5789 temperature changes following forest harvest in the headwaters of the Trask River watershed,
5790 Oregon Coast Range. *Ecohydrology*, 13(3), e2178. <https://doi.org/10.1002/eco.2178>

5791

5792 This paper investigates the effects of different riparian forest harvest treatments on stream
5793 temperature. Stream temperature data was collected from 2006 to 2016 for multiple small (<50
5794 ha), non-fish-bearing headwater stream watersheds in the Trask River Watershed of the
5795 northwestern Oregon Coast range. The experiment followed a BACI design with four treatments,
5796 1) clearcut, no buffer (CC_NB; n = 4), 2) clearcut with 10-m no cut buffer (CC_B; n = 3), 3)
5797 Thinning with 10 m no-cut buffer (TH_B; n = 1), and 4) unharvested, reference streams (REF; n
5798 = 7). Temperature data was collected at 30-minute increments for all streams using continuously
5799 recording thermistors. Harvest operations occurred in the Summer of 2012 giving 6 summers of

5800 pre-treatment and 4 summers of post-treatment data collection. Temperature data was separated
5801 into 5th, 25th, 50th, 75th, and 95th percentiles, with each percentile being treated as independent
5802 response variables in a linear mixed model. Treatments were compared to reference watersheds
5803 to check for significant differences in temperature percentiles. For ecological context, the
5804 researchers also quantified the percentage of summer where temperatures were above 16 and 15
5805 °C, the preferred thermal regime limits for two local amphibian larvae (coastal tailed frog,
5806 coastal giant salamander). Results showed that even the small (10 m buffer; CC_B, TH_B) buffer
5807 was efficient in maintaining similar temperature changes throughout the summers compared to
5808 reference streams. There were no significant changes in the buffered watersheds with
5809 temperature responses in these watersheds ranging from negative values to negative values close
5810 to zero. The treatments with no buffer (CC_NB), however, showed significant increases in
5811 temperature for all percentiles with the greatest increases occurring in the 95th percentile,
5812 showing a mean increase of 3.6 °C (SE = 0.4). For the 5th percentile, the CC_NB also showed a
5813 mean temperature response 1.7°C (SE = 0.3; range from 1.5 - 2.8°C). Temperature changes were
5814 more severe in the CC_NB watersheds with no leave trees (4.2 and 4.4°C), however, this
5815 difference was not analyzed. The percentage of time the post-harvest, no-buffer treatments spent
5816 above the 16 and 15 °C thresholds were 1.3% and 4.7%, respectively. This was an increase from
5817 pre-harvest values that showed no instances of temperatures above 16°C, and only 0.2% of the
5818 recorded time above 15°C. The authors conclude that their evaluation of temperature responses
5819 as potential biologically significant changes adds context to the changes and fluctuations
5820 observed in each harvest design. While significant changes in mean and percentile changes in
5821 temperature were observed, the amount of time spent above critical temperature thresholds for
5822 important amphibian species was minimal.

5823

5824 **SHD, Stream temperature**

5825

5826 Chan et al., 2004 (Removed from focal list, significant results only apply to fauna)

5827

5828 Chan, S.S., Anderson, P.D., Cissel, J., Larsen, L., Thompson, C., 2004. Variable density
5829 management in Riparian Reserves: lessons learned from an operational study in managed forests
5830 of western Oregon, USA. USDA Forest Service. <https://doi.org/10.1016/j.foreco.2013.06.055>

5831

5832 The purpose of this study was to assess the ability of variable retention thinning and riparian
5833 buffers at accelerating late-seral habitat, facilitating rare species management, and maintaining
5834 ecological functions within riparian zones of 40–70-year-old headwater forests in western
5835 Oregon. This study evaluated 13 separate sites each averaging ~ 100 ha whereby 4 buffer width
5836 treatments adjacent to variable retention thinning prescriptions were assessed. Buffer treatments
5837 include: (1) one site potential tree; (2) two-site potential trees; (3) variable buffer width based on
5838 vegetation and/or topographic site factors; (4) streamside buffer of only the first tree whereby

5839 thinning treatments applied up to 6 m of stream. Thinning treatments included: (1) Unthinned
5840 control - 500-750 trees per hectare; (2) High density retention - 70-75% of area thinned to 300
5841 TPH, 25-30% unthinned riparian reserves or leave islands; (3) Moderate density retention - 60-
5842 65% area thinned to 200 TPH, 25-30% unthinned riparian reserves or leave islands with 10%
5843 circular patch openings; (4) Variable density retention - 10% area thinned to 100 TPH, 25-30%
5844 thinned to 200 TPH, 25-30% thinned to 300 TPH, 20-30% unthinned riparian reserves or leave
5845 islands with 10% circular patch openings. Variables measured include stand development
5846 metrics, understory vegetation, microclimate, aquatic ecology, invertebrates, lichens, and
5847 bryophytes. Early findings from this study show that relatively small changes in the riparian
5848 environment are attributed to different residual thinning densities and different buffer widths.
5849 According to the results, the most suitable habitat for many species of fauna is consistently found
5850 within 5 m of the stream. The largest changes in relative humidity in warm and dry summer
5851 conditions occur within 15 m of the stream channel and begin to stabilize at 25 m. In summary,
5852 the early findings of this study indicate the near-stream riparian environment provides critical
5853 functions and habitat for a wide variety of organisms.

5854

5855 **Sediment**

5856

5857 Reiter et al., 2009

5858

5859 Reiter, M., Heffner, J. T., Beech, S., Turner, T., & Bilby, R. E. (2009). Temporal and Spatial
5860 Turbidity Patterns Over 30 Years in a Managed Forest of Western Washington 1. *JAWRA Journal*
5861 *of the American Water Resources Association*, 45(3), 793-808. [https://doi.org/10.1111/j.1752-](https://doi.org/10.1111/j.1752-1688.2009.00323.x)
5862 1688.2009.00323.x

5863

5864 This study evaluates the efficacy of the changes in a forest practices plan developed in 1974 to
5865 reduce sediment inputs into streams in the Deschutes River watershed of western Washington. To
5866 test this, the researchers analyzed 30 years of data (1975-2005) on water levels, discharge,
5867 suspended sediment, turbidity, and water and air temperature from four permanent sampling sites
5868 representing a range of basin sizes from small tributary headwaters to the mainstem of the
5869 Deschutes River. In the 1970s roughly 30% of the watershed had been harvested and
5870 approximately 63% of the existing road network had been constructed. Timber harvest continued
5871 until the early 1990s and the road network was completed in the late 1970s but updated to
5872 include culverts and sediment traps in the early 2000s. The researchers used turbidity as a proxy
5873 for suspended sediment correlation and corrected for typical seasonal increases in streamflow.
5874 The results showed a declining trend in turbidity at all permanent sampling sites during the study
5875 period even with active forest management. Following the road construction and harvest
5876 activities of the 1980s turbidity levels continued to decline until the year 2000 when they

5877 returned to pre-logging levels. The authors interpret these results as evidence that management's
5878 increased attention to reducing sediment is responsible for the reduction in sediment transport.

5879

5880 **Effect of debris torrents on shade, vegetation, and stream temperature**

5881

5882 D'Souza et al., 2011

5883

5884 D'Souza, L.E., Reiter, M., Six, L.J., Bilby, R.E., 2011. Response of vegetation, shade and stream
5885 temperature to debris torrents in two western Oregon watersheds. *Forest Ecology and*
5886 *Management* 261, 2157–2167. <https://doi.org/10.1016/j.foreco.2011.03.015>

5887

5888 The purpose of this study was to examine the effects of debris torrents on vegetation, shade, and
5889 stream temperature eight years after an extreme storm-related disturbance. This study examined
5890 two separate managed watersheds which were affected by storm-related debris torrents in 1996.
5891 This study addressed several questions regarding the patterns and rate of vegetation, shade and
5892 water temperature change post-disturbance: (1) What is the relationship between vegetation and
5893 local landform and substrate types along the study streams? (2) Does vegetation composition and
5894 structure, stream shade and water temperature in debris torrented streams differ between the two
5895 watersheds? and (3) How does recovery of stream temperature relate to vegetation and shade
5896 recovery and does this differ through time between watersheds? Data was gathered from
5897 multiple headwater streams following the disturbance in 1996 at 2 managed watersheds: the
5898 Williams River watershed (WRW), and the Calapooia River watershed (CRW). Data for stream
5899 temperature, to analyze stream temperature recovery, was collected immediately following the
5900 disturbance event in 5 streams, 3 at the CRW (2 disturbed; 1 reference), and 3 at the WRW (1
5901 disturbed, 1 reference) and for 8 years through the summer of 2004. Eight years post-disturbance
5902 12 disturbed streams (n = 6 for each watershed) were selected for data collection to examine the
5903 relationships between riparian vegetation, shade, and stream temperatures. Data on landform,
5904 substrate, and vegetation (density, species, and seedlings) were collected at each stream. Stream
5905 shade was estimated using hemispherical photographs taken 1 m above the stream center during
5906 summer and winter months and compared using t-tests. Stream temperature data was collected
5907 using continuously recording thermistors. Data were averaged and analyzed using t-tests, chi-
5908 square tests, simple linear regression, Pearson's correlation coefficient, and analysis of
5909 covariance. Results from this study show early successional species red alder and willow species
5910 dominated areas affected by debris torrents. All red alder variables (density, basal area, and
5911 height) showed a significant relationship with vegetation-related shade. Red alder showed a
5912 significantly higher density ($p = 0.0277$) and basal area ($p = 0.0367$) in the WRW sites. While
5913 stem density of red alder was similar in both watersheds, the size of the trees differed suggesting
5914 that colonization and/or growth of red alder in the WRW occurred more rapidly than in the CRW.
5915 However, there was no statistical difference in landforms or site factors between watersheds that

5916 explained these differences. The only correlations found were a negative relationship between
5917 alder density and rock; and a positive relationship between alder basal area and moss suggesting
5918 a relationship between moisture availability and red alder establishment and growth. The authors
5919 note that the WRW sites experienced greater precipitation in the years following disturbance and
5920 may have contributed to the greater growth rates of red alder, but no analysis was conducted.
5921 Total shade was also significantly higher in the WRW ($p = 0.0049$). Mean maximum daily
5922 temperature fluctuations ($p = 0.0483$), and 7-day maximum temperatures ($p=0,0483$) were also
5923 significantly lower in the WRW streams. Mean max daily stream temperatures were lower in the
5924 WRW streams but the difference was not significant ($p = 0.0779$). The authors conclude that
5925 even though the debris torrents resulted in poor soil conditions, the ability of red alder to thrive
5926 in these conditions resulted in rapid recovery of shade and thermal control.

5927

5928 **Stream temperature, shade and climate**

5929

5930 Reiter et al., 2015

5931

5932 Reiter, M., Bilby, R. E., Beech, S., & Heffner, J. (2015). Stream temperature patterns over 35
5933 years in a managed forest of western Washington. *JAWRA Journal of the American Water*
5934 *Resources Association*, 51(5), 1418-1435. <https://doi.org/10.1111/1752-1688.12324>

5935

5936 This study was an analysis of long-term stream temperature data in a western Washington
5937 watershed to evaluate the effects of forest management, before and after implementation of
5938 riparian forest best management practices, and climate change on stream temperatures. Stream
5939 temperature data from four permanent sampling stations in the Deschutes River Watershed.
5940 Stream and air temperature data was analyzed on a monthly basis from 1975-2009. This long-
5941 term dataset allowed for the examination of changes in stream temperature in four basins of
5942 varying size across a period from before stream buffers were implemented, during their
5943 implementation, and several instances of buffer expansion. Because the study period covered
5944 such a long time the changes in stream temperature based on climate change needed to be
5945 accounted for as well. The recovery of shade was estimated using the shade recovery function
5946 developed by R. Summers of Oregon State University (1983), whereby stream shade is estimated
5947 by angular canopy density (ACD) as a function of the age of stream-adjacent harvest units. To
5948 detect correlations of stream and air temperature change with land management activity
5949 separately from climate changes the data was fit to a model that included the effects of climate.
5950 The researchers accomplished this with a technique for deriving the residuals between stream
5951 temperature and climate called locally weighted scatterplot smoothing (LOWESS). The four
5952 watersheds varied in size from small (2 sites: Hard Creek, 2.4 km²; Ware Creek, 2.9 km²),
5953 medium (1 site: Thurston Creek, 9.3 km²), and large (1 site: The Deschutes River Station, 150
5954 km²). In the 1970s nominal buffer widths were required along fish-bearing streams, which

5955 expanded in the 1980s (requirements not listed), again in the mid-1990s to 23 m, and again to 30
5956 m in 2001. Methods for stream temperature data collection varied at different periods resulting in
5957 a margin of error for monthly temperatures of 0.14°C for 1975 - 1983, 0.09°C for 1984 – 1999,
5958 and 0.02°C. for 2000 – 2009. Because these margins of error were smaller than what the authors
5959 expected from climate and management, they were not accounted for in confidence intervals and
5960 p-values. The results for air temperature changes showed a statistically significant ($p \leq 0.05$)
5961 increasing trend in regional air temperatures for July TMAX_AIR and June and July
5962 TMIN_AIR. The trend for TMAX_AIR for July resulted in a trend magnitude of +0.07°C per
5963 year, for a total increase of 2.45°C over the 35-year record. For minimum air temperatures the
5964 magnitude of the June trend was +0.03°C per year while July TMIN_AIR had a trend magnitude
5965 of +0.04°C per year. The resulting increases in minimum temperatures for the period of record
5966 are 1.05°C and 1.40°C for June and July TMIN_AIR, respectively. Results for trends in stream
5967 temperature over the 35-year study period without adjustment for climate change showed no
5968 statistically significant trend in water temperature changes for the large watershed, while the
5969 medium watershed (Thurston Creek) showed decreasing trends in TMAX_WAT for June, July,
5970 and August, ranging in magnitude from 0.05°C (August) to 0.08°C (July) per year. For the
5971 smaller watershed, Hard Creek (Ware Creek was not included in this analysis), had significant
5972 decreasing trends in TMAX_WAT for July, August, and September. The magnitude of these
5973 trends was yearly decreases of TMAX_WAT by 0.05, 0.08, and 0.05°C, for July, August, and
5974 September, respectively. Significant changes in trends for TMIN_WAT were only found for the
5975 large basin site with yearly increases of 0.04, 0.03, and 0.04°C for July, August, and September,
5976 respectively. Results for stream temperature trends after adjusting for changes in air temperature
5977 (climate) showed significant decreasing trends in TMAX_WAT for the large basin by 0.04, 0.03,
5978 and 0.04°C yearly, for July, August, and September, respectively. For the medium basin, trends
5979 showed yearly decreases in TMAX_WAT of 0.07, 0.08, 0.06, and 0.03 for June, July, August,
5980 and September, respectively. For the small basin, climate adjusted trends in TMAX_WAT
5981 showed significant decreases in yearly trends by 0.05, 0.08, and 0.05 for July, August, and
5982 September, respectively. When stream temperature was examined with its correlation with
5983 estimated annual shade recovery from initial harvest (indexed by ACD). Significant correlations
5984 were found for monthly temperature metrics that were adjusted for climate, for all basins. The
5985 strongest correlations were for the smallest basin (Ware Creek) with correlation coefficients for
5986 climate adjusted maximum water temperatures (CTMAX_WAT) with ACD valuing -0.66, -0.78,
5987 -0.65, and -0.69 for June, July, August, and September, respectively. Correlation coefficients for
5988 Ware Creek CTMIN_WAT with ACD were -0.46, -0.64, -0.71, and -0.52 for June July, August,
5989 and September respectively. The largest basin (The Deschutes River) only showed significant
5990 correlations of CTMAX_WAT with ACD with July (-0.39) and August (-0.25); and only showed
5991 significant correlations of CTMIN_WAT with ACD for the months of August (+0.27), and
5992 September (+0.37). The authors interpret their results as evidence that following canopy
5993 recovery after implementation of riparian harvest rules the larger mainstem of the Deschutes
5994 River decreased in average maximum temperatures by approximately 1.3 °C when accounting for
5995 climate driven changes. The effects of canopy closure cooling were even more dramatic in the
5996 smaller headwater streams by 2.67 and 1.6 °C during the study period when accounting for
5997 climate driven changes (this includes a 0.5 °C correction based on climate warming). However,

5998 following re-initiation of timber harvest in 2001 for the area, when riparian protection buffers of
5999 30 m minimum were required, there was no detectable change in stream temperatures. The
6000 authors conclude that the results of this study show evidence that implementation of protection
6001 buffers in this area were sufficient in maintaining stream temperatures. Conversely, this study
6002 also shows evidence that despite these protections from land management induced stream
6003 temperature changes, these protections have been somewhat offset by the warming climate
6004 conditions.

6005

6006 **Overstory structure effects on understory light and vegetation**

6007

6008 Giesbrecht et al., 2017 (removed from focal, not relevant to questions, essentially a case study)

6009

6010 Giesbrecht, I.J.W., Saunders, S.C., MacKinnon, A., Lertzman, K.P., 2017. Overstory structure
6011 drives fine-scale coupling of understory light and vegetation in two temperate rainforest
6012 floodplains. *Can. J. For. Res.* 47, 1244–1256. [dx.doi.org/10.1139/cjfr-2016-0466](https://doi.org/10.1139/cjfr-2016-0466)

6013

6014 The purpose of this paper was to characterize the overstory structure and understory light
6015 regimes of temperate rainforest floodplains, and to assess the role of light and other site variables
6016 in driving stand vegetation patterns and processes. This study took place along two 1-ha coastal
6017 BC, Canada floodplain sites. These sites were selected as representative examples of floodplain
6018 forests in the Coastal Temperate Rainforest (CTR) as part of a larger network of long-term, old-
6019 growth monitoring plots. These sites were in the submontane variant of the very wet maritime
6020 subzone of the Coastal Western Hemlock zone (CWHvm1) of the B.C. coast. In each stand, the
6021 largest overstory trees are *Picea sitchensis* (Bong.) Carr., with several individuals taller than 60 m
6022 in height (maximum of 62 to 93 m). Based on coring a sample of main canopy trees, stand age at
6023 Kitlope is at least 95 years. Stand age at Carmanah is at least 350 years, based on a core from a
6024 50 m tall *P. sitchensis*. All trees ≥ 5 cm were measured along with all understory vegetation
6025 within 25 2m x 2m subplots. Stand characteristics were recorded as well as information on gap
6026 origins. Hemispheric canopy photographs were taken to estimate understory light penetration.
6027 Visual estimations of organic material, mineral layer, CWD, and other substrates were taken in
6028 each vegetation subplot. Relationships among measures of light transmission, vegetation
6029 structure, and diversity were analyzed with linear correlation analysis. Nonmetric
6030 multidimensional scaling was used to describe variation in species composition on multivariate
6031 axes. Results from this study show both sites as having a relatively high degree of canopy
6032 openness (11-11.6%) and light transmission (median 18% full sun) compared to many other
6033 tropical and temperate forests. Light transmission at both sites is however significantly lower
6034 than a number of old-growth sites in Quebec and northern BC. The origins of canopy openness
6035 and stand shade differ between both sites indicating distinct stand processes and different stages
6036 of stand development. Further, light levels vary substantially within short distances at each site

6037 reflecting a complex overstory structure. Although results from this study are reflective
6038 specifically of the coastal temperate rainforests of BC, the descriptive assessment of these two
6039 separate floodplain forests reveal a natural disturbance history which fostered a high degree of
6040 canopy openness and structural heterogeneity which may ultimately aid in informing future
6041 temperate rainforest floodplain restoration efforts.

6042

6043 **LW**

6044

6045 Reid & Hassan, 2020

6046

6047 Reid, D. A., & Hassan, M. A. (2020). Response of in-stream wood to riparian timber harvesting:
6048 Field observations and long-term projections. *Water Resources Research*, 56(8),
6049 e2020WR027077. <https://doi.org/10.1029/2020WR027077>

6050

6051 This paper proposes a conceptual model of wood storage response to different harvesting
6052 intensities. The model predicts how LW in streams is expected to change spatially and
6053 temporally following three different harvest patterns. The model was developed with 45 years of
6054 LW data retrieved from the Pacific coastal region of Vancouver Island, British Columbia. The
6055 Carnation Creek watershed, which supports gravel bed forested streams, contains riparian forests
6056 that have received a wide range of harvest plans implemented. During logging in the 1970s and
6057 '80s riparian forests of one region were harvested with buffer widths ranging from 1 – 70 meters
6058 in upstream reaches, and another region with near complete or complete removal of vegetation to
6059 the streams edge in downstream reaches. In-stream wood volume and characteristics data has
6060 been collected in eight of these study reaches since 1973 (pre-harvest). The researchers used this
6061 data with simulation modelling to develop a reach-scale wood budget model that predicts wood
6062 loss and recover patterns for 300 years (1900-2200). This paper has two objectives: (i) to use this
6063 field data and modeling approach to examine LW storage changes, the time to minimum wood
6064 load, and wood load recovery times as a result of riparian timber harvesting and forest
6065 regeneration, and (ii) to describe the characteristics of in stream wood, with particular focus to
6066 spatial and temporal patterns in wood storage over the multidecade scale following harvesting in
6067 riparian areas. The model was based upon the proposed response outlined by Murphy and Koski
6068 (1989). Wood budget responses were estimated using three management scenarios. Scenario 1 is
6069 a no harvest scenario, in this configuration, the loss of wood supply from the landscape has little
6070 to no impact on input from wood mortality or bank erosion, and therefore in-stream storage,
6071 decay, and transport of wood is not affected. Scenario 2 represents partial loss of forested area in
6072 the riparian zone, which will lead to a near-immediate reduction in wood recruitment to the
6073 channel from mortality and bank erosion along harvested areas. Wood decay and other
6074 components of wood loss will exceed rates of input, leading to a reduction in storage until time
6075 T_{min} , the point where wood recruitment equals losses as the forest regrows in riparian areas and

6076 the greatest overall reduction in storage has occurred (ΔS_{max}). Wood storage increases
6077 thereafter, eventually recovering to preharvest levels after time T_{rec} . Scenario 3 represents an
6078 intensive harvest scenario where most of the riparian area has undergone harvesting over a short
6079 period of time, a major reduction of input from bank erosion and mortality occurs. This greater
6080 reduction leads to a much larger ΔS_{max} than in Figure 1b as wood losses exceed recruitment.
6081 However, as the dominant wood sources recover at the same rate, the time to T_{min} and T_{rec} is
6082 similar under both the moderate and intensive harvest scenarios. Results of the model show
6083 evidence that wood storage in streams of harvested reaches, hits its minimum value in 50 years
6084 or more following loss of LW input, decay, and export of current stock. Recovery of LW volume
6085 in-streams following harvest is estimated to take approximately 150-200 years. The pattern and
6086 intensity of the harvesting operation had little effect on LW loss and recovery times but did affect
6087 the estimated magnitude of LW volume loss in the first 50 – 80 years. These results show
6088 evidence that timber harvest has a long-term effect on LW storage and loading dynamics even
6089 with protective buffers. However, buffers can ameliorate the magnitude of LW loss during the
6090 recovery period. The one caveat of this model is it doesn't account for as much variability on
6091 stream configuration or valley morphologies that are likely to affect LW storage.

6092

6093 **Buffers and LW Recruitment**

6094

6095 Grizzel et al., 2000 (Removed)

6096

6097 Grizzel, J., McGowan, M., Smith, D., Beechie, T., 2000. STREAMSIDE BUFFERS AND
6098 LARGE WOODY DEBRIS RECRUITMENT: EVALUATING THE EFFECTIVENESS OF
6099 WATERSHED ANALYSIS PRESCRIPTIONS IN THE NORTH CASCADES REGION
6100 (Timber/Fish/Wildlife Monitoring Advisory Group and the Northwest Indian Fisheries
6101 Commission). fp_tfw_mag1_00_003

6102

6103 This study analyzed the effectiveness of the Washington Watershed Analysis (WWA)
6104 prescriptions at recruiting large woody debris. This study took place at 10 riparian sites
6105 distributed across 5 watershed administrative units in the Northern Cascades of Washington. Ten
6106 sites were randomly chosen with gradients and buffer width classes in compliance with WWA
6107 indices. To analyze WWA effectiveness, debris frequency and size at each site were compared to
6108 targets derived from WWA. In addition, debris recruitment was compared between three buffer
6109 width classes. Geometric mean diameter and geometric mean length of debris was calculated
6110 based on measurements of midpoint diameter and total lengths. This data was then compared to
6111 targets derived from a channel width-dependent regression. Results show post-harvest mortality
6112 substantially decreasing stand density at several sites. In stream frequency targets were met at
6113 most sites; however, debris categorized as "good" for habitat was only achieved at four out of ten
6114 sites. At the time of data collection, a large portion of debris recruited from buffers was either

6115 above or outside the bankfull flow zone. The authors point out that the degree to which the debris
6116 will influence fluvial processes in the future will depend on whether or not they are recruited into
6117 the stream and will also depend on the size and state of decay. The size of debris recruited from
6118 buffers was significantly smaller than recruited from unmanaged old-growth stands.
6119 Interestingly, data shows recruitment occurring from the outermost margins of the widest buffers
6120 (20-30 m, >30 m), suggesting narrow buffers may limit recruitment. The authors point out that
6121 the large degree of variability in recruitment from site to site suggests windthrow as an important
6122 causal factor. In channels oriented perpendicular to damaging winds (east-west), there was a
6123 higher likelihood of potential recruitment as compared to channels oriented parallel to damaging
6124 winds. The authors conclude with multiple recommendations for future study. First, they suggest
6125 integrating habitat inventory with recruitment to achieve a better understanding of relationships.
6126 Second, they suggest future study into the fate of debris suspended above channels given much
6127 of our current understanding is based on assumptions of decay and breakage. Finally, they
6128 recommend study into factors influencing windthrow in riparian buffers.

6129

6130 **Sediment**

6131

6132 Rachels et al., 2020

6133

6134 Rachels, A. A., Bladon, K. D., Bywater-Reyes, S., & Hatten, J. A. (2020). Quantifying effects of
6135 forest harvesting on sources of suspended sediment to an Oregon Coast Range headwater stream.
6136 *Forest Ecology and Management*, 466, 118123. <https://doi.org/10.1016/j.foreco.2020.118123>

6137

6138 This study uses sediment source fingerprinting techniques to quantify the proportional
6139 relationship of sediment sources (hillslope, roads, streambanks) in harvested and un-harvested
6140 watersheds of the Oregon Coast Range. The researchers used sediment traps, and chemical
6141 analysis to estimate the origin of suspended sediment in the stream and to quantify magnitude of
6142 sediment stored in protection buffers. The study included one catchment (Enos Creek) that was
6143 partially clearcut harvested in the summer of 2016 and an unharvested reference catchment
6144 (Scheele Creek) located ~3.5 km northwest of Enos Creek. The paired watersheds had similar
6145 road networks, drainage areas, lithologies and topographies. The treatment watershed was
6146 harvested with a skyline buffer technique in the summer of 2016 under the Oregon Forest
6147 practices Act policy that requires a minimum 15 m no-cut buffer. The proportion of suspended
6148 sediment sources were similar in the harvested ($90.3 \pm 3.4\%$ from stream bank; $7.1 \pm 3.1\%$ from
6149 hillslope) and unharvest ($93.1 \pm 1.8\%$ from streambank; $6.9 \pm 1.8\%$ from hillslope) watersheds.
6150 However, the harvested watershed contained a small portion of sediment from roads ($3.6 \pm$
6151 3.6%), while the unharvested reference watershed suspended sediment contained no sediment
6152 sourced from roads. In the harvested watersheds the sediment mass eroded from the general
6153 harvest areas (96.5 ± 57.0 g) was approximately 10 times greater than the amount trapped in the

6154 riparian buffer (9.1 ± 1.9 g), and 4.6 times greater than the amount of sediment collected from
6155 the unharvested hillslope (21.0 ± 3.3 g). These results suggest that the riparian buffer was
6156 efficient in reducing sediment erosion relative to the harvested area. The caveat of this study was
6157 the limited sample size (1 treatment, 1 paired reference watershed) and does not incorporate the
6158 effects of different watershed physiography on sediment erosion.

6159

6160 **SED**

6161

6162 Puntenney-Desmond et al., 2020

6163

6164 Puntenney-Desmond, K. C., Bladon, K. D., & Silins, U. (2020). Runoff and sediment production
6165 from harvested hillslopes and the riparian area during high intensity rainfall events. *Journal of*
6166 *Hydrology*, 582, 124452. <https://doi.org/10.1016/j.jhydrol.2019.124452>

6167

6168 This study uses simulation modeling to evaluate the differences in run-off rates, sediment
6169 concentrations, and sediment yields between watershed harvested areas, along the interface of
6170 harvested areas and riparian buffers, and within riparian buffers during periods of high-intensity
6171 rainfall events. The model simulations were calibrated with soil and watershed characteristic data
6172 collected from the Star Creek catchment located in southeastern Alberta. Fifteen plots were
6173 selected for rainfall simulations along three transects on a north facing hillslope (aspect: $\sim 358^\circ$)
6174 and along two transects on a southeast facing hillslope (aspect: $\sim 129^\circ$). Each transect consisted
6175 of three plots that were spaced ~ 20 m apart along the planar hillslopes. Each plot was one
6176 square-meter, which was bounded by a three-sided steel frame that was inserted into the soil with
6177 the open side facing down the slope. The plots were located either (a) within the general harvest
6178 area, (b) along the edge of the riparian buffer at the interface with the harvested area, or (c)
6179 within the riparian buffer. The high-intensity rainfall events were calibrated to mimic 100-year,
6180 or greater, storm events of the Northern Rocky Mountains (1-hour high intensity rainfall). The
6181 results showed runoff rates and surface and shallow subsurface were greatest in the buffer areas
6182 than in the harvested areas or in the harvest-buffer interfaces especially during dry conditions.
6183 During the dry condition rainfall simulations, the general pattern of runoff rates (surface/shallow
6184 subsurface flow) was riparian buffer (175.6 ± 17.3 [SE] ml min^{-1}) > harvest-riparian edge
6185 (125.8 ± 18.2 ml min^{-1}) > general harvest area (37.2 ± 8.5 ml min^{-1}). Mean runoff rates within
6186 the riparian buffer plots were greater than within the general harvest area plots ($t = 2.90$, $p = .03$).
6187 Runoff ratios were only statistically greater in the riparian buffer plots ($13.9 \pm 3.1\%$) relative to
6188 the general harvest area ($2.9 \pm 1.5\%$) during the dry conditions. All runoff ratios declined during
6189 the wet condition rainfall simulations relative to the dry condition simulations with no evidence
6190 for differences between any of the plot positions ($p > .27$ for all pairwise comparisons). During
6191 the dry condition rainfall simulations, the general patterns of sediment concentrations and
6192 sediment yields were opposite of the runoff rates, with the general harvest area > harvest-riparian

6193 edge > riparian buffer. The sediment concentration was (a) 424.8 mg l⁻¹ (151.0–1195.3 mg l⁻¹)
6194 in the general harvest area, (b) 100.9 mg l⁻¹ (45.8–222.1 mg l⁻¹) along the harvest riparian
6195 edge, and (c) 26.9 mg l⁻¹ (12.2–59.1 mg l⁻¹) in the riparian buffer. Statistically, there was
6196 strong evidence for differences in sediment concentrations between the general harvest area and
6197 along the harvest-riparian edge ($t = 3.21$, $p = .01$) and between the harvest area and the riparian
6198 buffer ($t = 6.17$, $p < .001$). Statistically, there was no evidence for differences in sediment yields
6199 between any of the plot positions. Sediment concentration among plot positions remained the
6200 same during the wet rainfall simulations as the dry rainfall simulations—general harvest area >
6201 harvest-riparian edge > riparian buffer. The geometric mean and 95% confidence intervals (back-
6202 transformed) for the sediment concentration was (a) 285.7 mg l⁻¹ (67.9–1201.5 mg l⁻¹) in the
6203 general harvest area, (b) 79.6 mg l⁻¹ (36.5–173.5 mg l⁻¹) along the harvest-riparian edge, and
6204 (c) 22.3 mg l⁻¹ (3.5–141.7 mg l⁻¹) in the riparian buffer. However, while sediment
6205 concentrations differed most strongly between the general harvest area and the riparian buffer (t
6206 $= 3.51$, $p = .01$), other pairwise comparisons were not significant ($p > .20$). Statistically, there
6207 was no evidence for differences in sediment yields between any of the plot positions for rainfall
6208 simulations during wet conditions. The authors speculate this was likely due to the greater soil
6209 porosity in the disturbed, harvested areas. Sediment concentration in the runoff, however, was
6210 approximately 15.8 times higher for the harvested area than in the riparian buffer, and 4.2 times
6211 greater than in the harvest-buffer interface. Total sediment yields from the harvested area (runoff
6212 + sediment concentration) were approximately 2 times greater than in the buffer areas, and 1.2
6213 times greater in the harvest-buffer interface (however, these proportions were not statistically
6214 different). Replication of the model showed high levels of variability in total run off rate,
6215 sediment concentrations, and sediment yields but the relationships between timing and relative
6216 magnitudes between the three experimental areas were consistent. The authors speculate that
6217 these results will become more relevant as climate change is expected to increase the frequency
6218 of high-intensity rainfall events following dry periods in this area. They suggest expanding
6219 similar methods to understand these effects in areas of different hydro-climatic settings.

6220

6221 **Stream Temperature**

6222

6223 Pollock et al., 2009

6224

6225 Pollock, M. M., Beechie, T. J., Liermann, M., & Bigley, R. E. (2009). Stream temperature
6226 relationships to forest harvest in western Washington I. JAWRA Journal of the American Water
6227 Resources Association, 45(1), 141-156. <https://doi.org/10.1111/j.1752-1688.2008.00266.x>

6228

6229 This study investigates the effect of watershed harvest percentage, and time since harvest on
6230 summer stream temperatures at different scales in the Olympic Peninsula, Washington. The
6231 researchers examined recorded stream temperature data in 40 small watersheds that experienced

6232 a range of harvest from 0 – 100% (7 unharvested, 33 harvested between 25-100%), with
6233 regrowth age groups binned for analysis as recently clear cut (< 20 years old) and less recently
6234 clearcut (mostly < 40 years old). Unharvested sites were estimated as being >150-years old.
6235 Clearcut is defined in this paper as removing any protective canopy cover for streams. This study
6236 tested 3 hypotheses: (1) the condition of the riparian forest immediately upstream of a site
6237 primarily controls stream temperature, (2) the condition of the entire riparian forest network
6238 affects stream temperature, and (3) the forest condition of the entire basin affects stream
6239 temperature. These hypotheses were test by examining correlations of stream temperature with
6240 the condition of the immediate upstream riparian forest, or more correlated with forest conditions
6241 more spatially distant and on a coarser scale, such as the entire upstream riparian forest network
6242 or the forest condition of the entire basin. To avoid site effects in their analysis sites were chosen
6243 from a narrow range of subbasin sizes (approximately 1-10 km²) and elevation (75-400 m).
6244 Further, all sites were underlain by sedimentary rock and had perennial flow. Each hypothesis
6245 was tested with linear regression to evaluate the correlations of each age group at each scale with
6246 stream temperature data. The researchers also used AIC value comparisons for model selection to
6247 assess the correlation of other physiographic features (elevation, basin area, aspect, slope, or
6248 geologic composition) with stream temperatures. Results of general temperature patterns showed
6249 that average daily maximum (ADM) were strongly correlated with average diurnal fluctuations
6250 ($r^2 = 0.87$, $p < 0.001$, $n = 40$), indicating that cool streams also had more stable temperatures. For
6251 basin-level harvest effects on stream temperatures. The percentage of the basin harvested
6252 explained 39% of the variation in the ADM among subbasins ($r^2 = 0.39$, $p < 0.001$, $n = 40$) and
6253 32% of variation in the average daily range (ADR) ($r^2 = 0.32$, $p < 0.001$, $n = 40$). The median
6254 ADM for the unharvested subbasins was 12.8 °C (mean = 12.1 °C), which was significantly
6255 lower than 14.5 °C, the median (and average) ADM for the harvested subbasins ($p < 0.001$).
6256 Likewise, the median (and average) ADR for the unharvested subbasins was 0.9 °C, which was
6257 significantly lower than 1.6 °C, the median ADR (average = 1.7 °C) for the harvested subbasins
6258 ($p < 0.001$). Results for the correlations between the riparian network scale forest harvest and
6259 stream temperature showed that the total percentage of the riparian forest network upstream of
6260 temperature loggers harvested explained 33% of the variation in the ADM among subbasins ($r^2 =$
6261 0.33 , $p < 0.001$, $n = 40$) and 20% of variation in the ADR ($r^2 = 0.20$, $p = 0.003$, $n = 40$).
6262 However, the total percentage of upstream riparian forest harvested within the last 20 years was
6263 not significantly correlated to ADM or ADR. Results for near upstream riparian harvest and
6264 stream temperature showed either non-significant, or very weakly significant correlations. For
6265 example, there were no significant correlations between the percentage of near upstream riparian
6266 forest recently clear-cut and ADM temperature ($r^2 = 0.03$, $p = 0.79$, $n = 40$), the ADR of stream
6267 temperatures ($r^2 = 0.02$, $p = 0.61$, $n = 40$) or any other stream temperature parameters. The
6268 proportion of total harvested near upstream riparian forest (avg = 0.66, SD \pm 0.34, range = 0.0-
6269 1.0) was weakly correlated with ADM ($r^2 = 0.12$, $p = 0.02$, $n = 40$) and not significantly
6270 correlated with ADR ($r^2 = 0.07$, $p = 0.06$, $n = 40$). Even when the upstream riparian corridor
6271 length was shortened to 400 m and then to 200 m, and the definition of recently harvested was
6272 narrowed to <10 year, no significant relationships between temperature and the condition of the
6273 near upstream riparian forest was found. Results for the effect of physical landscape variables on
6274 stream temperature found that the variables of elevation, slope, aspect, percent of the basin with

6275 a glacial surficial geology, upstream distance of the site to sedimentary (bedrock) geology, and
6276 the percent of sedimentary surficial geology in the basin individually explain between 5% and
6277 14% more of the variability relative to basin harvest. Adding any one of these variables to the
6278 model increases the r^2 from 0.40 up to between 0.48 and 0.51. However, the coefficient for
6279 percent of basin harvested and its standard error stay essentially the same, thus the authors
6280 concluded that adding additional variables to the model did not change the basic finding that
6281 there is a strong relationship between ADM and total amount of harvest in a basin. Thus, for
6282 these models, the percentage of basin area harvested was the best predictor of variation in mean
6283 maximum stream temperatures. The probability of stream temperatures increasing beyond DOE
6284 standards (16 °C for seven-day average of maximum temperatures) increased with percent
6285 harvest. Nine of the 18 sites with 50-75% harvest and seven of the nine sites with >75% harvest
6286 failed to meet these standards. The authors interpret these results as evidence that the total
6287 amount of forest harvested within a basin, and within a riparian stream network are the most
6288 important predictors of changes in summer stream temperatures. They conclude that watersheds
6289 with 25-100% of their total area harvested had higher stream temperatures than watersheds with
6290 little or no harvest. Furthermore, they speculate that past basin-wide timber management can
6291 impact stream temperatures over long periods of time in a way that riparian buffer treatments
6292 cannot entirely ameliorate.

6293

6294 **Stream Temperature**

6295

6296 Groom et al., 2011a

6297

6298 Groom, J.D., Dent, L., Madsen, L.J., 2011. Stream temperature change detection for state and
6299 private forests in the Oregon Coast Range. *Water Resources Research* 47.
6300 <https://doi.org/10.1029/2009WR009061>

6301

6302 The purpose of this study was to evaluate the effectiveness of private and state forest buffer rules
6303 on state water quality stream temperature antidegradation standards in the Oregon Coast Range.
6304 According to the Department of Environmental Quality (DEQ), under the Protecting Cold Water
6305 (PCW) criterion, anthropogenic activities are not permitted to increase stream temperature by
6306 more than 0.3 °C above its ambient temperature. In addition, the cumulative amount of
6307 anthropogenic temperature increase allowed in streams with temperature total maximum daily
6308 loads (TMDLs) is 0.3 °C for all sources combined. Stream temperature and riparian stand
6309 conditions were measured pre- and post-harvest between 2002 and 2008 at 33 sites (18 private-
6310 owned, 15 state-managed). Treatment stands included 26 clear-cuts and 7 partial cuts (leave tree
6311 requirements not specified), all of which were harvested in adherence to FPA (private) and FMP
6312 (state) standards. Private sites followed FPA rules whereby the riparian management area
6313 (RMA)s are 15 and 21 m wide on small and medium fish-bearing streams, respectively, with a 6

6314 m no-cut zone immediately adjacent to the stream. State sites followed the state management
6315 plan whereby a 52 m wide buffer is required for all fish-bearing streams, with an 8 m no cut
6316 buffer immediately adjacent to the stream. Stream temperature data was collected for at least 2
6317 years prior to harvest. Reference reaches were located immediately upstream from the harvested
6318 reaches. Generalized least square regression was used to model ambient conditions while
6319 accounting for temporal autocorrelation. The authors examined prediction intervals to assess the
6320 rule exceedance (>0.3 °C increase in temperature). Results indicate that sites harvested according
6321 to FPA standards exhibited a 40.1% probability that a pre harvest to post harvest comparison of
6322 2 years of data will detect a temperature change of > 0.3 °C. Conversely, harvest to state FMP
6323 standards resulted in an 8.6% probability of exceedance that did not significantly differ from all
6324 other comparisons. The a priori and secondary post hoc multimodel comparisons did not indicate
6325 that timber harvest increased the probability of PCW exceedance at state sites. The authors point
6326 out that the 0.3 °C change threshold still lies 1 or 2 orders of magnitude lower than previous
6327 findings from studies which took place prior to the enactment of the riparian protection
6328 standards. The authors recommend further research looking into the potential persistence of
6329 stream temperature change downstream after harvest. In addition, they recommend looking into
6330 the biological significance of increases in stream temperature change particularly to aquatic life.

6331

6332 **Stream and subsurface water temperature**

6333

6334 Guenther et al., 2014

6335

6336 Guenther, S.M., Gomi, T., Moore, R.D., 2014. Stream and bed temperature variability in a
6337 coastal headwater catchment: influences of surface-subsurface interactions and partial-retention
6338 forest harvesting. *Hydrological Processes* 28, 1238–1249. <https://doi.org/10.1002/hyp.9673>

6339

6340 This study documented changes in stream and subsurface water temperature in response to forest
6341 harvesting in two paired headwater catchments. Specifically, the researchers hypothesized that
6342 post-logging changes in bed temperatures should be greatest in locations experiencing hyporheic
6343 downwelling (DW) and least in areas with lateral inflow/groundwater discharge. This study took
6344 place in the University of British Columbia Malcolm Knapp Research Forest near Vancouver,
6345 Canada. As a part of an ongoing study into the effects of riparian buffers on stream ecology. The
6346 catchments of 3 southerly-aspect first order streams were harvested using partial retention (50%
6347 removal of basal area including riparian zone) methods resulting in approximately 14% reduction
6348 in canopy cover on average; 3 other southerly-aspect streams served as unharvested controls.
6349 Before thinning treatments, the harvested riparian forests were dominated by western hemlock,
6350 (*Tsuga heterophylla*), western red cedar (*Thuja plicata*), and Douglas-fir (*Pseudotsuga*
6351 *menziesii*). The forests were mature second growth forests with trees approximately 30–40 m tall,
6352 and canopy closure than 90%. Harvest operations began in September 2004 and completed in

6353 November of 2004. Temperature data was summarized from 10-minute intervals to daily
6354 minimum, maximum, and mean temperatures for stream and bed temperatures for one-year prior
6355 to, and one year following harvest. An analysis of the post-harvesting effects was conducted
6356 using a paired-catchment analysis. Results from this study show treatment sites resulted in higher
6357 daily maximum stream and bed temperatures after harvest but smaller changes in daily minima.
6358 Daily maximum post-harvest stream temperatures averaged over July and August ranged from
6359 1.6°C to 3°C at different locations. Post harvest changes in bed temperature at the lower reaches
6360 were smaller than changes in stream temperature, but was greater at sites with downwelling (DF)
6361 flow, and decreased with depth at upwelling (UW) and DF sites dropping to approximately 1°C
6362 at a depth of 30 cm. Changes did not vary significantly with depth at the middle reach, and
6363 averaged approximately 1°C change in daily maximum bed temperature over July and August. In
6364 summary, stream temperature responses differed at different locations within the cutblock. Bed
6365 temperatures also differed between UW and DW zones as well as between reaches with different
6366 contributions of lateral inflow. Given evidence that stream/bed temperature is shown to change
6367 spatially and with differences in hyporheic exchange and lateral inflow, the authors conclude by
6368 suggesting further research into the how these results might impact biological and ecological
6369 processes.

6370

6371 **Stream Temperature and evaporation/wind speed**

6372

6373 Guenther et al., 2012 (not in focal, does not separate the effects of shade reduction from wind
6374 speed/)

6375

6376 Guenther, S. M., Moore, R. D., & Gomi, T. (2012). Riparian microclimate and evaporation from
6377 a coastal headwater stream, and their response to partial-retention forest harvesting. *Agricultural
6378 and Forest Meteorology*, 164, 1-9.

6379

6380 The purpose of this study was to (1) develop and test an evaporimeter designed specifically to
6381 measure stream surface evaporation from headwater streams; (2) fit a wind function for
6382 computing evaporation from meteorological observations, and to compare it to previously
6383 published wind functions for evaporation from streams; and (3) quantify the influence of partial-
6384 retention forest harvesting on riparian microclimate and evaporation. This study was conducted
6385 in the University of British Columbia Malcom Knapp Research Forest (MKRF), approximately
6386 60 miles east of Vancouver, Canada and focused on the headwater stream of Griffith Creek. The
6387 harvesting treatment involved removal of 50% of the basal area from within the cut block,
6388 including the riparian zone. Smaller stems were removed, leaving the larger stems for harvest at
6389 a later date. creek. Analysis of paired pre- and post-logging hemispherical photographs indicated
6390 that canopy closure decreased by about 14% due to the logging treatment. Air temperature and
6391 relative humidity were measured by a Campbell Scientific CS500 sensor with stated accuracies

6392 of ± 0.5 °C for temperature and ± 3 – 6% for relative humidity. Wind speed was measured with a
6393 Met One anemometer with a stall speed of 0.447 m s⁻¹. Instruments were scanned every 10 s by
6394 a Campbell Scientific CR10x data logger; observations were averaged and stored every 10
6395 minutes. Evaporation was measured using four specially designed evaporimeters comprising an
6396 evaporation pan connected to a Mariotte cylinder. Results showed that Daily mean wind speeds
6397 increased following harvest, but were still consistently lower than wind speeds at the control site,
6398 with a maximum of 1.09 m s⁻¹. Vapor pressure was generally lower after harvesting. Vapor
6399 pressure deficit (vpd) increased following harvesting, but tended to remain lower than vpd
6400 measured at the control site. After harvesting, the relatively high wind speeds in the afternoon
6401 generally coincided with higher water temperatures, which in turn are associated with higher vpd
6402 at the water surface and a stronger vapor pressure gradient to drive evaporation. After harvest,
6403 wind speeds and vapor pressure gradients were higher and stability was weaker, consistent with
6404 the observed increase in evaporation. The authors conclude that the generally stronger relations
6405 between riparian and open microclimate variables after harvesting suggest that the riparian zone
6406 became more strongly coupled to ambient climatic conditions after harvesting as a result of
6407 increased ventilation. Further, that stream evaporation increased markedly as a result of partial
6408 retention harvest, consistent with the decrease in atmospheric vapor pressure, the increase in
6409 stream vapor pressure, the increase in wind speed and the decreased stability. In fact, prior to
6410 harvest, vapor pressure gradients often favored condensation rather than evaporation.

6411

6412 **LW**

6413

6414 Opperman, 2005 (Not in focal)

6415

6416 Opperman, J. J. (2005). Large woody debris and land management in California's hardwood-
6417 dominated watersheds. *Environmental Management*, 35(3), 266-277. DOI:10.1007/s00267-004-
6418 0068-z

6419

6420 The purpose of this paper was to evaluate the effects of stream and riparian area characteristics
6421 (bankfull width, gradient, basal area), and land ownership (public vs. private) on LW loading,
6422 and frequency, and debris jam frequency (response variables) in 21 hardwood-dominated forests
6423 of a Mediterranean climate region of northern California. The relationship between the stream
6424 and riparian area characteristics (explanatory variables: basal area of riparian trees, bankfull
6425 width, and gradient), and the response variables (woody debris loading and frequency, and
6426 debris-jam frequency) were evaluated with linear regression. The characteristics were then
6427 combined with ownership categories and their relative weight in explaining LW loading,
6428 frequency and pool frequency were assessed with a multi-variate analysis. Debris jam frequency
6429 was also analyzed by channel position with a chi-square. Results showed that debris jam
6430 frequency in the 21 reaches analyzed were strongly influenced by living standing trees rooted at

6431 the margins of the bank, especially in channel positions near the stream bank, but also spanning
6432 the channel partially, or completely. In general, LW loading was significantly higher in reaches
6433 adjacent to public lands (104 ± 13 m³/ha) than in those adjacent to private lands (46 ± 8 m³/ha;
6434 $P = 0.0015$). The strongest relationship for LW loading was with bankfull width ($r^2 = 0.32$; $p =$
6435 0.0006), and riparian basal area ($r^2 = 0.22$; $p = 0.006$) riparian basal area. This is likely the cause
6436 of the difference in public vs. private, as the public lands had significantly higher basal area in
6437 the riparian areas at distances >5 m from the stream, than the private lands. Debris jam frequency
6438 was also significantly influenced by riparian area gradient ($r^2 = 0.14$; $p = 0.03$) and basal area (r^2
6439 $= 0.11$; $p = 0.05$). The author concludes that landownership, and thus, land-management
6440 practices are driving factors in LW dynamics in this region.

6441

6442 **LW**

6443

6444 Nowakowski & Wohl, 2008

6445

6446 Nowakowski, A. L., & Wohl, E. (2008). Influences on wood load in mountain streams of the
6447 Bighorn National Forest, Wyoming, USA. *Environmental Management*, 42(4), 557-571.
6448 DOI:10.1007/s00267-008-9140-4

6449

6450 The purpose of this paper is to evaluate the relationship between riparian area characteristics, and
6451 land management practices with in-stream wood-loads in the Bighorn National Forest of
6452 northern Wyoming. The authors hypothesized that 1) valley geometry correlates with wood load,
6453 2) stream gradient correlates with wood load, 3) wood loads are significantly lower in managed
6454 watersheds than in similar unmanaged watersheds. The study analyzed data from 19 conifer
6455 dominated, forested headwater reaches in the bighorn mountains. Study reaches were separated
6456 by two watersheds, managed and unmanaged, with similar drainages, elevation, and lithology.
6457 Unmanaged watersheds were defined as having a history of minimal anthropogenic influences.
6458 The managed watershed had a history of different harvest prescriptions from unregulated in the
6459 late 1800s, clearcutting in the mid-1900s with tie floating practices. The relationship between in-
6460 stream wood loads (m³/ha) was analyzed with 11 valley-scale (elevation, forest type, forest stand
6461 density, etc.) and 13 channel-scale (reach gradient, channel width, etc.) variables with linear
6462 regression. Results support the first and third hypotheses. Across all streams, the highest
6463 explanatory power of all models tested produced land use (managed vs unmanaged), and basal
6464 area as a significant predictor of wood loads ($r^2 = 0.8048$). For the unmanaged watershed the
6465 model produced stream valley sideslope gradient as the single best predictor of wood load ($r^2 =$
6466 0.5748) supporting the first hypothesis. Shear stress was the best predictor of wood load in the
6467 managed watersheds ($r^2 = 0.2403$), These results did not directly support the second hypothesis.
6468 The authors suggest that while shear stress is a function of stream gradient (shear stress and
6469 stream gradient were significantly correlated, $r^2 = 0.9392$), gradient itself did not have the

6470 highest explanatory power of wood load in any of the models tested. Valley characteristics
6471 consistently explained more of the variability in wood load (42-80%) than channel characteristics
6472 (21-33%). When land use (managed vs. Unmanaged) effect on wood loads was analyzed the
6473 number of wood pieces per 100 m of stream was marginally significant ($p = 0.0565$), and the
6474 difference in wood volume per channel was significant ($p = 0.0200$) supporting the third
6475 hypothesis. When the significant valley and channel characteristics of the managed and
6476 unmanaged watersheds were controlled for, the significant difference in wood loads between
6477 managed and unmanaged watersheds were enhanced ($p = 0.0006$). Managed watersheds (1.1
6478 $m^3/100\ m$) had, on average, 2-3 times lower in-stream wood loads than unmanaged (3.3 $m^3/100$
6479 m) watersheds. These results suggest watersheds with a history of timber harvest have a decrease
6480 in stream wood loads than unmanaged watersheds, and that wood load dynamics can be driven
6481 by valley morphology, specifically, slope.

6482

6483 **Harvesting Practices on Suspended Sediment Yields**

6484

6485 Hatten et al., 2018

6486

6487 Hatten, J.A., Segura, C., Bladon, K.D., Hale, V.C., Ice, G.G., Stednick, J.D., 2018. Effects of
6488 contemporary forest harvesting on suspended sediment in the Oregon Coast Range: Alsea
6489 Watershed Study Revisited. *Forest Ecology and Management* 408, 238–248.
6490 <https://doi.org/10.1016/j.foreco.2017.10.049>

6491

6492 The objectives of this study were to (1) determine the effects of contemporary harvesting
6493 practices on suspended sediment yields and concentration, and (2) determine if contemporary
6494 harvesting practices produce lower sediment yields than historic practices. This study took place
6495 in the central Oregon Coast Range and consisted of a paired watershed study whereby Flynn
6496 Creek (FC) served as a reference watershed and Needle Branch (NB) served as a treatment
6497 watershed. A third watershed, Deer Creek (DC) served as a secondary control to compare
6498 historical vs contemporary harvest practices. The upper section of the treatment watershed was
6499 clearcut harvested using contemporary harvest practices (no buffer in non-fish-bearing streams
6500 with equipment exclusion zones, and a 15 m no-cut-buffer in fish-bearing streams) adhering to
6501 BMP's. Daily precipitation, discharge, and suspended sediment were collected at all three
6502 watersheds from October 2005 to June 2016. The upper half of the treatment watershed, (35 ha;
6503 measured at the Needle Branch Upper Gage or NBUG) was harvested in 2009 (Phase I) and the
6504 lower half (NBLG) was harvested in the fall of 2014 and mid-summer 2015 (Phase II). A model
6505 was developed using step wise linear regression to compare suspended sediment concentration
6506 (SSC). Differences in SSC among downstream sites and across harvest entries were compared
6507 utilizing an analysis of covariance. Results of the stepwise multiple linear regression showed
6508 strong evidence ($p < .001$) that all covariates (hydrograph limb, cumulative area discharge within

6509 water year, day of water year, daily precipitation, previous day's precipitation) were related to
6510 SSC across all watersheds. Both the mean and maximum SSC were greater in the reference
6511 catchments (FCG and DCG) compared to the harvested catchment (NBLG) across all water
6512 years. In NBLG the mean SSC was 32 mg L⁻¹ (~63%) lower after the Phase I harvest and 28.3
6513 mg L⁻¹ (~55%) lower after the Phase II harvest when compared to the pre-harvest
6514 concentrations. Compared to the reference watersheds, the mean SSC was 1.5-times greater in
6515 FCG (reference) compared to NBLG during the pre-harvest period. After the Phase I harvest the
6516 mean SSC in FCG (reference) was 3.1-times greater and after the Phase II harvest was 2.9-times
6517 greater when compared to the SSC in NBLG, the harvested watershed. Data from historical and
6518 contemporary harvests indicate contemporary practices are more effective at mitigating
6519 sedimentation. Historical data from the original study show harvesting without buffers, road
6520 building, and slash burning resulted in ~2.8 times increase in annual sediment yields and aquatic
6521 ecosystem degradation. The authors conclude that contemporary harvesting practices (i.e., stream
6522 buffers, smaller harvest units, no broadcast burning, leaving material in channels) using buffers
6523 were shown to sufficiently mitigate sediment delivery to streams, especially when compared to
6524 historic practices.

6525

6526 **Riparian Vegetation Removal Effects on Inputs and Production.**

6527

6528 Hetrick et al., 1998 (Removed, outside of timeline)

6529

6530 Hetrick, N.J., Brusven, M.A., Meehan, W.R., Bjornn, T.C., 1998. Changes in Solar Input, Water
6531 Temperature, Periphyton Accumulation, and Allochthonous Input and Storage after Canopy
6532 Removal along Two Small Salmon Streams in Southeast Alaska. *Transactions of the American*
6533 *Fisheries Society* 127, 859–875. [https://doi.org/10.1577/1548-](https://doi.org/10.1577/1548-8659(1998)127<0859:CISIWT>2.0.CO;2)
6534 [8659\(1998\)127<0859:CISIWT>2.0.CO;2](https://doi.org/10.1577/1548-8659(1998)127<0859:CISIWT>2.0.CO;2)

6535

6536 The purpose of this study was to assess whether or not the removal of second growth riparian
6537 vegetation would affect the production of juvenile coho salmon. In addition, this study aims to
6538 understand whether perceived effects are due to changes in habitat or food availability. This
6539 study took place in the Tongas National Forest on Prince of Wales Island, Alaska. Experimental
6540 reaches were divided into untreated and treated sections whereby treated sections had all
6541 vegetation on both sides of the streambank 6-15 m back removed. Stream discharge, water
6542 temperature, periphyton accumulation, allochthonous inputs, and storage of benthic organic
6543 matter were assessed during the summer and fall of 1988-1989. Differences in measured
6544 variables were assessed with a split-block analysis of variance. Results from this study show
6545 average light intensities reaching the water surface was significantly greater ($P < 0.01$) in the
6546 open canopy block than in the closed canopy block and was influenced significantly by weather
6547 conditions. Removal of riparian vegetation in both sections of the study significantly increased

6548 the accumulation of periphyton biomass and chlorophyll a ($P < 0.01$), and significantly decreased
6549 the amount of allochthonous organic inputs to streams ($P < 0.01$). Average daily allochthonous
6550 input rates for closed and open canopy conditions at Eleven creek were 789 and 6 mg AFDM/m²
6551 respectively, while input rates for closed and open canopy conditions at Woodsy creek were 805
6552 and 6 mg AFDM/m². Average daily water temperatures in open and closed canopy blocks at
6553 Eleven Creek were similar in 1988 but were significantly higher in the open blocks than in the
6554 closed blocks in 1989 ($P < 0.01$). The authors conclude by suggesting a thorough investigation
6555 into the interactions and responses of higher trophic levels to increases in periphyton biomass
6556 production and decreases in allochthonous inputs resulting from removal of riparian vegetation.
6557 Furthermore, the authors point out that the ability of stream segments to retain organic inputs
6558 through in-stream large woody debris may be a more important factor for allochthonous input
6559 processing by stream biota than the amount of allochthonous inputs entering a stream.

6560

6561 **Wood Recruitment and Retention**

6562

6563 Hough-Snee et al., 2016

6564

6565 Hough-Snee, N., Kasprak, A., Rossi, R.K., Bouwes, N., Roper, B.B., Wheaton, J.M., 2016.
6566 Hydrogeomorphic and Biotic Drivers of Instream Wood Differ Across Sub-basins of the
6567 Columbia River Basin, USA. *River Research and Applications* 32, 1302–1315.
6568 <https://doi.org/10.1002/rra.2968>

6569

6570 The purpose of this study was to understand the hydrogeomorphic and ecological processes
6571 which lead to wood recruitment and retention in seven sub-basins of the interior Columbia River
6572 Basin (CRB), USA. To achieve this, in-stream wood volume and frequency are quantified across
6573 sub basins. Following this, the riparian, geomorphic, and hydrologic attributes which are most
6574 strongly correlated to in-stream wood loads were determined. Random forest models were used
6575 to identify relationships between ecological and hydrogeomorphic attributes that influence in-
6576 stream wood within each sub-basin. Non-metric multidimensional scaling was performed on a
6577 matrix of hydrogeomorphic and forest cover variables, excluding instream wood frequency and
6578 volume to visualize reaches and sub-basins' relative similarity. To determine how wood
6579 predictors differed between sub-basins, ordinary least squares regression models of wood volume
6580 and frequency were built within each sub-basin. Results from this study show that in stream
6581 wood volume and frequency were distinctly different across all seven sub-basins. Across the
6582 CRB, wood frequency ranged from 0 to 2117.0 pieces km⁻¹, while volume ranged from 0 to 539
6583 m³ km⁻¹. Large wood volume (PERMANOVA $F= 5.1$; $p = 0.001$) and frequency
6584 (PERMANOVA $F= 5.4$; $p = 0.001$) differed significantly between sub-basins. According to
6585 random forest (RF) models, mean annual precipitation, riparian large tree cover, and individual
6586 watershed were the three most important predictors of wood volume and frequency. Watershed

6587 area was the fourth strongest predictor of wood frequency, while catchment-scale and reach-scale
6588 forest cover were the fourth and fifth strongest predictor of wood volume. In contrast, sinuosity
6589 and measures of streamflow and stream power were relatively weak predictors of wood volume
6590 and frequency. Taken together, wood volume and frequency increased with precipitation and
6591 large riparian tree cover and decreased with watershed area. Final RF models explained 43.5% of
6592 the variance in volume and 42.0% of the variance in frequency of in stream wood loads. Results
6593 for drivers of wood frequency and volume between sub-basins were highly variable either
6594 showing no relationship between candidate models and predictive power (e.g., $r^2 \leq 0.12$; Entiat
6595 sub-basin). The highest predictive models for wood volume ($r^2 > 0.55$) and wood frequency (r^2
6596 ≤ 0.45) were for the John Day sub basin. Depending on the sub basin wood volume and
6597 frequency was positively correlated with forest cover, watershed area, large tree cover, 25-year
6598 flood event stream power, riparian conifer cover, and precipitation. Negative correlations,
6599 depending on sub basin, of wood volume and frequency with baseflow discharge, riparian woody
6600 cover, watershed area, and large tree cover. Given the heterogeneous results across all sub-basins
6601 studied, the authors conclude by emphasizing the importance of incorporating local data and
6602 context when building wood models to inform future management decisions.

6603

6604 **Stream Temperature**

6605

6606 Hunter, 2010 (not in focal, treatments and results not relevant to questions)

6607

6608 Hunter, M.A., 2010. Water Temperature Evaluation of Hardwood Conversion Treatment Sites
6609 Data Collection Report (Data Collection Report). Cooperative Monitoring, Evaluation, and
6610 Research (CMER). Fp_cmer_05_513

6611

6612 The purpose of this study is to evaluate the response of stream temperature to changes in canopy
6613 cover using a before-after-control-impact design. This study took place along nine hardwood-
6614 dominated riparian stands in Western Washington. Variables measured among locations and years
6615 include riparian conditions, canopy cover, channel dimensions, substrate, flow and stream
6616 temperature. Results from this study show that hardwood conversion buffers (HCB -
6617 approximately 15 m width) intended to convert hardwood-dominated riparian areas to conifer-
6618 dominated riparian areas usually resulted in decreased canopy cover of streams. Mean Global
6619 Site Factor (GSF - the proportion of global radiation under a plant canopy relative to the amount
6620 in an open area) increased in most study sites with HCB's. However, mean GSF did not change
6621 substantially at sites with buffers closer to standard (~ 18 – 45 m) non-hardwood conversion
6622 buffers. Temperature was highly variable over time and among locations suggesting stream
6623 temperature is affected by many factors that might differ among locations and throughout time.
6624 Longitudinal patterns of warming and cooling were consistent at all sites indicating the potential

6625 importance of careful site selection to account for changes in the longitudinal distribution of
6626 temperatures.

6627

6628 **Influence of Stream Geomorphology on Water Temperature**

6629

6630 Hunter & Quinn, 2009

6631

6632 Hunter, M.A., Quinn, T., 2009. Summer Water Temperatures in Alluvial and Bedrock Channels
6633 of the Olympic Peninsula. *Western Journal of Applied Forestry* 24, 103–108.
6634 <https://doi.org/10.1093/wjaf/24.2.103>

6635

6636 The purpose of this study was to understand how stream geomorphology influences water
6637 temperature in managed stands on the Olympic Peninsula, Washington. Sites chosen for this
6638 included an alluvial study site and a bedrock study site whose overall characteristics were
6639 otherwise comparable apart from geomorphology. The alluvial study site was a 1.6-km reach of
6640 Thorndyke Creek. The bedrock study site was a 1.4-km reach of the South Fork Pysht River.
6641 Both channels were located in 35–50-year-old managed forests dominated by Douglas-fir
6642 (*Pseudotsuga menziesii*) in the uplands and red alder (*Alnus rubra*) in the riparian zone. Surface
6643 substrate at the alluvial channel was composed mostly of gravel, whereas the bedrock channel
6644 was composed of mostly bedrock, boulder, and cobble. The mean solar input (GSF: global site
6645 factor) did not differ between streams. Water temperature was recorded at 75-m intervals along
6646 each channel during the summers of 2003 and 2004. Results from this study show consistent
6647 differences in stream temperature response in alluvial versus bedrock channels. Seasonal
6648 maximum and minimum average daily temperatures varied less at the alluvial site compared to
6649 the bedrock site. This, the authors suggest may be due to hyporheic exchange in alluvial channels
6650 helping to buffer surface water temperatures from gaining or losing heat. In addition,
6651 groundwater may also contribute to the increased stability at the alluvial site. Two same-day
6652 measurements at each site showed the alluvial site gaining 8% of its flow, as compared to the
6653 bedrock site whose flow decreased by approximately 15%. The bedrock site was also shown to
6654 have the highest variation in reach-scale water temperatures during low flow. The authors
6655 conclude that stream geomorphology may have profound impacts on spatial and temporal
6656 patterns of channel water temperature. The authors suggest temperature reading from a single
6657 location may not accurately represent the entire channel. Additional research involving collection
6658 of temporal and longitudinal data will be needed to tailor riparian buffers to channel type.

6659

6660 **Stream temperature, sediment, nutrient**

6661

6662 Murray et al., 2000

6663

6664 Murray, G. L. D., Edmonds, R. L., & Marra, J. L. (2000). Influence of partial harvesting on
6665 stream temperatures, chemistry, and turbidity in forests on the western Olympic Peninsula,
6666 Washington. Northwest science., 74(2), 151-164. Handle: <https://hdl.handle.net/2376/1065>

6667

6668 This study investigates the effects of partial watershed harvest (7-33%) on stream temperature,
6669 chemistry, and turbidity relative to an unharvested old-growth watershed in the western Olympic
6670 Peninsula, Washington. Both harvested watersheds (Rock and Tower creeks) originally contained
6671 old-growth forests. Rock Creek had 7% of its watershed harvested in 1981, and Tower Creek had
6672 33% of its watershed harvested between 1985 and 1987. Logging extended to the stream edge
6673 near the in-stream monitoring sites. Data for stream daily maximum, minimum, and mean
6674 temperatures, chemistry, and turbidity was recorded and monitored from June 1996 to June 1998
6675 (10-15 years post-harvest). Differences in variables between treatment and reference watersheds
6676 were compared with a one-way ANOVA with a posthoc Tukey HSD test. Results showed higher
6677 maximum summer stream temperatures (15.4 °C), and lower winter maximum stream
6678 temperatures (3.7 °C) in the two treatment watersheds compared to the unharvested reference
6679 watershed (12.1 °C and 6.0 °C for summer max, and winter max, respectively). Winter minimum
6680 temperatures for one of the harvested watersheds reached 1.2 °C (Rock Creek) compared to a
6681 winter minimum of 6 °C Thus, seasonal variation of stream maximum temperatures and winter
6682 minimum temperatures were more extreme in the treatment watershed than in the control. There
6683 were no seasonal patterns or significant differences between watersheds in stream chemistry
6684 except for calcium and magnesium concentrations being consistently higher in the unharvested
6685 watersheds. Turbidity was low and not significantly different between watersheds. The authors
6686 interpret these results as evidence of partial harvest having minimal impact on stream
6687 temperatures, chemistry, and turbidity long-term (after 10-15 years). The stream temperature
6688 changes were significant but did not exceed the 16 °C threshold used as a standard for salmonid
6689 habitat. However, there was no data collection during the first decade following harvest.

6690

6691 **Channel Habitat, Particle Size, Stream Temperature, and Woody Debris Response to**
6692 **Harvest**

6693

6694 Jackson et al., 2001

6695

6696 Jackson, C.Rhett., Sturm, C.A., Ward, J.M., 2001. Timber Harvest Impacts on Small Headwater
6697 Stream Channels in the Coast Ranges of Washington1. JAWRA Journal of the American Water
6698 Resources Association 37, 1533–1549.

6699 <https://doi.org/10.1111/j.1752-1688.2001.tb03658.x>

6700

6701 The purpose of this study was to evaluate changes in stream temperature, particle size
6702 distributions of bed material, and channel habitat distributions in 15 first- or second order
6703 streams located on the Coast Range of Western Washington. Four of the fifteen stream basins
6704 were not harvested and served as references; three streams were cut with unthinned riparian
6705 buffers; one with a partial buffer; one with a buffer of non-merchantable trees; and six were
6706 clearcut to the stream edge. Buffer widths varied by operation; the average buffer width varied
6707 from 15 – 21 meters. The narrowest buffer measured on one side of the stream was 2.3 meters.
6708 Data for woody debris, sediment concentrations, turbidity, and stream temperatures were
6709 recorded for one-year prior to harvest (1998). Harvest was conducted in the spring and early
6710 summer of 1999, and post-harvest data was collected for about a month after operations were
6711 complete. Thus, the results presented in this study represent changes in stream attributes and
6712 characteristics immediately following harvest. Results from this study show that logging without
6713 buffers had immediate and dramatic effects on channel morphology. Without buffers, and the
6714 relatively steep topography of the study sites logging debris tended to accumulate at the bottom
6715 of slopes thereby burying or covering many headwater streams. Covered channels were defined
6716 in this study as having flow completely obscured by organic debris, but a recognizable channel
6717 still exists below the debris. Buried channel was defined as having so much organic detritus in
6718 the flow cross-section that the channel was no longer definable. Needles, twigs, whole branches,
6719 and logs buried headwater streams with a mean depth of 0.94 meters of organic debris (range:
6720 0.5 - 2.0 meters). Of the clearcut streams the percent of stream buried with organic matter ranged
6721 from 6 to 90%, and the percent covered by organic matter ranged from 8 to 85%. The sum of
6722 buried and covered for each stream ranged from 72 to 100%. On the other hand, most buffered
6723 streams had 0% covered or buried by organic matter post-harvest with the only exception being
6724 one stream that experienced blowdown post-harvest that covered 29% of the stream. While
6725 debris accumulation tended to protect streams from the effects of solar radiation, organic logging
6726 debris was also shown to trap fine sediment in the channels which, in the near term, greatly
6727 reduced downstream sediment movement. As a result of increased roughness and additional bank
6728 failures within the clearcut sites, sediment size shifted towards finer particles growing from 12 to
6729 44 percent. In contrast, particle size distributions continued nearly unchanged in buffered and
6730 reference sites. In the first summer after logging, significant increases were detected in overall
6731 macroinvertebrate densities, collector densities, shredder abundance and biomass, and organic
6732 and inorganic matter accretion. However, these responses were not detected one year following
6733 logging. For stream temperature changes, because the data collection was for such a short period
6734 of time (1-year pre- and 1-month post-harvest), and because the summer of 1999 was much
6735 cooler than 1998, the assessment of harvest effects on stream temperature changes was difficult.
6736 Thus, to interpret significant changes in stream temperatures from pre- to post- harvest, daily
6737 maximum temperatures were plotted against the appropriate reference stream, and a regression
6738 equation was calculated. The slopes of the regression lines were compared with a student's t-test
6739 to determine significant differences. Of the seven clearcut streams, three showed no significant
6740 changes in temperature, one became cooler (-1.1 °C), one became slightly warmer (+0.8 °C), and

6741 the other 2 became warmer or colder depending on location with decreases in temperature
6742 upstream (-2.2 and -1.7 °C) and increases in temperature downstream (+5.2 and +15.1 °C). The
6743 buffered streams had significant but less dramatic changes in temperature with one decreasing in
6744 temperature (-0.3 °C), and 2 increasing in temperature (+1.6 and +2.4 °C). The one site with the
6745 non-merchantable buffer had much higher temperature increases (+3.7 and +6.6 °C). The authors
6746 posit that sites which retained riparian buffers succeeded in keeping debris out of streams as well
6747 as served to protect streambanks from failure or erosion. Some mature trees left within buffers
6748 experienced blow down and spanned the channel. While the clearcut streams had nearly all
6749 canopy cover removed, the buildup of slash and LW in the stream also provided shade and
6750 insulation that caused reductions in stream temperatures, or slight increases with one exception
6751 (+15.1 °C) The authors point out that this study only served to point out immediate effects of
6752 logging on physical channel conditions. Although important, there are still many questions about
6753 how channel conditions will evolve over time.

6754

6755 **LW**

6756

6757 Meleason et al., 2003

6758

6759 Meleason, M. A., Gregory, S. V., & Bolte, J. P. (2003). Implications of riparian management
6760 strategies on wood in streams of the Pacific Northwest. *Ecological Applications*, 13(5), 1212-
6761 1221. <https://doi.org/10.1890/02-5004>

6762

6763 This study used simulation modeling to evaluate the potential effects of three different riparian
6764 and watershed harvest scenarios on the standing stock of large wood in a hypothetical stream in
6765 the Pacific Northwest. The three scenarios involved harvest 1) clearcut to the streambank, 2)
6766 riparian management buffer widths ranging from 6-75 m, and 3) riparian buffers of various
6767 widths with upland forest plantation. The effects of each scenario on wood load dynamics were
6768 simulated with OSU STREAMWOOD for four harvest rotation periods (no harvest, 60, 90, and
6769 120 years) over the course of 720 years. Results for scenario one (clear-cut to stream) showed
6770 minimal accumulation of wood into the stream with little change over time due to the lack of a
6771 forested riparian management zone. Results for scenario two showed the maximum standing
6772 stock of in-stream wood loads required ≥ 30 m no-cut buffer zones for 500-year-old forests.
6773 Wood loads in streams with 6 m wide buffers showed 32% of standing wood load stocks after
6774 240 years. Results from scenario three showed minimal amounts of wood contributed into
6775 streams from forest plantations when > 10 m wide buffers were used. The authors interpret these
6776 results as evidence that riparian buffer widths and forest age are more important for estimating
6777 changes in wood loads over time than the harvest rotation age of plantation forests.

6778

6779 **LW**

6780

6781 Martin & Grotefendt, 2007

6782

6783 Martin, D. J., & Grotefendt, R. A. (2007). Stand mortality in buffer strips and the supply of
6784 woody debris to streams in Southeast Alaska. *Canadian Journal of Forest Research*, 37(1), 36-49.
6785 <https://doi.org/10.1139/x06-209>

6786

6787 This study compared riparian stand mortality and in-stream LW recruitment characteristics
6788 between riparian buffer strips with upland timber harvest and riparian stands of unharvested
6789 watersheds using aerial photography. This study was conducted in the northern and southern
6790 portions of Southeast Alaska at multiple sites in nine timber harvest areas. All study sites were
6791 along moderate- and low-gradient streams with channel widths ranging from 5 m to 30 m wide.
6792 All buffer strips were conifer dominated and a minimum of 20 m wide that included selective
6793 harvest within the 20 m zone. Reference sites were along unharvested reaches in the same area.
6794 Stand mortality was estimated by the proportion of downed trees within a buffer strip.
6795 Differences in downed tree proportions relative to reference streams were assumed to be caused
6796 by timber harvest, accounting for selective in-buffer harvests. A one-tailed paired t-test or a
6797 Wilcoxon signed rank test was used to check for statistical differences between treatment and
6798 reference sites. Results showed significantly higher mortality (based on cumulative stand
6799 mortality: downed tree counts divided by standing tree counts + downed tree counts),
6800 significantly lower stand density (269 trees/ha in buffer units and 328 trees/ha in reference units),
6801 and a significantly higher proportion of LW recruitment from the buffer zones of the treatment
6802 sites than in the reference sites. Densities within all units ranged from 0 – 1334 trees/ha
6803 depending on location. Overall, mean stand density in the buffer units was 18% lower than in the
6804 reference units. Results also showed that mortality varied with distance to the stream.
6805 Differences in mortality for the treatment sites were similar to the reference sites for the first 0-
6806 10 m from the stream (only a 22% increase in the treated sites). However, mortality in the outer
6807 half of the buffers (10-20 m) from the stream in the treatment sites was more than double (120%
6808 increase) what was observed in the reference sites. This caused a change in the LW recruitment
6809 source distance curves, with a larger proportion of LW recruitment coming from greater
6810 distances in logged watersheds. LW recruitment based on the proportion of stand recruited (PSR)
6811 was significantly higher in the buffered units compared to the reference units. However, PSR
6812 from the inner 0-20 m was only 17% greater in the buffer units than in the reference units, while
6813 PSR of the outer unit (10 – 20 m) was more than double in the buffered units than in the
6814 reference units. The researchers conclude that the increase in mortality was caused by an
6815 increased susceptibility to windthrow. They estimate that future recruitment potential from the
6816 logged sites diminished by 10% relative to the unlogged reference sites.

6817

6818 **Stream temperatures**

6819

6820 Macdonald et al., 2003b

6821

6822 Macdonald, J. S., MacIsaac, E. A., & Herunter, H. E. (2003). The effect of variable-retention
6823 riparian buffer zones on water temperatures in small headwater streams in sub-boreal forest
6824 ecosystems of British Columbia. *Canadian journal of forest research*, 33(8), 1371-1382.
6825 <https://doi.org/10.1139/x03-015>

6826

6827 This study investigates the impacts of forest harvest on stream temperatures under three variable
6828 retention buffer treatments in headwater streams of the interior sub-boreal forests of British
6829 Columbia. Temperature data were recorded for two years pre- and five years post-harvest from
6830 five harvested streams and two unharvested reference streams. Differences between pre- and
6831 post-harvested stream temperatures were compared with the paired reference streams using
6832 repeated measures ANOVA. Treatment riparian areas were harvested with the following
6833 prescriptions: 1) low-retention – removal of all merchantable timber >15 or >20 cm DBH for
6834 pine or spruce respectively, within 20 m of the stream 2) high-retention – removal of
6835 merchantable timber >30 cm DBH within 20-30 m of the stream, and 3) Patch-cut – high
6836 retention for the lower 60% of watershed approaching streams and removal of all vegetation in
6837 the upper 60% of the watershed. Eight first-order streams were included in this study: two

6838 in the Gluskie Creek watershed (G5, G7) and six in the Baptiste Creek watershed (B1–B6). Five
6839 of these streams were within the harvested boundaries (2 high-retention, 2 low-retention, and 1
6840 patch cut), and 3 reaches outside of the harvest boundary served as controls. Results showed a
6841 significant increase in stream temperatures ranging from 4 – 6 °C at five years post-harvest, and
6842 increased ranges of diurnal temperature fluctuations for all treatment streams relative to the
6843 reference streams. Streams that had summer maximum mean weekly temperatures of 8°C before
6844 harvesting had maximum temperatures near 12°C or more following harvesting. Daily ranges of
6845 1.0–1.3°C before harvesting became 2.0–3.0°C following harvesting. Greater temperature ranges
6846 occurred in low-retention and patch treatments than the high-retention or control treatment
6847 locations. The high-retention buffer treatment mitigated temperature increases for the first three
6848 years. Still, increased mortality (windthrow) caused a reduction in the canopy that increased
6849 stream temperatures equivalent to other treatment streams by year five. The results of this study
6850 show evidence that high-retention buffers are no more effective in preserving stream temperature
6851 changes than small retention buffers when treatment areas have a high susceptibility to
6852 windthrow.

6853

6854 **Sediment delivery pathways**

6855

6856 Litschert & MacDonald, 2009

6857

6858 Litschert, S. E., & MacDonald, L. H. (2009). Frequency and characteristics of sediment delivery
6859 pathways from forest harvest units to streams. *Forest Ecology and Management*, 259(2), 143-
6860 150. <https://doi.org/10.1016/j.foreco.2009.09.038>

6861

6862 This study investigates the frequency of sediment delivery pathways (“features”) in riparian
6863 management areas and measures the physical characteristics and connectivity of these pathways
6864 following timber harvest. The results of this study were then used to develop models for
6865 predicting the length and connectivity of pathways formed from harvest units. Data was collected
6866 from over 200 harvest units with riparian management areas in the Eldorado, Lassen, Plumas,
6867 and Tahoe National Forests in the Sierra and Cascade mountains of northern California. Riparian
6868 buffer widths for this area are 90 m and 45 m for perennial and annual streams respectively. No
6869 machinery is allowed in the riparian management areas. Data collected and analyzed for the
6870 pathways included years since harvest, mean annual precipitation, soil depth, soil erodibility,
6871 hillslope gradient, aspect, and elevation. Characteristics of pathway length, gradient, and
6872 roughness were also collected. Relationships between site variables and pathway variables were
6873 assessed using linear regression. The site variables with the most significant relationships with
6874 the pathway variables were used in a multivariate regression model to predict pathway length.
6875 Only 19 of the 200 harvest units had sediment development pathways. Pathways ranged in age
6876 (time since harvest) from 2 to 18 years, and in length from 10 m to 220 m. Of the 19 pathways,
6877 only six were connected to streams, and five of those originated from skid trails. Pathway length
6878 was significantly related to mean annual precipitation, cosine of the aspect, elevation, and
6879 hillslope gradient. The authors conclude that timber prescription practices for these National
6880 Forests are effective in reducing sediment delivery pathways. The authors interpret these results
6881 as evidence that skid trails should be directed away from streams, maintaining surface roughness,
6882 and promptly decommissioning skid trails.

6883

6884 **LW**

6885

6886 Liquori, 2006

6887

6888 Liquori, M. K. (2006). POST-HARVEST RIPARIAN BUFFER RESPONSE: IMPLICATIONS
6889 FOR WOOD RECRUITMENT MODELING AND BUFFER DESIGN 1. *JAWRA Journal of the*
6890 *American Water Resources Association*, 42(1), 177-189. [https://doi.org/10.1111/j.1752-](https://doi.org/10.1111/j.1752-1688.2006.tb03832.x)
6891 [1688.2006.tb03832.x](https://doi.org/10.1111/j.1752-1688.2006.tb03832.x)

6892

6893 This study investigates the differences in treefall characteristics in riparian management areas
6894 based on ecological and physiographic variables to give insight on the variables important for
6895 wood recruitment modeling. Data were collected from 20 riparian buffer sites that had all been
6896 clearcut within three years of sampling with standard no-cut buffers 25 ft. An additional 50-100
6897 ft buffer was applied to fish-bearing streams depending on stream type, in a managed tree farm in
6898 the Cascade Mountains of western Washington. These riparian buffers generally consisted of
6899 naturally regenerated, second-growth conifer stands about 45 to 70 years old. “Very modest”
6900 thinning was applied to some stands to meet wildlife objectives and any downed wood not
6901 affecting the channel was removed. Tree characteristic data collected included tree size (DBH
6902 and height), species, fall direction, tree fall angles, estimated cause of mortality, and distance to
6903 the stream. Site characteristics included stream gradient, valley morphology, and time since
6904 harvest. Tree recruitment probability curves were developed as a function of tree height using
6905 methods described by Beschta, (1990). Results showed that wind-caused mortality and tree fall
6906 rates were significantly higher, up to three times higher, than competition-induced mortality
6907 within buffers for three years following treatment. The median observed treefall per site was
6908 15% of all trees in each buffer, ranging from 1 to 57%. total treefall at each site for one, two, and
6909 three years since harvest was $16 \pm 10\%$, $28 \pm 21\%$, and $10 \pm 10\%$, respectively. Total treefall
6910 percentage for each site was not correlated to years since harvest (Spearman $R = 0.11$; $p = 0.34$).
6911 The mean and standard deviation of the total normalized treefall for one-year old sites was $405 \pm$
6912 394 trees/km ($n = 9$), for two-year old sites was 264 ± 280 trees/km ($n = 7$), and for three-year
6913 old sites was 556 ± 316 trees/km ($n = 4$). Treefall varied significantly by species. Downed red
6914 alder (*Alnus rubra*), western red cedar (*Thuja plicata*), and Douglas-fir (*Pseudotsuga menziesii*)
6915 comprised 3 percent to 8 percent of all downed trees; these species had treefall rates ranging
6916 from 5 percent to 9 percent of the total number of trees of the same species. By contrast, treefall
6917 rates for western hemlock (*Tsuga heterophylla*) and Pacific silver fir (*Abies amabilis*) ranged
6918 from 23 percent to 26 percent. Treefall rates also varied somewhat by size, with the 31 to 41 cm
6919 (12 to 16 in) diameter class having the greatest treefall rates (All trees were grouped into size
6920 classes based on diameter at breast height: 1 to 8 in; 8 to 12 in; 12 to 16 in; 16 to 20 in; and more
6921 than 20 in). Treefall following harvest greatly exceeded the expected competition induced
6922 mortality rates (posited by Franklin, 1970) of 0.5%, and the model of average competition
6923 mortality used in Rainville et al. (1985), which ranged from 0.7 - 1.6%, and 2% per year for bank
6924 undercutting. Treefall direction was heavily biased towards the channel regardless of channel or
6925 buffer orientation and tree fall probability was highest in the outer areas of the buffers (adjacent
6926 to the harvest area). Fall direction bias increased significantly in the inner portions of the buffer.
6927 Within the 0 to 7 m zone and 7 to 15 m zone, 68% and 67% of the trees, respectively, fell toward
6928 the channel ($n = 125$ and 153 , respectively). Only 44% of the outer zone (> 15 m) downed trees
6929 fell toward the channel ($n = 403$). Generally, recruitment was negatively correlated to buffer
6930 width ($r^2 = 0.40$). Treefall was generally highest at the outside edges of buffers (50+ feet),
6931 representing about 60% of the total observed treefall, while the 0–25-foot zone represented
6932 ~18%, and the 25–50-foot zone represented ~22%. The authors interpret their results as evidence
6933 that tree fall models that use a random fall direction may underrepresent the probability of LW
6934 recruitment into streams. Further, they suggest that the increase in windthrow mortality and the
6935 probability of tree fall with increasing distance from the stream should be considered.

6936

6937 **LW**

6938

6939 Lininger et al., 2021 (removed from focal list, this is a case study)

6940

6941 Lininger, K. B., Scamardo, J. E., & Guiney, M. R. (2021). Floodplain large wood and organic
6942 matter jam formation after a large flood: Investigating the influence of floodplain forest stand
6943 characteristics and river corridor morphology. *Journal of Geophysical Research: Earth Surface*,
6944 126(6), e2020JF006011. <https://doi.org/10.1029/2020JF006011>

6945

6946 This study examines how river corridor morphology and forest stand density influence LW and
6947 coarse particulate matter (CPOM) deposition patterns in the flood plain resulting from a 400-year
6948 flood event in West Creek in the Colorado Front Range in 2013. The researchers tested the
6949 hypothesis that if river corridor geomorphology affects LW and CPOM deposition then there
6950 should be an inverse relationship between elevation above and distance from the stream's edge.
6951 Further, that deposition frequency would be higher in unconfined portions of the corridor.
6952 Considering forest stand structure, the researchers hypothesized that LW/CPOM jams would be
6953 pinned by trees, higher in intermediate forest densities, and decrease in size with increasing
6954 forest stand density. Field data of LW/CPOM jams were analyzed with non-parametric Spearman
6955 correlation tests to determine the strength of their relationship with channel and stand
6956 characteristics. Results showed support for most of the hypotheses. LW accumulations did
6957 decrease in size with distance from the stream, but CPOM did not. Confined channels (steeper
6958 reaches) contained fewer LW/CPOM loads per unit area. The authors speculate that these reaches
6959 had higher flow rates and thus lower deposition during the flood. CPOM jams increased in
6960 number per area with increasing stand density with most jams pinned against live trees. The
6961 authors conclude that the effect of riparian forest stand density is evidence that riparian forests in
6962 the floodplains should be preserved to increase LW and CPOM trapping probability.

6963

6964 **Stream Temperature**

6965

6966 Janisch et al., 2012

6967

6968 Janisch, J.E., Wondzell, S.M., Ehinger, W.J., 2012. Headwater stream temperature: Interpreting
6969 response after logging, with and without riparian buffers, Washington, USA. *Forest Ecology and*
6970 *Management* 270, 302–313. <https://doi.org/10.1016/j.foreco.2011.12.035>

6971

6972 The purpose of this study was to assess the stream temperature response to three different
6973 harvesting treatments in small, forested headwater catchments in western Washington. The pre-
6974 logging calibration period lasted 1–2 summers and stream temperatures were monitored for two
6975 or more summers after logging. Harvest treatments occurred between September 2003 and July
6976 2005; catchments were clustered by harvest year for analysis. A before-after-control-impact
6977 study design was used to contrast stream temperature responses for three forest harvest
6978 treatments: clearcut logging to the stream (n=5), a continuous buffer (n=6) with widths 10-15 m
6979 on each side of the channel, and a patched buffered (n=5) where portions of the riparian forests
6980 ~50-110 m long were retained in distinct patches along some portion of the channel with the
6981 remaining riparian area clearcut. For the patch buffers there was no standard width, the buffer
6982 spanned the full width of the floodplain area and extended well away from the stream. Upland
6983 areas adjacent to buffers were clearcut. Regression relationships were developed between
6984 temperatures measured in the treatments and corresponding reference catchments. A simple
6985 ANOVA model was used that only included fixed effects for treatment, years since treatment,
6986 and day of year. Because of the unbalanced experimental design and variation in time of harvest,
6987 clustering of treatments caused the sample sizes to become too small to apply a more complex
6988 nested, repeated measures ANOVA could not be used. Correlation analysis was conducted
6989 between post-harvest stream temperatures and descriptive variables on a subset of catchments to
6990 examine possible factors that might control post-harvest thermal responses. Results from this
6991 study show significant increases in stream temperature in all treatments. Although temperature
6992 responses were highly variable within treatments, July and August daily maximum temperatures
6993 increased in clearcut catchments during the first year after logging by an average of 1.5°C (range
6994 0.2 to 3.6°C), in patch-buffered catchments by 0.6°C (range – 0.1 to 1.2°C), and in continuously
6995 buffered catchments by 1.1°C (range 0.0 to 2.8°C). Canopy cover in all streams averaged 95%
6996 prior to harvest and did not differ between treatment and reference streams. Following treatment,
6997 canopy cover in the clearcut catchments averaged 53%, canopy cover in the patch buffer
6998 treatment averaged 76%, and canopy cover in the continuous buffer treatment averaged 86%.
6999 Following treatment, the canopy cover of the clearcut and patch buffer treatments were
7000 significantly lower than in the reference streams. The continuous buffer treatments did not differ
7001 significantly from the reference streams for canopy cover. Further analyses which attempted to
7002 identify variables responsible for controlling the extent of stream temperature responses showed
7003 the amount of cover retained in the riparian buffer was not a strong explanatory variable. Post-
7004 treatment temperature changes suggested that treatments ($p = 0.0019$), the number of years after
7005 treatment ($p = 0.0090$), and the day of the year ($p = 0.0007$) were all significant effects
7006 explaining observed changes in temperature. Wetland area ($r^2 = 0.96$, $p < 0.01$) and length of
7007 surface flow ($r^2 = 0.67$, $p = 0.05$) were strongly correlated with post-logging temperature
7008 changes. Regression analysis of these variables showed streams with fine-textured substrates
7009 responded differently than coarse textured substrates. The authors speculate this is possibly due
7010 to groundwater interactions which can buffer thermal responses of small streams. In summary,
7011 the authors conclude that their results suggest small headwater streams may be fundamentally
7012 different than larger streams partly because factors other than canopy shade can greatly influence
7013 stream energy budgets to moderate stream temperatures despite changes and/or removal of the
7014 overstory canopy.

7015

7016 **Large woody debris**

7017

7018 Jones et al., 2011 (Removed from focal list, study not relevant to focal questions)

7019

7020 Jones, T.A., Daniels, L.D., Powell, S.R., 2011. Abundance and function of large woody debris in
7021 small, headwater streams in the Rocky Mountain foothills of Alberta, Canada. *River Research*
7022 *and Applications* 27, 297–311. <https://doi.org/10.1002/rra.1353>

7023

7024 The purpose of this study was to assess LW abundance in the upper foothills of the Rocky
7025 Mountains in Alberta, Canada. This study also sought to understand key processes that underlie
7026 changes in LW function. Finally, this study used results to develop a LW recruitment, decay and
7027 interaction model. This research was conducted in 21 headwater streams spanning two
7028 watersheds. At each site, all LW was sampled and was classified according to decay, orientation,
7029 position and function. LW frequency, total volume, and total in-stream volume were calculated
7030 and analyzed for differences using a one-way ANOVA followed by a Tukey post hoc test to
7031 differentiate among significant classes. Results show LW frequency was greater in the Alberta
7032 foothills (64.0 ± 3.3 LW 100 m¹) than in many small, headwater streams in mountain (46.2 ± 3.6),
7033 coastal (47.6 ± 3.8), mixed broad-leaf (47.0 ± 4.2) and boreal (31.0 ± 3.0) streams. This, the
7034 authors suggest, is likely due to the narrow bankfull width channels characteristic of the Alberta
7035 foothills which are less able to transport LW downstream. LW with ≥ 20 cm was more frequent in
7036 coastal streams, and overall LW volume was also greatest in coastal streams (721.0 ± 99.9 m³ ha⁻¹).
7037 The authors note that large LW volumes in coastal streams are likely due to geomorphic
7038 disturbances alongside large, long-lived, decay resistant tree species. According to Harmon et al.
7039 1986, much of the variation in LW recruitment is due to differences in species life history and
7040 forest type which together govern log size and decay rates.

7041

7042 **Suspended Sediment**

7043

7044 Karwan et al., 2007

7045

7046 Karwan, D., Gravelle, J., Hubbart, J., 2007. Effects of timber harvest on suspended sediment
7047 loads in Mica Creek, Idaho. *Forest Science* 53, 181–188.
7048 <https://doi.org/10.1093/forestscience/53.2.181>

7049

7050 The purpose of this study was to examine the effects of forest road construction and timber
7051 harvest on total suspended solids (TSS) in a forested watershed. This study took place at the
7052 Mica Creek Experimental Watershed in northern Idaho. The study area consisted of dense,
7053 naturally regenerated, even-aged stands ~65 years old and ~300 trees per acre. Timber harvesting
7054 and heavy road use began in 2001. Treatments in the paired-watershed experiment consisted of
7055 (1) commercial clearcut of the watershed area of 50%, and was broadcast burned and replanted
7056 by the end of May 2003, (2) partial cut in which half the canopy was removed in 50% of the
7057 watershed in 2001, with final 10% of log processing and hauling in early summer of 2002. and
7058 (3) a no-harvest control. All harvests were carried out according to best management practices
7059 and in accordance with the Idaho Forest Practices Act. At the time of the study this involved a
7060 22.86 m (75 ft) stream protection zones (SPZs) on each side of fish-bearing (Class I) streams.
7061 The inner 50 ft is an equipment exclusion zone where no ground-based skidding machinery is
7062 allowed. Timber harvesting is allowed in Class I SPZs, but 75% percent of existing shade must
7063 be retained. Along non-fish-bearing (Class II) streams, harvesting equipment was excluded from
7064 entering within 9.14 m (30 ft) of definable stream channels and any cut trees were felled away
7065 from the stream; however, there were no tree retention requirements. In the clearcut and partial
7066 cut units, line skidding was used on slopes in the watershed exceeding approximately 20%, while
7067 tractor skidding was used on the lower gradient slopes. On all skid trails, drainage features, such
7068 as water bars, were installed for erosion control at the end of the harvest period. Time series data
7069 were compiled for all measured TSS values from 1991 through 2004. Data was collected via
7070 seven stream monitoring flumes located within the Mica Creek Watershed. Monthly TSS loads
7071 were compared across watersheds for five time intervals: (1) pretreatment: ~6 years, (2)
7072 immediate post-road construction: ~1 year, (3) recovery post-road construction: ~3 years, (4)
7073 immediate post-harvest: ~1 year, and (5) recovery post-harvest: ~3 years. Trends in the
7074 relationship between treatment and control watersheds were statistically examined for each of the
7075 time intervals. Treatments in the paired-watershed experiment consisted of (1) commercial
7076 clearcut of the watershed area of 50%, and was broadcast burned and replanted, (2) partial cut in
7077 which half the canopy was removed in 50% of the watershed (3) a no-harvest control. All
7078 harvests were done according to best management practices and the Idaho Forest Practices Act.
7079 This included equipment exclusion zones of 50- and 30-feet for fish- and non-fish-bearing
7080 streams, respectively. On all skid trails, drainage features, such as water bars, were installed for
7081 erosion control at the end of the harvest period. Analysis of covariance was used for each
7082 treatment-control watershed pair. Results show monthly TSS loads from watersheds 1 (clearcut),
7083 2 (partial cut), and 3 (no-harvest) ranged from 0.4 kg km⁻² to above 10,000 kg km⁻², with a
7084 maximum in the spring months and minimum in the winter and late summer months similar to
7085 intra-annual trends in water yield. Road construction in both watersheds did not result in
7086 statistically significant impacts on monthly sediment loads in either treated watershed during the
7087 immediate or recovery time intervals. A significant and immediate impact of harvest on monthly
7088 sediment loads in the clear-cut watershed ($p = 0.00011$), and a marginally significant impact of
7089 harvest on monthly sediment loads in the partial-cut ($p = 0.081$) were observed. Total sediment
7090 load from the clearcut over the immediate harvest interval exceeded predicted load by 152%
7091 (6,791 kg km⁻²); however, individual monthly loads varied around this amount. The largest
7092 increases in percentage and magnitude occurred during snowmelt months, namely April 2002

7093 (560%, 2,958 kg km⁻²) and May 2002 (171%, 3,394 kg km⁻²). Neither treatment showed a
7094 statistical difference in TSS during the recovery time (clearcut: p = 0.2336; partial-cut: p =
7095 0,1739) compared to calibration loads (pre-treatments). The authors conclude that best
7096 management practices for road construction, including improvement of existing roads, did not
7097 produce significant changes in TSS. Significant changes in TSS only occurred immediately after
7098 harvest. However, after one year, the TS load became statistically indistinguishable from the
7099 control.

7100

7101 **Harvest effects on Instream light**

7102

7103 Kaylor et al., 2017

7104

7105 Kaylor, M.J., Warren, D.R., Kiffney, P.M., 2017. Long-term effects of riparian forest harvest on
7106 light in Pacific Northwest (USA) streams. *Freshwater Science* 36, 1–13.
7107 <https://doi.org/10.1086/690624>

7108

7109 The purpose of this study was to evaluate relationships between riparian forest stand age and
7110 stream light availability. The specific goals dealt with evaluating characteristics of late-
7111 successional forest light regimes, and whether canopy openness and light differed between
7112 streams flowing through harvested units and late-successional forest units. This study took place
7113 at the HJ Andrews Experimental Forest in the Cascade Mountain, Oregon. Approximately 11.5
7114 km of stream length were sampled in the McCrae Basin which consists mostly of old-growth
7115 forests Douglas-fir forests with small patch clear cuts. All treatment sites were harvested within
7116 50 to 60 years before the study. Clearing up to both stream banks occurred at two of seven
7117 treated sites and clearing up to one bank occurred on all other treated sites. Stream bank-full
7118 width, wetted width, canopy openness, % red alder, and estimated photosynthetically active
7119 radiation (PAR) were quantified at 25-m intervals to evaluate relationships between channel and
7120 riparian characteristics and stream light. Results from this study show mean estimated PAR
7121 reaching the streams was lower in the recovering harvested units (50-year post-treatment) than
7122 in up and downstream reaches bordered by old growth for all comparisons (n=14), while only 6
7123 were significant (p<0.05). All in all, old growth reaches averaged 1.7 times greater PAR values
7124 than in nearby harvested units with the greatest differences occurring when harvest was
7125 implemented on both banks. Mean canopy openness was higher in late-successional forests (>
7126 300 years old) than in young second growth forests (30–100-year-old forests), 18% and 8.7%
7127 respectively. Results also indicate the relationship between canopy openness and PAR was
7128 stronger at the reach scale than at individual locations with mean canopy openness explaining
7129 78% of the variance in mean PAR estimates. The researchers also conducted a review of
7130 available literature of studies that contained information on the effects of Northwest Douglas-fir
7131 forest growth dynamics on canopy cover and light availability. The researchers concluded from

7132 this review that canopy closure, and thus lower light availability, occurs approximately 30 years
7133 after growth and maintained until after 100 years of growth when the canopy structure begins to
7134 open and produce gaps. Altogether, this study suggests stream light regimes are affected by
7135 initial canopy removal and subsequent recovery. Depending on forest type, dominant species and
7136 the age of the stand, different stages of stand development may reflect complex overstory
7137 structures allowing variable levels of light to the stream.

7138

7139 **Stream Temperatures**

7140

7141 Kibler et al., 2013

7142

7143 Kibler, K.M., Skaugset, A., Ganio, L.M., Huso, M.M., 2013. Effect of contemporary forest
7144 harvesting practices on headwater stream temperatures: Initial response of the Hinkle Creek
7145 catchment, Pacific Northwest, USA. *Forest Ecology and Management* 310, 680–691.
7146 <https://doi.org/10.1016/j.foreco.2013.09.009>

7147

7148 The purpose of this study was to investigate the effects of contemporary forest harvesting
7149 practices on headwater stream temperatures using a BACI design. This study was conducted as
7150 part of the Hinkle Creek paired Watershed Study (HCPWS). This study consisted of a nested,
7151 paired watershed study in which harvesting treatments in accordance with the Oregon Forest
7152 Practices Act (FPA) were applied to four headwater catchments in southern Oregon. Oregon FPA
7153 does not require retention of fixed-width buffer strips adjacent to non-fish-bearing streams. Thus,
7154 as a part of the harvest activities, fixed-width buffer strips containing merchantable overstory
7155 conifers were not left adjacent to the non-fish-bearing streams. Clearcut harvest took place
7156 between August 2005 and May 2006. Streamflow and temperature were measured at 8 locations
7157 within the basin from autumn 2002 until autumn of 2006 giving 3 years of pre-harvest data and
7158 <1 year of post-harvest data. Treatment and reference catchments were paired based on similarity
7159 in catchment area, aspect, stream orientation, stream length, and discharge. Significant
7160 differences between pre- and post-harvest daily max temperature measurements were detected
7161 across all sites, however, magnitude and direction of changes were inconsistent. Results for daily
7162 mean maximum stream temperatures show a variable response across all four harvested streams
7163 ranging from 1.5°C cooler to 1.1°C warmer relative to pre-harvest years. No statistically
7164 significant changes in max, mean, or minimum daily stream temperatures to timber harvest were
7165 observed. The authors suggest possible explanations for lack of consistent temperature increases
7166 to shading provided by logging slash. Interestingly, statistically significant changes to
7167 relationship between treatment and reference site pairs with respect to minimum and mean
7168 stream temperatures resulted in decreased minimum daily stream temperatures on days where
7169 high temperatures were observed in reference streams. At one treatment site, mean minimum
7170 temperatures across the warm season decreased 1.9°C relative to pre-harvest years, and the

7171 minimum temperature on the warmest day decreased by 2.8°C relative to pre-harvest years.
7172 Except for one treatment-reference pair, highly significant changes to slope and intercept
7173 parameters of minimum daily stream temperatures were detected for each stream pair ($p < 0.001$).
7174 The authors suggest decreases in daily minimum stream temperature is a likely consequence of
7175 timber harvest.

7176

7177 **Shade and Stream temperature**

7178

7179 Cupp & Lofgren, 2014

7180

7181 Cupp, C.E. & Lofgren, T.J. (2014). Effectiveness of riparian management zone prescriptions in
7182 protecting and maintaining shade and water temperature in forested streams of Eastern
7183 Washington. Cooperative Monitoring Evaluation and Research Report CMER 02-212.
7184 Washington State Forest Practices Adaptive Management Program. Washington Department of
7185 Natural Resources, Olympia, WA.

7186

7187 The purpose of this study was to assess the percent reduction in canopy cover, and the response
7188 in stream temperatures following riparian timber harvest under the “all available shade” rule
7189 (ASR), and the standard rule (SR) in eastern Washington. The ASR is applied to areas in the Bull
7190 Trout Habitat Overlay (BTO; map of bull trout habitat) that requires retention of all available
7191 shade within 75 feet of the stream. Under the standard shade rule (SR) some harvest is allowed
7192 within the 75-foot buffer depending on elevation and pre-harvest canopy cover. The primary
7193 objectives of this study were to (1) Quantify and compare differences in post-harvest canopy
7194 closure between the SR and the ASR riparian prescriptions of eastern Washington; and (2)
7195 Quantify and compare differences in stream temperature effects of the two riparian prescriptions:
7196 the SR and the ASR. This study was conducted at 30 sites in eastern Washington. Sites were
7197 between 65-100 years old and were situated along second to fourth order streams with harvest-
7198 regenerated or fire-regenerated forests. Reference reaches were located upstream from treatment
7199 reaches where harvest was applied. Eighteen sites were located on state owned and managed
7200 forests and 12 sites were located on private industrial forests. Prior to harvest treatments, canopy
7201 closure measurements ranged from 89% to 97%, with a mean of 93%. The riparian management
7202 zone (RMZ) consists of three zones: The core zone is nearest to the edge of the stream and
7203 extends out 30 feet horizontally from the bankfull edge or outer edge of the channel migration
7204 zone (CMZ), whichever is greater. The inner zone is situated immediately outside of the core
7205 zone. For streams with a bankfull width of less than or equal to 15 feet wide, the inner zone
7206 width is 45 feet wide. All streams assessed in this study were less than or equal to 15 feet wide.
7207 The outer zone of the RMZ is the zone furthest from the water and its width varies according to
7208 stream width and site class for the land. The specific site class (a measure of site productivity) at
7209 each treatment site would vary the outer zone width from 0 to 55 feet wide. Seven sites had up to

7210 four years pre-harvest temperature data with only two years post-harvest data. Nine sites had
7211 three years pre-harvest data and one site had only one year pre-harvest data. The remaining 13
7212 sites had two years pre-harvest data. Following harvest treatments, all 30 sites had at least two
7213 years post-harvest temperature data collection, although 21 of the 30 sites had at least three years
7214 post-harvest monitoring. Data collection included twice hourly stream and air temperature data
7215 during each sample period. Canopy, shade, riparian, and channel data were collected during the
7216 first-year pre-harvest and the first year post-harvest. Stream temperature data were collected at
7217 30-minute intervals between 1 July and 15 September for a total of 77 days each year a site was
7218 investigated. Stream canopy closure and shade were quantified at 75-ft intervals within each
7219 reach using a hand-held densiometer (for canopy closure measurements) and a self-leveling
7220 fisheye lens digital camera (for shade measurements). A t-test was used to evaluate differences in
7221 pre-harvest canopy cover between reference and treatment reaches, and between ASR and SR
7222 sites. A correlation analysis between post-harvest change in shade and the descriptive riparian
7223 and channel values (e.g., trees per acre, basal area, channel gradient, etc.) was also used to
7224 examine possible factors that may control post-harvest changes in shade. A linear mixed effects
7225 model was used to quantify and compare differences in daily max stream temperatures (DMAX)
7226 between no harvest, ASR and SR prescriptions. Results showed post-harvest shade values
7227 decreased in SR sites (mean effect of -2.8%, $p = 0.002$), as did the canopy closure values (mean
7228 effect of -4.5%, $p < 0.001$). Shade and canopy closure values did not significantly change in the
7229 treatment reaches of the ASR sites. Mean shade reduction in the SR treatment sites exceeded the
7230 mean shade reduction in the ASR sites by 3%. Canopy closure reduction was also greater in the
7231 SR sites than in the ASR sites by a mean of 4%. Specifically, the mean shade reduction in ASR
7232 sites was 1% with a maximum reduction of 4%. The mean reduction of shade in the SR sites was
7233 4% with a maximum reduction of 10%. Mean shade contribution of upland trees (trees outside of
7234 the RMZ) per study site was calculated as $< 1\%$. Shade reduction levels did not differ between
7235 the sites receiving RMZ-harvest only and the sites receiving standard operational upland harvest.
7236 Site seasonal means of daily maximum stream temperature treatment responses in the first two
7237 years following harvest ranged from $-0.7\text{ }^{\circ}\text{C}$ to $0.5\text{ }^{\circ}\text{C}$ in the ASR reaches and from -0.3 to 0.6 in
7238 the SR reaches. Site seasonal mean post-harvest background responses in reference reaches
7239 ranged from $-0.5\text{ }^{\circ}\text{C}$ to $0.6\text{ }^{\circ}\text{C}$ in the first two years following harvest. Mean daily maximum
7240 stream temperature increased $0.16\text{ }^{\circ}\text{C}$ in the SR harvest reaches, whereas stream temperatures in
7241 both the ASR sites and in the no-harvest reference reaches increased on average by $0.02\text{ }^{\circ}\text{C}$.
7242 Seasonal mean stream temperature responses of up to $0.5\text{ }^{\circ}\text{C}$ in the no-harvest references were
7243 common during the post-harvest test period. Sample period means of daily maximum
7244 temperature responses varied from $-1.1\text{ }^{\circ}\text{C}$ to $0.7\text{ }^{\circ}\text{C}$ in the first two years post-harvest for the
7245 ASR sites, from -0.5 to $0.8\text{ }^{\circ}\text{C}$, in the SR sites, and -0.5 to $0.9\text{ }^{\circ}\text{C}$ in the reference sites. The
7246 authors interpret these results as evidence that temperature effects of the SR, and ASR were
7247 similar to reference conditions along sampled reaches for small streams in the mixed fir zone
7248 mid-successional forests of eastern Washington. Further, that processes not directly related to
7249 canopy cover alteration over streams may be primarily responsible for the small variations
7250 observed in stream temperatures following harvest.

7251

7252
7253 Ehinger et al., 2021
7254
7255 Ehinger, W.J., W.D. Bretherton, S.M. Estrella, G. Stewart, D.E. Schuett-Hames, and S.A. Nelson.
7256 2021. Effectiveness of Forest Practices Buffer Prescriptions on Perennial Non-fish-bearing
7257 Streams on Marine Sedimentary Lithologies in Western Washington. Cooperative Monitoring,
7258 Evaluation, and Research Committee Report CMER 2021.08.24, Washington State Forest
7259 Practices Adaptive Management Program, Washington Department of Natural Resources,
7260 Olympia, WA.
7261
7262 The purpose of this study was to assess the effectiveness of riparian management zone
7263 prescriptions in maintaining functions and processes in headwater perennial, non-fish-bearing
7264 streams in incompetent (easily eroded) marine sedimentary lithologies in western Washington.
7265 Specifically, this study used a multiple before after control impact (MBACI) design to compare
7266 unharvested reference sites to sites harvested under the western Washington Forest Practices for
7267 non-fish-bearing streams to assess the effects of these rules on riparian vegetation and wood
7268 recruitment, canopy closure and stream temperature, stream discharge and downstream transport
7269 of suspended sediment and nitrogen, and benthic macroinvertebrates. The Forest Practices rules
7270 for non-fish-bearing streams in the study area includes clearcut harvest with a two-sided 50-foot-
7271 wide riparian buffer along at least 50% of the riparian management zone, including buffers
7272 prescribed for sensitive sites and unstable slopes. Due to the additional unstable slope buffers,
7273 total buffer area was 18 to 163% greater than the current minimum buffer rule. Ten study sites
7274 were chosen with first-, second-, and third-order non-fish-bearing streams. Data was collected
7275 for 1-2 years of pre-harvest, during the harvest period (2012 – 2014), and at least 2 years post-
7276 harvest at all sites. ~~Because of unstable slopes, total buffer area was 18 to 163% greater than the~~
7277 ~~50-foot buffer.~~

TRT2



Basin Size (ha [ac])	10 (25)	Stream Length (m [ft])	591 (1939)
Bankfull Width (cm)	114	Buffers	
Wetted Width (cm)	52	Length Buffered (%)	54
Stream Slope (%)	12	Minimum Area (ha [ac])	1.00 (2.47)
Valley Wall Slope (%)	58	Actual Area (ha [ac])	1.19 (2.93)
Wetted Channel (%)	65	Increase in Area (%)	18
Average Precip (cm [in])	262 (103)	Mean Buffer Width (m [ft])	15 (48)

Commented [WB26]: For the author: Green lines are the riparian stand and LWD transects where data was collected at each of the treatment sites (this example is treatment site 2). Some of those transects were in buffers that were exactly 50ft wide (FP buffers), some were less than 50ft (e.g. near T2), and some were unbuffered (e.g. near T3). This is why data was divided up into buffer types and not analyzed at the site level. Treatment sites, like the Hard Rock study, were a full clearcut harvest of a Type N watershed with the current forest practice rules buffers applied to the entire Np network. See Appendix B for more information.

Commented [WB27R26]: -Authors can delete comment and map after reading-

7278
 7279 ~~This resulted in~~ The riparian stand and wood recruitment data was summarized by 4 different
 7280 buffer types 1) FP Buffers, encompassing the full width (50 feet), 2) <50ft buffers, 3)

7281 Unbuffered, harvested to the edge of the channel, and 4) Reference sites in unharvested ~~forests~~
7282 ~~basins~~. ~~Because of the separation into multiple treatments, sample sizes became small and~~
7283 ~~unbalanced. Thus, no formal~~ statistical analyses were conducted, ~~and~~ only descriptive statistics
7284 were applied for changes in stand structure and wood loading due to the low replication of
7285 reference sites and differences in harvest timing and distribution of buffer types at the treatment
7286 sites. Density decreased by 33 and 51% and basal area by 26 and 49% in the full and <50ft
7287 buffers, respectively, with high variability among sites. Nearly all trees were removed from
7288 Unbuffered ~~sites-areas~~ during harvest (>99% of basal area). ~~In the reference plots/sites,~~
7289 cumulative post-harvest mortality during the 3-year post-harvest interval was only 6.5% of live
7290 density. In contrast, mean post-harvest mortality ~~in of the full-FP buffers sites~~ and the <50 ft
7291 buffers ~~_sites~~ were 31 and 25% of density, respectively. However, there was considerable
7292 variation in mortality among sites exceeding 65% in the FP buffers at two of the two full buffer
7293 treatment sites. Windthrow and physical damage from falling trees accounted for ~75% of
7294 mortality in the full-FP and <50 ft buffers. In contrast to the ~~treated-treatment sites~~, <10% of
7295 trees died due to wind or physical damage in the reference sites. There was little post-harvest
7296 large wood input in reference sites: an average of 4.3 pieces and 0.34 m³ of combined in- and
7297 over-channel volume per 100 m of channel. In contrast, the full buffer sites FP and <50 ft buffers
7298 ~~sites~~ received an average of 23 and 10 pieces/100 m and 2.3 and 0.7 m³/100 m of large wood,
7299 respectively. The majority of recruited large wood pieces had stems with roots attached
7300 (SWRW); 60, 70, and 100% in the reference, full buffer FP, and <50 ft buffer types, respectively.
7301 Pre-harvest channel large wood loading ranged from 55.8 to 111 pieces/100 m and from 9.8 to
7302 25.2 m³/100 m among buffer types. Piece counts remained stable in the reference sites through
7303 year 3 post-harvest, increased in the full buffer FP buffers and unbuffered sites-areas (8 and 13%,
7304 respectively), and decreased in the <50 ft buffers (15%). The authors noted confidence in
7305 interpretation of the descriptive statistics due to the large differences between the post-harvest
7306 treatments and the reference sites. Also, these results were of similar direction and magnitude
7307 that was seen in the preceding Hard Rock and BCIF studies.

7308 For effects of treatment on shade, data was analyzed with generalized linear mixed-effects
7309 models. For effects of treatment on stream temperature, data was analyzed for the seven-day
7310 average in a linear-mixed-effects model analysis of variance. Mean canopy closure decreased in
7311 the treatment sites from 97% in the pre-harvest period to 75%, 68%, and 69% in the first, second,
7312 and third post-harvest years, respectively, and was related to the proportion of stream buffered
7313 and to post-harvest windthrow within the buffer. The seven-day average temperature response
7314 increased by 0.6°C, 0.6°C, and 0.3°C in the first, second, and third post-harvest years,
7315 respectively. During and after harvest, mean monthly water temperatures were higher, but
7316 equaled or exceeded 16.0°C only in 2 treatment sites by up to 1.8°C at one site and by 0.1°C at
7317 another. None of the three REF sites exceeded 16.5°C during the study. ~~Predictive models could~~
7318 ~~not be fitted to the temperature data for statistical analysis.~~

7319 Results for changes in nutrient concentrations post-harvest were highly variable. Harvest
7320 treatment effects on nutrient concentrations, discharge, and suspended sediment export could not
7321 be calculated because discharge prediction equations could not be developed.

7322

7323

7324 McIntyre et al., 2018

7325

7326 McIntyre, A.P., M.P. Hayes, W.J. Ehinger, S.M. Estrella, D. Schuett-Hames, and T. Quinn
7327 (technical coordinators). 2018. Effectiveness of Experimental Riparian Buffers on Perennial
7328 Non-fish-bearing Streams on Competent Lithologies in Western Washington. Cooperative
7329 Monitoring, Evaluation and Research Report CMER 18-100, Washington State Forest Practices
7330 Adaptive Management Program, Washington Department of Natural Resources, Olympia, WA.

7331

7332 The purpose of the study was to evaluate the effectiveness of forest management prescriptions in
7333 maintaining aquatic conditions and processes for small non-fish-bearing (Type N) headwater
7334 stream basins underlain by competent “hard rock” lithologies (i.e., volcanic or igneous rock) in
7335 western Washington. Specifically, this study quantified and compared the effects of timber
7336 harvest adjacent to Type N streams on riparian stand structure and tree mortality, in stream wood
7337 loading and recruitment, stream temperature and canopy cover, stream discharge, turbidity, and
7338 suspended sediment export, nitrogen export, and response of stream-associated amphibians. This
7339 study used a before-after control-impact (BACI) study design. This involved evaluation of four
7340 experimental treatments: (1) unharvested reference (n = 6); (2) 100% treatment (n = 4), a
7341 [minimum](#) two-sided 50-ft riparian buffer along the entire Riparian Management Zone (RMZ);
7342 (3) FP treatment (n = 3), a [minimum](#) two-sided 50-ft riparian buffer along at least 50% of the
7343 RMZ, consistent with the current Forest Practices buffer prescription for Type N streams [and](#)
7344 [including This treatment also included a circular buffer](#) protecting the uppermost points of
7345 perennial flow (PIP), [headwall and side-slope seeps, and tributary junctions](#); [and](#) (4) 0%
7346 treatment (n = 4), clearcut to stream edge (no [riparian](#) buffer). [Study sites comprised the entire](#)
7347 [non-fish-bearing \(Type N\) basin when possible and The](#) upland forests of all treatments were
7348 clearcut harvested. The study design included [data collection for](#) at least two years [of](#) pre-harvest
7349 (2006 –2008); [and up to](#) three years of post-harvest data [collection](#) (2009 – 2011). Results for
7350 stand structure and tree mortality showed that in the RMZs, the proportional changes in stem
7351 count (dstems) and basal area (dBA) were similar for the reference (mean dstems: -11.8, SE 5.3;
7352 dBA: -6.9, SE 5.4) and 100% (mean dstems: -3.8, SE 5.9; dBA -6.7, SE 6.0) treatment. In
7353 contrast, the magnitude of decrease was significantly greater in the FPB (portion of FP
7354 containing trees; mean dstems: -29.6, SE 6.5; dBA 124.4, SE 6.7) treatment than in either the
7355 reference or 100% treatment. The pattern was similar in the PIPs. 2 years post-harvest tree
7356 mortality was mostly (70%) attributed to wind/mechanical agents (pre-harvest wind/mechanical
7357 agent caused mortality was 70%). In the reference sites, trees that died post-harvest had smaller
7358 diameters (mean 10.3 in) and fewer came from the overstory crown class (59.0%) than the other
7359 treatments. In contrast, in the 100% and FPB treatments, ~70% of trees that died were from the
7360 overstory crown class and their mean diameters were 1 (11.2 in) and 2 (12.2 in) in greater than
7361 those in the reference sites, respectively. Results for wood recruitment and loading showed that

7362 tree fall rates were highly variable during the pre-harvest period between sites ranging from 0 to
7363 239.9 trees/ha/yr. Large wood (LW) recruitment rates in the pre-harvest period were also highly
7364 variable ranging from 0 to 121.6 pieces/ha/yr, along with recruitment volume (0-16.2 m³/ha/yr).
7365 2 years post-harvest recruitment rates in the reference riparian management zones (RMZs) were
7366 lower and less variable (5.9 to 37.3 trees/ha/yr) than in buffer treatments. Tree fall rates for the
7367 100% treatment ranged from 7.7 to 76.4 trees/ha/yr, and for the FPB treatments tree fall rates
7368 ranged from 4.2 to 152.2 trees/ha/yr. Post-harvest LW recruitment volumes in reference RMZs
7369 were relatively low, ranging from 0.7 to 2.2 m³/ha/yr. Post-harvest LW recruitment volumes
7370 were generally higher and more variable in the 100% and FPB RMZs, ranging from 0.3 to 14.0
7371 m³/ha/yr in the 100% treatment and 0 to 7.6 m³/ha/yr in the FPB. Because of the high variability
7372 between sites in all treatments the p values for comparisons between treatments were generally
7373 high ($p \geq 0.35$), except for the FPB vs. reference comparison for piece count which was nearly
7374 significant ($p = 0.13$). The only significant differences were for the 0% treatments which had
7375 significantly lower LW recruitment by volume than the Reference RMZ ($P = 0.02$). For PIPs, LW
7376 recruitment in the 100% treatment was over 12 times the reference rate by piece count ($P = 0.03$)
7377 and 30 times the reference rate by volume ($P = 0.04$). Recruitment in the FPB PIPs was also
7378 high, over nine times the reference rate by piece count ($P = 0.08$) and 18 times the reference rate
7379 by volume ($P = 0.11$). The amount of change in the number of LW pieces per meter from pre-
7380 harvest to post-harvest depended on treatment ($P < 0.01$). Analysis estimated the changes in
7381 100%, FP and 0% treatments to be different from the change in the reference ($P < 0.001$, 0.03 and
7382 < 0.01 , respectively). The percentage of the stream channel length covered by newly recruited
7383 wood in the second post-harvest year ranged from 0 to 11% in the reference, 1 to 15% in the
7384 100% treatment and 0 to 10% in the FP treatment and was 0% in all four of the 0% treatments.
7385 The percent of stream channel covered by new wood differed between the 0% treatment and the
7386 reference ($P = 0.03$), 100% ($P < 0.01$), and FP treatments ($P = 0.03$). Overall, the authors
7387 estimated a mean between-treatment increase of 60% (95% CI: 0–150%), 70% (95% CI:
7388 0–190%) and 170% (95% CI: 80–330%) in the number of SW pieces per stream meter in the
7389 100%, FP and 0% treatments compared with the reference, respectively. Also, a between-
7390 treatment increase of 60% (95% CI: 30–110%), 40% (95% CI: 0–100%) and 50% (95% CI:
7391 10–90%) in the number of LW pieces per stream meter in the 100%, FP and 0% treatments
7392 compared with the reference, respectively. The authors conclude that windthrow was responsible
7393 for much of the increase in LW. However, they also posit that the timing and magnitude of wood
7394 inputs was inconsistent, resulting in considerable variability between and within sites, especially
7395 in the FP treatment. Results for shade response to treatments post-harvest was greatest in the 0%
7396 treatment than in either the 100% or the FP treatment. Effective shade decreased to 77, 52, and
7397 14% 2 years post-treatment, in the 100%, FP, and 0% buffer treatments, respectively. Canopy and
7398 Topographic Density (CTD), defined as the percentage of the photograph obscured by vegetation
7399 or topography decreased from an average of 95% pre-harvest to 86, 71, and 43% 2 years post-
7400 harvest in the 100%, FP, and 0% buffer treatments, respectively. All were significantly lower
7401 than the reference (92% 2 years post-treatment). Results for stream temperature showed
7402 maximum daily water temperatures increased post-harvest in all but one of the harvested sites
7403 and was elevated over much of the year at most of the sites. Daily temperature response (TR)
7404 increased in late winter or early spring, reached a maximum in July–August and was still

7405 elevated well into the fall. This pattern was observed at most of the sites. For the Buffer
7406 Treatment locations, 94 of the 131 calculated mean monthly temperature responses (MMTRs)
7407 were significant and 91 of these significant responses were positive. In comparison, only 52 of
7408 156 MMTR values calculated for the reference sites were significant and these were nearly
7409 evenly split with 25 positive and 27 negative responses. This strongly suggests that the pattern of
7410 post-harvest increases in daily maximum water temperature is real even though the magnitude of
7411 some of the individual MMTRs is relatively small ($<0.5^{\circ}\text{C}$). Warming tended to be greatest in
7412 July or August with MMTR ranging from 0.5°C to 2.3°C in the 100%, -0.4°C to 1.8°C in the FP,
7413 and 1.0°C to 3.5°C in the 0% treatments. Post-harvest, Max7D (seven-day-average maximum
7414 stream temperature) was higher at 36 of the 40 locations within the harvest units across all 11
7415 buffer treatment sites regardless of presence or absence of a buffer, buffer width, and
7416 longitudinal location along the stream. Relative to the unharvested sites, there were summertime
7417 temperature increases throughout the stream length and across all buffer treatment sites. The
7418 authors conclude that none of the buffer treatments were successful in preventing significant
7419 increases in maximum stream temperature. The generalizable conclusions made by the authors
7420 from this portion of the study are that 1) Buffer widths greater than 50 ft (15.2 m) are needed to
7421 prevent shade loss and (2) Maximum water temperature decreased below the harvest unit after
7422 flowing through approximately 100 m of intact forest but was still elevated compared to pre-
7423 harvest conditions. Results for nitrogen and phosphorus concentrations showed that post-harvest
7424 changes for total-N or total-P were not significant for any of the treatments relative to the
7425 Reference. The only significant difference detected within 2 years post-harvest was for nitrate-N
7426 concentration between the 0% buffer treatment and all other treatments. However, for annual
7427 export, total-N and nitrate-N export increased post-harvest at all sites, with the smallest increase
7428 in the 100% treatment and the largest in the 0% treatment. Compared to the reference sites, the
7429 GLMM analysis showed a relative increase in total-N export post-harvest of 5.52 ($P = 0.051$),
7430 11.52 ($P = 0.0007$), and 17.16 ($P < 0.0001$) $\text{kg ha}^{-1} \text{ yr}^{-1}$ in the 100%, FP, and 0% treatments. The
7431 GLMM analysis showed a relative increase in nitrate-N export post-harvest of 4.83 ($P = 0.048$),
7432 10.24 ($P = 0.001$), and 15.35 ($P < 0.0001$) $\text{kg ha}^{-1} \text{ yr}^{-1}$ in the 100%, FP, and 0% treatments,
7433 respectively, only slightly less than the changes in total-N. Total-P export increased post-harvest
7434 by a similar magnitude in all treatments: 0.10 ($P = 0.006$), 0.13 ($P = 0.001$), and 0.09 ($P = 0.010$)
7435 $\text{kg ha}^{-1} \text{ yr}^{-1}$ in the 100%, FP, and 0% treatments, respectively. The increase in N, total-N and
7436 nitrate-N, from the treatment watersheds post-harvest was strongly correlated with the increase
7437 in annual runoff ($R^2 = 0.970$ and 0.971 ; $P = 0.001$ and 0.001 , respectively) and with the
7438 proportion of the basin harvested ($R^2 = 0.854$ and 0.852 ; $P = 0.031$ and 0.031 , respectively). The
7439 correlation with the proportion of stream length buffered was weaker ($R^2 = 0.761$ and 0.772 ; P
7440 < 0.079 and 0.072 , respectively). In contrast, total-P export was uncorrelated with all three
7441 variables. Overall, the authors concluded that mean flow-weighted concentration of total-N and
7442 nitrate-N increased at all buffer treatment sites post-harvest, however the magnitude was variable
7443 and significant only for the 0% buffer treatment. However, the export of total-N increased in the FP and
7444 0% treatments and nitrate-N increased in all buffer treatments. Increases in N export was
7445 correlated with increased stream discharge and the proportion of the site that was harvested. Pre-
7446 harvest total-P concentration was low and remained so post-harvest, although P export increased
7447 slightly post-harvest in all treatments due to the increase in discharge. Results for changes in

7448 water turbidity and suspended sediment concentrations (SSC) showed both turbidity and SSC
7449 increased with increasing discharge during storm events but then rapidly fell off. Analysis of
7450 treatment effects revealed no significant effects of harvest and no clear pattern regarding the
7451 relative effectiveness of buffer treatments at mitigating the effects of clearcut harvests on
7452 suspended sediment export (SSE). The general conclusions made by the authors were that all
7453 sites appeared to be supply limited both pre- and post-harvest. Results for litterfall input showed
7454 a decrease in TOTAL litterfall input in the FP (P = 0.0034) and 0% (P = 0.0001) treatments
7455 between pre- and post-treatment periods. LEAF litterfall (deciduous and conifer leaves
7456 combined) input decreased in the FP (P = 0.0114) and 0% (P <0.0001) treatments in the post-
7457 treatment period. In addition, CONIF (conifer needles and scales) litterfall input decreased in the
7458 FP (P = 0.0437) and 0% (P <0.0001) treatments, DECID (deciduous leaves) in the 0% (P
7459 <0.0001) treatment, WOOD (twigs and cones) in the FP (P = 0.0044) and 0% (P = 0.0153)
7460 treatments, and MISC (e.g., moss and flowers) in the 0% (P = 0.0422) treatment. Results for
7461 comparison of the post-harvest effects between treatments showed LEAF litterfall input
7462 decreased in the 0% treatment relative to the reference (P = 0.0040), 100% (P = 0.0008), and FP
7463 (P = 0.0267) treatments. Likewise, there was a decrease in DECID litterfall input in the 0%
7464 treatment relative to the Reference (P = 0.0001), 100% (P <0.0001), and FP (P = 0.0015)
7465 treatments. Results for detritus with comparisons between the pre- and post-treatment periods
7466 showed an increase in TOTAL detritus export in the 100% treatment (P = 0.0051) and a decrease
7467 in the 0% treatment (P = 0.0046; Table 12-9). Likewise, there was an increase in CPOM,
7468 WOOD, MISC, and FPOM detritus export in the 100% treatment (P <0.05), but a decrease in the
7469 0% treatment (P <0.05). The authors for this portion of the study conclude that overall, total
7470 litterfall input was slightly higher after harvest in the 100% treatment, lower in the FP treatment
7471 and lowest in the 0% treatment; however, statistical differences were only detected for
7472 deciduous inputs between the 0% treatment and the other treatments. Total detritus export
7473 decreased in the 0% treatment relative to the reference, and in the FP and 0% treatments relative
7474 to the 100% treatment.

7475

7476

7477 McIntyre et al., 2021

7478

7479 McIntyre, A.P., M.P. Hayes, W.J. Ehinger, S.M. Estrella, D.E. Schuett-Hames, R. Ojala-Barbour,
7480 G. Stewart and T. Quinn (technical coordinators). 2021. Effectiveness of experimental riparian
7481 buffers on perennial non-fish-bearing streams on competent lithologies in western Washington –
7482 Phase 2 (9 years after harvest). Cooperative Monitoring, Evaluation and Research Report CMER
7483 2021.07.27, Washington State Forest Practices Adaptive Management Program, Washington
7484 Department of Natural Resources, Olympia, WA.

7485

7486 This study was a follow-up study to the hard-rock Phase 1 study (McIntyre et al., 2018) to assess
7487 changes over longer time periods (up to 9 years post-harvest). The purpose of the study was to
7488 evaluate the effectiveness of forest management prescriptions in maintaining aquatic conditions
7489 and processes for small non-fish-bearing (Type N) headwater stream basins underlain by
7490 competent “hard rock” lithologies (i.e., volcanic or igneous rock) in western Washington.
7491 Specifically, this study quantified and compared the effects of timber harvest adjacent to Type N
7492 streams on riparian stand structure and tree mortality, in stream wood loading and recruitment,
7493 stream temperature and canopy cover, stream discharge, turbidity, and suspended sediment
7494 export, nitrogen export, and response of stream-associated amphibians. This study used a
7495 before-after control-impact (BACI) study design. This involved evaluation of four experimental
7496 treatments: (1) unharvested reference (n = 6), (2) 100% treatment (n = 4), a two-sided 50-ft
7497 riparian buffer along the entire Riparian Management Zone (RMZ), (2) FP treatment (n = 3), a
7498 two-sided 50-ft riparian buffer along at least 50% of the RMZ, consistent with the current Forest
7499 Practices buffer prescription for Type N streams, (3) 0% treatment (n = 4), clearcut to stream
7500 edge (no-buffer). The upland forests of all treatments were clearcut harvested. The study design
7501 included data collection for at least two years pre-harvest (2006–2008), and up to nine years
7502 post-harvest from 2009 (harvest began in 2008) until 2016 or 2017 depending on the variable
7503 (e.g., wood loading, shade, etc.). Results for stand structure showed that in the buffered portions
7504 of the FP treatments (FPB) density, basal area and relative density (RD) decreased by 59%, 55%
7505 and 54%, respectively, 8 years after harvest. For the same variables, reductions in the 100%
7506 RMZs were 30%, 14%, and 17%, respectively. In contrast, stand structure in the reference RMZs
7507 was more stable, with a 17% decrease in density and little change in basal area or RD. Change in
7508 live basal area did not differ statistically between 100% and REF RMZs for any time interval
7509 although the differences increased over time. The FPB–REF contrast was not significant in the
7510 first interval (years 1 and 2 post-harvest), but it was in subsequent intervals (5- and 8-years post-
7511 harvest) as the magnitude of change in FPB RMZs increased over time. The FPB–100% contrast
7512 was not significant until the last interval when basal area stabilized in the 100% treatment but
7513 continued to decline in FPB. Between treatment comparison of cumulative change in live basal
7514 area (m²/ha) between the 100% treatment and the Reference was -2.9 (CI: -16.9, 11.0), -6.0 (CI:
7515 -20.0, 8.0), and -6.8 (CI -20.8, 7.1) for the first-, second-, and third-time intervals respectively
7516 (none were significant). Comparison between the FPB and Reference were -10.2 (CI: -25.5, 5.2),
7517 -16.1 (CI: -31.4, -0.8), and -21.1 (CI: -36.4, -5.8) for the first-, second-, and third-time intervals
7518 respectively (differences for intervals 2 and 3 were significant). For tree mortality, results
7519 showed that by year 8 post-harvest mortality as a percentage of pre-harvest basal area was lower
7520 in the reference (16.1%) than in the 100% (24.3%) and FPB (50.8%). The FPB–Reference
7521 contrast was not significant 2 years post-harvest, but it was at 5- and 8-years post-harvest as
7522 mortality in FPB increased relative to the reference. The contrast between the 100% and Ref
7523 were not significant for any time interval 8 years post-harvest. The contrasts 100% vs. REF and
7524 FPB vs. 100%—were not significant for any time interval. This may have been because of the
7525 high variability in the data. There was a temporal pattern to mortality in 100% and FPB RMZs.
7526 Annual rates of mortality as percentage of live basal area and density were highest in the first
7527 two years after harvest, then decreased. Wind/physical damage was the primary cause of
7528 mortality. In the 100% treatment it accounted for 78% and 90% of the loss of basal area and

7529 density, respectively; in FPB it accounted for 78% and 65% of the loss. Wind accounted for a
7530 smaller proportion of mortality in reference RMZ (52%). Large wood recruitment to the channel
7531 was greater in the 100% and FPB RMZs than in the reference for each pre- to post-harvest time
7532 interval. Eight years post-harvest mean recruitment of large wood volume was two to nearly
7533 three times greater in 100% and FPB RMZs than in the references. Large wood recruitment rates
7534 were greatest during the first two years, then decreased. However, these differences were not
7535 significant between any treatment comparisons, again, likely due to the high variability in the
7536 data. Mean large wood loading differed significantly between treatments in the magnitude of
7537 change overtime. Results showed a 66% ($P < 0.001$), 44% ($P = 0.05$) and 47% ($P = 0.01$) increase
7538 in mean large wood density in the 100%, FP and 0% treatments, respectively, in the first 2 years
7539 post-harvest compared with the pre-harvest period and after controlling for temporal changes in
7540 the references. Five years post-treatment the mean LW density in the FP continued to increase
7541 42% ($P = 0.08$), and again 8 years post-treatment (41%; $P = 0.09$). Results for canopy cover
7542 showed that riparian cover declined after harvest in all buffer treatments reaching a minimum
7543 around 4 years post-harvest. The treatments, ranked from least to most change, were REF, 100%,
7544 FP, and 0% for all metrics and across all years. Effective shade results showed decreases of 11,
7545 36, and 74 percent in the 100%, FP, and 0% treatments, respectively. Significant post-harvest
7546 decreases were noted for all treatments and all years. Results for stream temperature showed that
7547 within treatment mean post-pre-harvest difference in the REF treatment never exceeded 1.0°C.
7548 In contrast, the mean within treatment difference in the 100% treatment was 2.4°C in 2009 (Post-
7549 harvest year 1) but never exceeded 1.0°C in later years. The mean difference in the FP treatment
7550 exceeded 1.0°C immediately after harvest then again in 2014–2016 (post-harvest years 6–9)
7551 while in the 0% treatment the mean difference was 5.3°C initially, then decreased over time to
7552 near, but never below, 0.9°C. Stream temperature increased post-harvest at most locations within
7553 all 12 harvested sites and remained elevated in the FP and 0% treatments over much of the nine
7554 years post-harvest. Temperature responses varied by treatment, by season, and over the years. In
7555 three out of the first four post-harvest years there was, at least, a weak ($r < -0.48$) negative
7556 correlation between July monthly mean temperature response (MMTR) and the change in
7557 riparian cover based on each of the four shade metrics. The correlation was generally weaker
7558 ($-0.4 < r$ and $P > 0.10$) after post-harvest year 4, except for post-harvest year 9 ($-0.6 < r < -0.4$).
7559 However, there were only eight data pairs available for Post 9, compared to ten to twelve for the
7560 other years, which affected the correlation coefficient and p-value. However, there was a great
7561 deal of variability in the correlation coefficient of July MMTR with shade across post-harvest
7562 years among sites and treatments with some sites showing negative correlations and others
7563 positive for some treatments in some years. Considering site characteristics, aspect showed an
7564 influence on stream temperature response. In the first five post-harvest years and in Post 7 the
7565 highest MMTR in each treatment was nearly always the site with a southern (SE or SW) aspect.
7566 No significant correlation between July MMTR and either mean July discharge or the post-
7567 harvest difference in discharge was observed. For the effects of harvest on stream discharge,
7568 cumulative results of regression analysis (forward and reverse regression approaches) indicated
7569 that discharge did increase following harvest. In relative terms, discharge increased by 5-7% on
7570 average in the 100% treatments while increasing between 26-66% in the FP and 0% treatments.
7571 The change in discharge following harvest was also affected by climate, weather, and physical

7572 hydrology of the watershed. In all basins, discharge varied with precipitation, but this was a
7573 complex relationship showing lag time between precipitation events and discharge rate response
7574 in some watersheds. This indicated a potential relationship with physical hydrology at some
7575 watersheds. Results for water turbidity and suspended sediment export (SSE) were stochastic in
7576 nature and the relationships between SSE export and treatment effects were not strong enough to
7577 confidently draw conclusions. Results for harvest effects on total nitrogen export following a
7578 generalized linear mixed effects model, however, showed significant ($P < 0.05$) treatment effects
7579 were present in the FP treatment post-harvest and in the 0% treatment in the post-harvest (2-
7580 years immediately following harvest) and extended periods (2015 – 2017; 7 and 8 years post-
7581 harvest) relative to the reference sites, but there were no significant differences in total-N export
7582 between the treatments. Analysis showed an increase in total-N export of 5.73 ($P = 0.121$), 10.85
7583 ($P = 0.006$), and 15.94 ($P = 0.000$) kg/ha/yr post-harvest in the 100%, FP, and 0% treatments,
7584 respectively, and of 6.20 ($P = 0.095$), 5.34 ($P = 0.147$), and 8.49 ($P = 0.026$) kg/ha/yr in the
7585 extended period. Results for nitrate-N export showed changes similar to but slightly less than
7586 those seen in the total-N analysis with a relative increase in nitrate-N export of 4.79 ($P = 0.123$),
7587 9.63 ($P = 0.004$), and 14.41 ($P < 0.001$) kg/ha/yr post-harvest in the 100%, FP, and 0% treatments,
7588 respectively. None of the changes in the extended period were significant. However, the authors
7589 note that there was high variability in the data for the extended period and nitrate-N export only
7590 returned to pre-harvest levels in one watershed. The increase in total-N and nitrate-N export
7591 tended to be highest during the high flow months in the fall and early winter. The authors
7592 conclude that the 100% treatment was generally the most effective in minimizing changes from
7593 pre-harvest conditions, the FP was intermediate, and the 0% treatment was least effective. The
7594 collective effects of timber harvest were most apparent in the 0% treatment in the two years
7595 immediately post-harvest.

7596

7597 **LW**

7598

7599 Johnston et al., 2011

7600

7601 Johnston, N. T., Bird, S. A., Hogan, D. L., & MacIsaac, E. A. (2011). Mechanisms and source
7602 distances for the input of large woody debris to forested streams in British Columbia, Canada.
7603 Canadian journal of forest research, 41(11), 2231-2246. <https://doi.org/10.1139/x11-110>

7604

7605 The purpose of this study was to determine whether the processes and source distances from
7606 which LW entered streams differed among channel types and sizes, to describe LW source
7607 distance curves for a wide range of undisturbed stream and forest types, and to characterize the
7608 relationships between LW input mechanism, source distance, and piece size. Input processes,
7609 source distances, and physical characteristics of approximately 2100 pieces of LW at 51
7610 anthropogenically undisturbed stream reaches throughout south and central British Columbia

7611 were determined. Large wood (LW) was defined in this study as pieces within or suspended
7612 above the active channel, with a minimum length of 1 m. and capable of inducing sediment scour
7613 or deposition. A delivery mechanism was assigned to each LW piece, when it could be
7614 determined, as bank erosion, landslide, windthrow of live trees, stem snap, or standing dead tree
7615 fall. Differences in the frequencies of count data among LW delivery mechanisms, LW positions,
7616 or LWD functions were assessed using chi-square tests. The effects of channel (type, width) and
7617 forest (maximum tree height) characteristics on the proportions of LWD pieces entering the
7618 channel by a given input mechanism were examined using ANCOVA. Channel type for this
7619 study was grouped into 3 categories; riffle-pool (RP), cascade-pool (CP), and step-pool (SP).
7620 Results showed that tree mortality was the most common entry mechanism at all channel types
7621 and width categories and accounted for 65% of all LW pieces sampled. Both channel and
7622 riparian forest characteristics influenced the proportion of LW pieces that entered streams by tree
7623 mortality ($P < 0.05$) but did not vary significantly among channel types ($P = 0.13$). The
7624 proportion of LW pieces recruited by tree mortality decreased with increasing channel width and
7625 with increasing maximum tree height. Bank erosion inputs accounted for 20%–25% of all LW
7626 pieces at the lower-gradient RP and CP sites but were much less important at the SP channels.
7627 Erosion inputs increased with increasing stream size within all channel types ($P = 0.0004$). Wind-
7628 induced inputs (windthrow and stem snap) accounted for 13%–20% of inputs over the channel
7629 types and generally increased in importance in the smaller channels. The proportion of LW
7630 recruited to the stream by stem breakage increased with increasing tree height ($P < 0.0001$) and
7631 varied among channel types ($P = 0.040$), being about twice as prevalent at SP channels as
7632 elsewhere. Landslide inputs of LWD were a minor delivery mechanism. There was considerable
7633 variability in distances from which LW entered the stream. However, based on the cumulative
7634 distributions over sites, 90% of the LW pieces or volume entering the channels originated within
7635 18 m of the stream in 90% of all cases (between 2 and 23 m in all cases). The distances from
7636 which LW entered the streams differed significantly among the various input mechanisms ($P <$
7637 0.001), the rank ordering of the mean source distances being bank erosion $<$ tree mortality $<$
7638 stem breakage $<$ windthrow $<$ landslides. Bank erosion and landslides delivered the largest LW
7639 pieces and tree mortality and stem breakage the smallest. In general, source distances increased
7640 with increasing tree height, with the effect being stronger in the steeper channel types and
7641 weaker in the wider channels for LW pieces and volume. However, all two-way interactions
7642 among variables were significant implying that the mechanisms through which vegetation and
7643 stream geomorphology influenced LW source distance were complex. Maximum tree height in
7644 the adjacent forest accounted for the greatest variance in in-stream LW source distance for all
7645 models.

7646

7647 **Nutrient**

7648

7649 Deval et al., 2021

7650

7651 Deval, C., Brooks, E. S., Gravelle, J. A., Link, T. E., Dobre, M., & Elliot, W. J. (2021). Long-
7652 term response in nutrient load from commercial forest management operations in a mountainous
7653 watershed. *Forest Ecology and Management*, 494, 119312.
7654 <https://doi.org/10.1016/j.geomorph.2013.11.028>

7655

7656 The purpose of this study was to quantify and compare the differences in nitrogen and
7657 phosphorus concentrations and loads between pre-disturbance, post road construction (post-
7658 road), post experimental harvest (PH-I), and post operational harvest (PH-II) from both a
7659 hydrological yield and nutrient concentration perspective. This study was carried out in the Mica
7660 Creek Experimental Watershed in Northern Idaho. For this analysis time periods have been
7661 broken into four distinct phases: Pre-disturbance (1992–1997), Post-road (1997–2001),
7662 experimental-harvest Phase I (PH-I) (2001–2007), and operational sequential harvest Phase II
7663 (PH-II) when the extent and frequency of harvests increased (2007–2016). PH-I represents an
7664 experimental treatment phase during which harvest activities were experimentally controlled
7665 (only upstream headwater watersheds were harvested and mature vegetation removal ranged
7666 between 24% and 47%) followed by site management operations including broadcast burning
7667 and replanting. PH-II represents the post-experimental phase where the study area transitioned to
7668 operational treatments that consisted of additional road construction and timber harvest, with site
7669 management operations including pile burning and competition release herbicide application.
7670 During this operational phase, the mature vegetation removal in the upstream and cumulative
7671 downstream watersheds ranged between 36% and 50% and 17–28%, respectively. Monthly
7672 annual grab samples of stream water were collected from seven flumes over the course of 25
7673 years (from pre- to post-treatments). The samples were analyzed for six parameters, specifically
7674 nitrate + nitrite (NO₃ + NO₂), total Kjeldhal nitrogen (TKN), total ammonia nitrogen (TAN)
7675 containing un-ionized (NH₃) and ionized (NH₄⁺) ammonia, total nitrogen (TN), total
7676 phosphorus (TP), and orthophosphate (OP). This study used a before-after, control-impact paired
7677 series design (BACIPS) to evaluate direct and cumulative effects of forest management practices
7678 on stream nutrient concentrations in paired and nested watersheds. Results for long-term trends
7679 in stream flow showed a statistically significant increasing trend in all the watersheds during the
7680 fall and winter seasons. Significant increases in summer streamflow only occurred in the control
7681 watersheds. There were minimal changes in TKN concentration with a slight observed reduction
7682 in long-term TKN loads. Overall, the cumulative mean TAN loads from all watersheds did not
7683 show large variations with sequential varying treatments over time. In contrast to TAN, there was
7684 a significant response in NO₃ + NO₂ following timber harvest. The response in NO₃ + NO₂
7685 concentrations was negligible at all treatment sites following the road construction activities.
7686 However, NO₃ + NO₂ concentrations during the PH-I period increased significantly ($p < 0.001$)
7687 at all treatment sites. Similar to the PH-I period, all watersheds experienced significant increases
7688 in NO₃ + NO₂ concentration during the PH-II treatment period. Overall, the cumulative mean
7689 NO₃ + NO₂ load from all watersheds followed an increasing trend with initial signs of recovery
7690 in one treatment watershed after 2014. Mean monthly TP concentrations showed no significant
7691 changes in the concentrations during the post-road and PH-I treatment periods. However, a
7692 statistically significant increase in TP concentrations ($p < 0.001$) occurred at all sites, including

7693 the downstream cumulative sites, during PH-II. Generally, OP concentrations throughout the
7694 study remained near the minimum detectable concentrations. A statistically significant increase
7695 in mean monthly OP concentrations occurred only at the cumulative downstream treatment site
7696 during both Post-road (p-value = 0.021) and PH-I (p-value < 0.001) treatment periods,
7697 respectively. The largest cumulative increase in mean annual loads was largely attributed to
7698 increased flow. The authors conclude that only relatively small increases in nutrient loads were
7699 detected suggesting that Idaho Forest Practices Act regulations and BMPs are effective in
7700 minimizing the delivery of particulate-bound pollutants. Forest management activities increased
7701 stream NO₃ + NO₂ concentrations and loads following timber harvest activities, but these effects
7702 were also attenuated in downstream reaches and reduced through time as vegetation regrowth
7703 occurred.

7704

7705 Quinn, T., G.F. Wilhere, and K.L. Krueger, technical editors. 2020. Riparian Ecosystems, Volume
7706 1: Science Synthesis and Management Implications. Habitat Program, Washington Department
7707 of Fish and Wildlife, Olympia.

7708 This publication is a synthesis of scientific literature concerning riparian areas (function, process,
7709 characteristics, etc.) for the purpose of informing management and the development of policies
7710 related to management of riparian areas and watersheds of Washington State. The most relevant
7711 information in the publication to this review are in chapters 3 (large wood), 4 (stream
7712 temperature), and 6 (Nutrient dynamics in Riparian ecosystems).

7713 The main conclusions from chapter 3 (large wood) state that the successful conservation, or
7714 restoration, of fish habitats in forested areas requires management practices that deliver adequate
7715 wood into aquatic systems. They purpose the main scientific uncertainties, from a management
7716 perspective, is (1) the shape of the wood recruitment curves under different watershed and site-
7717 level conditions. These curves describe the function of wood input into streams from greater
7718 distances from the stream (source distance curves) based on stand structure (e.g., young-old, tree
7719 height variability, density metrics, etc.), species compositions (especially conifer vs. hardwood),
7720 and site conditions (e.g., slope, moisture availability, soils, site index). The second uncertainty is
7721 the effects of the potential wood delivery mechanisms that occur outside of the riparian area
7722 (e.g., landslides, debris flows). They posit that management objectives for large wood
7723 recruitment potential should aim to restore site composition and structure that is similar to
7724 unmanaged riparian forests. The authors suggest that much is known about large wood
7725 recruitment potential from within the riparian forests based on site potential and stand structure.
7726 Previous work and the development of source distance curve equations show the range of
7727 “effective” tree heights (trees with the bulk of stem > 10 cm, functionally classified as large
7728 wood) is between 85 and 230 feet. This means 100% of a sites wood recruitment potential is
7729 within 85 -230 feet of the stream. However, these equations do not account for the presence of
7730 smaller trees or the potential of tree recruitment from outside of the riparian area, or from
7731 extreme channel migration events.

7732 The main conclusions from chapter 4 identify that the science surrounding stream thermal
7733 regimes is uneven. Scientists are certain that stream temperatures and thermal regimes are
7734 important to aquatic species, and thus it is important for management practices to restore and
7735 conserve these conditions. However, while the general conclusions of most studies show that
7736 land use changes (urbanization, and agriculture) and forest management within riparian areas
7737 leads to warmer stream temperatures, the spatial and temporal effects of any specific riparian
7738 management action remain uncertain. Recovery rates for stream temperature post-treatment vary
7739 greatly based on site location because of differences in stream size (width and depth), and
7740 physiography (climate, physical geography). Shade from the adjacent riparian area is widely
7741 accepted as the most important, and most directly manageable, factor affecting stream
7742 temperature. However, because of the variability in other factors affecting stream temperature,
7743 predicting changes in stream temperature from shade removal will likely always suffer from
7744 imprecision.

7745 The main conclusions from chapter 6 (Nutrient dynamics in Riparian ecosystems) list land use,
7746 forest age and composition, Climate and seasonality, elevation and topography, hydrology,
7747 nutrient concentrations, forms and inputs, soil properties and geology, and biota as the major
7748 factors influencing nutrient dynamics in riparian ecosystems. Riparian areas that are structurally
7749 diverse in physiography and soil are most likely to support diverse biota (vegetation, animals,
7750 and microbial communities. More diverse riparian communities are considered best in processing
7751 and assimilating nutrient loads. The authors identify headwater streams as important zones for
7752 active nutrient processing because they affect downstream nutrient loads. Also, maintaining the
7753 connection between the aquatic and terrestrial environments via floodplain conservation and
7754 restoration is important. While there is still a lot of uncertainty involved in the mechanisms
7755 responsible for nutrient transport through the system, it is clear that riparian areas are vital for the
7756 not only providing nutrients to stream, but also in filtering, processing, and storing nutrients in
7757 the short and long term. The results of most studies indicate that the storage and filtering of
7758 nitrogen is most effective in areas with wide vegetated buffers compared to narrower buffers or
7759 unvegetated riparian areas regardless of the type of vegetation present. However, the type of
7760 vegetation directly impacts the quality and quantity of nutrients available. Deciduous trees
7761 generally provide more litter with higher nutrient content. Coniferous trees, on the other hand,
7762 live longer and provide more shade and large wood input potential. Thus, the authors conclude
7763 that riparian management should consider both the structural and food web roles of each species
7764 present in a forested riparian area.

7765