WASHINGTON GEOLOGICAL SURVEY
MAP SERIES 2023-05

Geologic Map of the Kittitas and East Kittitas 7.5-minute Quadrangles, Kittitas County, Washington

Andrew J. Sadowski, Logan R. Wetherell, Megan L. Anderson, and Jack E. Powell

December 2023

Geologic Map of the Kittitas and East Kittitas 7.5-minute Quadrangles

©2023 Washington Geological Survey

www.dnr.wa.gov/geology



ABSTRACT

New geological and geophysical investigations of the Kittitas and East Kittitas quadrangles refine Neogene stratigraphy and characterize geologic structures in southern Kittitas Valley. New whole rock geochronometry (n=329) locally refines the middle Miocene chemostratigraphic framework of the Columbia River Basalt Group (CRBG). These strata are filled toward Kittitas Valley by faulting and folding at the mountain front. Abundant suprabasalt late Miocene to Pliocene volcaniclastic strata are less steeply tilted.

From twelve U-Pb analyses, we interpret eruptive ages of ~15.95 Ma for a tuff at the top of the Vantage Member and ~8.72 Ma for a pumice within the upper Ellensburg Formation. Younger sedimentary interbeds of the Ellensburg Formation are more common and thicker compared to older interbeds, suggesting a migrating and (or) deepening depositional center. Between ~5.4 and ~3.3 Ma, deposition transitioned from largely Cascade-sourced sediments of the Ellensburg Formation to basalt-derived sediments of overlying Pliocene(?) Quaternary deposits.

We identify reverse and thrust faults with oblique-slip on the flanks of bedrock highs. We interpret these range front faults as young, in-sequence thrust-type structures in a compressional or transpressional regime. Fault-related folds are associated with these structures, including several plunging, northeast- through west-trending anticlines and synclines. We map numerous northerly striking oblique-slip faults with varying offsets, especially where the folds of the Boylston Mountains bend northward.

Geophysical modeling of gravity and aeromagnetic data suggests laterally abrupt thickness changes in basaltic units. These thickness changes may be related to (1) ramp-flat geometries under uplifts, and (or) (2) periods of syn-eruptive development of local accommodation space, where CRBG rocks experienced concurrent footwall thickening (growth strata) and hanging-wall thinning (uplift-related erosion).

LIST OF MAP UNITS

Holocene to Pleistocene Nonglacial Deposits

af **Artificial fill (Holocene)**—Cobbles, pebbles, sand, and boulders; poorly sorted and unconsolidated; placed to elevate home sites, roads, and infrastructure.

mf **Modified land (Holocene)**—Sand–through boulder-sized material redistributed to modify topography for industrial, agricultural, and general uses, including, but not limited to, gravel pits, rock quarries, aggregate mines, and home sites.

Qp **Peat (Holocene to Pleistocene)**—Organic and organic-rich sediment; includes peat, gyttja, muck, silt, and clay; typically in closed depressions; mapped in natural or man-made wetlands, bog areas, and ephemeral water bodies.

Qls **Landslide deposits (Holocene to Pleistocene)**—Clastic aggregate and scree; generally loose and poorly consolidated; clay- to boulder-sized material; angular to subangular, typically matrix supported, and less commonly clast supported near talus piles; unstratified and structureless; unit contains rubble of sand, silt, clay, cobbles, pebbles, boulders, and diamicton of mostly basalt clasts.

Holocene to Pliocene Alluvial and Colluvial Deposits

Stream channel and stream flood (overbank) deposits and terraces. Deposits include pebbles, cobbles, sand, silt, clay, peat, and boulders, all in varying amounts and thicknesses. Colors range from light tannish gray to medium brown. Clasts are angular to well rounded; unconsolidated to moderately to weakly compacted and poorly to moderately cemented. The clasts are typically cobbles with sand and pebbles, well rounded, and moderately to well sorted. The deposits are pyritized near the Yakima River and mostly basaltic in the East Kittitas quadrangle.

Qa **Alluvium (Holocene)**—Stream channel deposits on active flood plains and commonly flanking creeks; unit is narrowly distributed throughout low elevations of the map area.

Qca **Intermediate-age alluvium (Holocene to Pleistocene)**—Stream-flood (overbank) and old channel deposits near active flood plains; deposited where ongoing alluvial deposition from overbank flow is possible but less evident compared to unit Qa; surfaces of unit Qca are slightly elevated relative to surfaces of unit Qa. Unit Qca is commonly indistinguishable from alluvial fan unit Qaf1 based on lithology and height relative to unit Qa. However, unit Qca tends to be found (1) closer to unit Qa than unit Qaf1, and (2) at slightly lower elevations than unit Qaf1.

Qaf **Alluvial fan deposits (Holocene to Pliocene?)**—Sand, gravel, and relict debris flows deposited in alluvial fans; generally brown; unconsolidated to poorly consolidated; silt- to boulder-sized clasts; angular to subrounded; unsorted to poorly sorted; matrix to clast supported; contains basalt of the Columbia River Basalt Group (CRBG) and rhyolite; thickness is generally less than 15 m but varies with age, where older units tend to be thicker. Unit is subdivided and numbered from 1 (lowest and youngest) to 4 (highest and oldest) based on relative elevation above the modern stream level and differences in surface morphologies.

Qtg **Terrace gravel deposits (Pleistocene)**—Gravel deposited in fluvial setting in Yakima canyon; light to medium brown; loose, poorly cemented; pebbles and cobbles with boulders and medium sand; rounded to subrounded; poorly sorted; matrix supported. Polymictic clasts are from Cascade Range. Mapped in detail by Kelsay and others (2017) and Bender and others (2016).

Rtg **Cobble gravel (Pliocene)**—Angular to well-rounded cobble gravel with clay to sand interbeds deposited in alluvial fans and the paleo-Yakima River channels, and conglomerates flanking Manastash Ridge and the Boylston Mountains. The unit forms flat incised surfaces, polymictic near the Yakima River corridor and basaltic dominated elsewhere; poorly to mildly consolidated; mildly to moderately cemented; thickness ranges from several meters to more than 30 m; coeval with Thorp Gravel.

Tertiary Sedimentary and Volcanic Bedrock

SEDIMENTARY ROCKS OF THE ELLENSBURG FORMATION

Tew **Ellensburg Formation, undivided (middle Miocene to early Pliocene?)**—Feldsarenitic and conglomeratic of fluvial, lacustrine, volcaniclastic, and colluvial origin underlying, intercalated with, and overlying rocks of the CRBG; light to medium brown to light to medium gray; mildly to strongly weathered; moderately indurated; where interbedded in the CRBG, unit is generally fine to coarse-grained sandstone; where unconformably overlying the CRBG, unit is locally volcaniclastic mudstones and sandstones overlain by conglomeratic colluvium on the northern flank of Manastash Ridge. Rounded to angular; well to moderately sorted; clasts supported. Ellensburg Formation sandstone is typically pale yellowish brown where upper Ellensburg unit overlying CRBG can be >120 m thick; commonly has abundant, conspicuously dark white mica, whereas older members interbedded with basalts commonly have white mica in varying amounts. Locally subdivided into:

Tcm **Conglomeratic rocks of the upper Ellensburg Formation (late Miocene)**—Conglomerate with fine sedimentary lenses and paleosols that overlie the CRBG; light brown to medium brown, mild to moderately weathered; mild to moderately cemented; pebbles, cobbles with few boulders, also containing coarse to fine sands; subangular to well rounded; poorly sorted, matrix supported. Structureless to laminated; polymict dominated by Cascade-sourced volcanic clasts near Yakima River corridor and basaltic dominated in East Kittitas.

Tvs **Volcaniclastic rocks of the upper Ellensburg Formation (late Miocene)**—Volcaniclastic sandstone, siltstone, and mudstone with some conglomerate directly overlying the Wanapum Basalt; grey, pinkish brown, or light brown, moderately weathered; moderately indurated to friable, mildly to moderately cemented; lithology varies from fine ashy mudstone to conglomerate; sandstone with pumice and lenses of accretionary lapilli ash fall tuff; subangular to rounded, poorly sorted, matrix supported; common paleosols and pedogenic sediments in coeval map area. Unit is >20 m thick and found in canals and ravines at mid elevations on Manastash Ridge.

Tlmc **Limuma Creek Member of the lower Ellensburg Formation (middle Miocene)**—Tuffaceous sandstone, that can be micaceous and rarely diatomaceous; light gray to white, mildly to moderately weathered, mildly to moderately cemented, medium grained, well sorted and matrix supported with common pumice and glassy to rare polymict pebble layers; less than 15 m thick. Member was formerly known by a derogatory term for Indigenous women that has since been replaced by the name Limuma Creek. Includes the Quincy diatomite.

Tvm **Vantage Member of the lower Ellensburg Formation (middle Miocene)**—Micaceous and volcaniclastic sandstone and siltstone; light brownish-gray to light gray or white; moderately to strongly weathered; mildly to moderately indurated, weakly cemented; subangular; fine to coarse grained, commonly well sorted, grain supported. The unit is less than 30 m thick and is stratigraphically between the basalt of Museum of the Sentinel Bluffs Member, unit M_{Mus}, and the overlying Wanapum Basalt.

Tvcc **Coleman member of the lower Ellensburg Formation (middle Miocene) (line unit only)**—Sandstone and siltstone underlying the Sentinel Bluffs Member of the CRBG; medium brown to light grey; generally fine- to medium-grained micaceous sandstone; estimated to be less than 20 m thick and mapped as a line unit in Yakima canyon in the southwestern corner of the map area.

VOLCANIC ROCKS OF THE COLUMBIA RIVER BASALT GROUP (CRBG)

Mwa **Wanapum Basalt, undivided (middle Miocene) (cross section only)**—Basalt, described in detail in the following units; dark grey to grayish brown, weathers to medium grey; fine grained, sparsely porphyritic, or weakly glomerocrystic; commonly with groundmass crystals at least 1.0 mm in size; less commonly aphyric than GRB units, <10% in thick. The formation unconformably overlies the Vantage Member (unit M_{Vac}), and upper Ellensburg units—especially unit M_{Tew}, which unconformably onlaps onto unit M_{Tew}. Locally subdivided into:

Mwp **Priest Rapids Member (middle Miocene)**—medium- or very coarse-grained basaltic andesite with groundmass crystals >2 mm in size; dark gray to grayish brown, weathers to light orangish brown; well indurated; mostly aphyric to weakly porphyritic, typically very coarse grained, sometimes with a gabbro-like texture (basaltic pegmatoid); unit is ~15 m thick at Vanderbilt Gap and poorly exposed in the canals on northern flank of Manastash Ridge.

Mwr **Roza Member (middle Miocene)**—Porphyritic basalt and basaltic andesite with medium to fine groundmass; dark gray to grayish brown, weathers dark to medium brown; diagnostically porphyritic (~2–10% total phenocrysts) with common centimeter-scale plagioclase phenocrysts typically >10 phenocrysts per hand sample(s); top of flow is more aphyric. The unit is ~15–30 m thick and invaded the Limuma Creek Member (unit M_{Tlmc}), where Roza lava flow(s) entered wet sediments that would become unit M_{Tlmc}, thereby the rocks of unit M_{Tlmc} underlie, intermingle, and overlie the Roza Member.

Mwf **Frenchman Springs Member, undivided (middle Miocene)**—Medium-grained basalt to basaltic andesite that is sparsely porphyritic to weakly glomerocrystic or aphyric; dark gray to grayish brown; well undurated; commonly sparsely porphyritic to glomerocrystic with <5% total phenocrysts; unit is commonly <30 m thick or <90 m thick from Umtanum Ridge and Badger Hollow but may thicken into valley. Less porphyritic than the Roza Member and finer grained than the Priest Rapids Member

Mwsf **Basalt of Sand Hollow (middle Miocene)**—Medium-grained basalt and basaltic andesite that is weakly porphyritic, with ~1% total phenocrysts, or aphyric; typically contains about two plagioclase phenocrysts per hand sample, and may contain glomerocrysts; medium to dark gray, weathers to dark reddish brown or reddish gray; well indurated; ~1 m wide columns can internally exhibit flat to curved platy foliation. Pillow breccia and palagonite are locally common. The unit is less than ~60 m thick and well exposed on north flank of the Boylston Mountains and summit of Manastash Ridge. The unit commonly overlies the Vantage Member in the western map area.

Mgr **Grande Ronde Basalt (GRB), undivided (middle Miocene) (cross section only)**—Basaltic andesite, described in detail in the following units. The Meridian well contains ~2,029 m of total basalt thickness (S. Reidel, written commun., 2020), presumably all Grande Ronde Basalt. The unit contains reverse (R) and normal (N) magnetostratigraphic units (MSU), from oldest to youngest: R1, N1, R2, and N2. Unit M_{Mg} is mapped where a lack of geochemistry prevented us from identifying a specific subunit, and in the cross section where the subunits are grouped at depth. Locally subdivided into:

Mgsf **Sentinel Bluffs Member, undivided (middle Miocene)**—Basaltic andesite; aphyric. The map area contains five subunits, from oldest to youngest: basalt of McCoy Canyon, California Creek, Spokane Falls, Slembert Creek, and Museum. Cumulatively, the Sentinel Bluffs Member is at least ~140 m thick, with the Meridian well showing a thickness range of 93–126 m (S. Reidel, written commun., 2020). We map unit M_{Mgs} as undivided (grouped) where geochemical results are unavailable to divide exposures into subunits but where we remain reasonably confident that the exposure belongs to the Sentinel Bluffs Member based on interpolations with nearby stratigraphic relationships; found on the northeastern portion of the map area. With available geochemistry, we subdivide the Sentinel Bluffs Member into:

Mgsm **Basalt of Museum (middle Miocene)**—Fine-grained basaltic andesite with groundmass crystals ranging in size from 0.05 to 0.6 mm and very rare phenocrysts of plagioclase up to 2.5 mm in size; most commonly aphyric. The unit is ~36–60 m thick, contains at least two flows, and is locally the top of the Sentinel Bluffs Member.

Mgsc **Basalt of Slembert Creek (middle Miocene)**—Fine-grained basaltic andesite with groundmass crystals ranging in size from 0.05–0.4 mm and very rare phenocrysts of plagioclase as large as 1.6 mm; most commonly aphyric. The unit is less than ~55 m thick and it contains at least two flows. The unit is sparsely mapped and less common than unit M_{Mgsm} and consistently overlies flows with Spokane Falls-type compositions in the eastern map area.

Mgsf **Basalt of Spokane Falls (middle Miocene)**—Fine-grained basaltic andesite with groundmass crystals ranging in size from 0.1–0.4 mm and very rare phenocrysts of plagioclase as large as 1.2 mm; most commonly aphyric. The unit is very common and <65 m thick and contains at least two flows. The unit may interfinger with Slembert Creek-type compositions in the southwest map area.

Mgsc **Basalt of California Creek (middle Miocene)**—Fine-grained basaltic andesite; medium gray, strongly weathered; moderately to well indurated; aphyric; ~50 m thick; found only at geochemistry site G284; found along the Boylston Fault in the southern portion of the map area.

Mgcc **Basalt of McCoy Canyon (middle Miocene)**—Fine-grained basaltic andesite with groundmass crystals ranging in size from ~0.1–0.5 mm and very rare phenocrysts of plagioclase as large as 1.2 mm; most commonly aphyric. The unit is ~65–90 m thick, contains two or three flows, and is locally the base of the Sentinel Bluffs Member.

Mwt **Winter Water Member (middle Miocene)**—fine to medium-grained basaltic andesite; locally aphyric, but may be sparsely to abundantly glomerocrystic or less commonly porphyritic; medium to dark gray, weathers medium to dark brown; well undurated; ~40–100 m thick and thickens eastward. The Meridian well shows a thickness range of 58–98 m for unit M_{Mgs} (S. Reidel, written commun., 2020). The unit contains at least two flows with well-developed middle entablature and a compositionally banded colonnade composed of stacks of centimeter-scale tabular columns. Synonymous with the "Umtanum flow."

Mor **Ortley member (middle Miocene)**—Fine- to medium-grained basaltic andesite with groundmass crystals up to 1 mm in size; aphyric. The unit is at least ~46 m thick in Yakima canyon and may thicken eastward at depth to ~230 m in the BISSA well and ~215–251 m in the Meridian well (S. Reidel, written commun., 2020); contains at least two flows. The unit is part of the N2 MSU, representing the base of N2 MSU in the map area, and is invasive into the Colonnade member of the Ellensburg Formation.

Mgc **Grouse Creek member (middle Miocene)**—Medium- to fine-grained basaltic andesite with groundmass crystals ranging in 1.5 mm in size; aphyric groundmass is very slightly porphyritic with <1% phenocrysts. Base of unit is not observed. From map patterns, the unit is >140 m thick, but may be considerably thicker north of the map area; at depth the unit is >90 m thick in the BISSA well and ~46–91 m thick at the Meridian well (S. Reidel, written commun., 2020). The unit contains at least one flow and is well exposed in Yakima canyon.

CORRELATION OF MAP UNITS

PERIODS AND EPOCHS BASED ON USGS FACT SHEET 2015-3009



Red font denotes remanent paleomagnetic polarity

GEOLOGIC SYMBOLS

Mass Wasting (overlay mw)—Mapped where hummocky landforms suggest mass movement on unstable slopes, but evidence for landslide deposits is inconclusive

Quaternary loess (overlay Ql)—Light brown to medium brown eolian loess; moderately weathered; low density; poorly compacted; <1 m thick; mostly silt and clay; mildly to moderately to poorly sorted; matrix supported; internally structureless; forms 1 meter-tall irregularly spaced mounds.

Miocene hyaloclastite (overlay hy)—Volcaniclastic aggregate in an hematin basalt flows pillow breccia with volcanic glass (tachylyte + sideromelane) and its alteration product palagonite. Color ranges from light yellowish brown or tan and generally strongly weathered, less dense than basalt, moderately well consolidated, generally composed of sand- to boulder-sized clasts in a very fine-grained matrix; angular to subangular; poorly sorted; some exposures may contain cobble- to boulder-size fragments of basalt pillows (centimeter to meter scale) that are matrix supported. Where pillows are absent, exposures are convoluted and structureless. Hyaloclastite forms from the quenching of hot lava flows in water, ice, or wet sediments. Where unconsolidated sediments this is called peperite hyaloclastite or peperite. Thickness varies but is generally less than 30 m. Overlay is common near the base of unit M_{Mgsm}, near the top of unit M_{Mgs}, and throughout unit M_{Mgr}.

Contact—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable.

Gradational contact—Identity and existence certain; location uncertain

Geologic boundary—Identity and existence certain; solid where location accurate; long-dashed where approximate; short-dashed where inferred. In this map, symbol shows internal contacts between lava flows of the same unit, or boundaries between alluvial fans of similar ages

Fault—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable

Thrust fault—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; sawtooth on upper plate

Reverse fault—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; rectangles on upthrown block

Strike-slip fault, right lateral offset—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; arrows show relative motion; rectangles on upthrown block

High-angle dip-slip fault—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; relative motion shown by U and D

Oblique-slip fault, high-angle right-lateral offset—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; relative motion shown by U and D; rectangles on upthrown block

Oblique-slip fault, high-angle left-lateral offset—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; relative motion shown by U and D

Anticline—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; arrows on line indicates plunge direction

Syncline—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable; arrows on line indicates plunge direction

Monocline, synclinal bend—Long-dashed where approximate; dotted where concealed; arrows show direction of dip; shorter arrow on steeper limb; arrow on line indicates plunge direction

Geologic unit too thin to show as a polygon—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; dotted where concealed; queried where identity or existence questionable

Scarp of unknown origin—Hachures point down slope

Dike—Identity or existence questionable; location accurate

Lineament of unknown origin

Cross section line

Perennial stream

Intermittent stream

Small, minor vertical or near-vertical joint—showing strike

Small, minor inclined fault—showing strike and dip

Small, minor vertical or near-vertical fault—showing strike

Inclined slickenline, groove, or striation on fault surface—showing bearing and plunge

Inclined bedding in unconsolidated sedimentary deposits or unconsolidated fragmental deposits of volcanic origin—showing strike and dip

Inclined lava flow banding in unconsolidated sedimentary deposits or unconsolidated fragmental deposits of volcanic origin—showing strike and dip

Inclined contact—showing dip value and direction

Inclined flow banding, lamination, layering, or foliation in igneous rock—showing strike and dip

Shear—showing strike and dip

Spring

Age site, radiometric

Age site, fossil

Age site, U-Pb, uranium-lead

Age site, infrared stimulated luminescence

Geochemistry sample location

Water well

Drill hole for hydrocarbon exploration or exploitation

Clast count sample

Surface data collected by Washington Geological Survey except for those shown as blue symbols (compiled surface data from R. Bentley and J. Powell, Central Washington University, unpub. mapping, 1980–1989).

Inclined bedding—showing strike and dip

Overturned bedding—showing strike and dip

Inclined bedding, where top direction of beds is known from local features—showing strike and dip

Small, minor inclined joint—showing strike and dip

Inclined fold hinge of small, minor syncline—showing bearing and plunge

FIGURE M1A, M1B, M1C
PHYSICAL FEATURES

BAF1—Boylston anticline fault 1
BAF2—Boylston anticline fault 2
BFZ—Boylston aeromagnetic low
BFZ—Boylston fault zone
FTL—Fault tip low
KVF—Kittitas Valley fault
KVG—Kittitas Valley gravity gradient
KVL—Kittitas Valley gravity low
MAH—Manastash aeromagnetic high
MAL—Manastash aeromagnetic low
MFZ—Manastash fault zone
MAH—Manastash aeromagnetic high
WAL—Wanapum aeromagnetic low



Figure M1A. Combined isostatic gravity, aeromagnetic (reduced-to-pole), and hillshade topography map. Gravity contours are 0.5 mGal. Gravity stations are light gray dots. Yellow dots are gravity mas-spots (explained in Pamphlet). X–X' and Y–Y' show the end points of the geophysical models in Figures M1B and M1C, respectively. Gray boxes mark the limits of the Kittitas and East Kittitas quadrangles.

Figure M1B. Two-dimensional potential-fields forward model of profile X–X'. Units and properties shown in the bottom panel predict gravitational and magnetic anomalies shown in the top two panels (p–saturated bulk density in kg/m³; χ–magnetic susceptibility in SI x 10^{-3}). Gravity data shown are no more than 750 m in map view from the model line. The diameters for the gravity data points are equal to the data error; magnetic data error is too small to show at print scale. "Right path" shows the elevation of the airplane that flew the aeromagnetic survey. Geologic units and the Boylston Fault property are shown in Figure M1A. Vertical exaggeration is 2:1. Datum is sea level. Geologic units are projected onto the model perpendicularly. Thicknesses of Ellensburg Formation units and Oligocene sediment are slightly shallower than in our model.



Figure M1C. Two-dimensional potential-fields forward model of profile Y–Y'. See geophysical figure table and legend for symbology and labeling explanation.

	Model Properties			Magnetic Remanence		Average Measured Rock Properties	
	P	X		Mr	Oxcl	P	X
Quaternary/Ellensburg Formation	1,800	0-6	0	—	—	1,810	6
Priest Rapids member (reversed)	2,770	8	2.5	-65	191	—	—
Ellensburg Formation-Limuma Creek	1,810	0	0	—	—	1,810	6
Roza member (reversed)	2,870	18	2.5	-38	198	2,870	9 (18 max)
Basalt of Sand Hollow	2,760	16	2.5	58	0	2,760	6 (16 max)
Basalt of Sand Hollow breccia	2,240	6	1–2.5	58	0	2,240	6
Basalt of Ginkgo	2,850	2	2.5	44	158	2,850	2
Ellensburg Formation-Vantage	1,810	0-6	0	—	—	1,810	6
Basalt of Museum	2,720–2,770	2	1.5	58	16	2,720	2
Basalt of Museum breccia	2,300	2	0–1.5	58	16	2,240	6
Basalt of Slembert Creek	2,620	10	1.5	58	16	2,620	10
Basalt of Spokane Falls	2,650	4	1.5	58	16	2,650	4
Basalt of Spokane Falls breccia	2,300	4	1.5	58	16	2,240	6
Basalt of McCoy Canyon	2,660	3	1.5	58	16	2,660	3
Winter Water member	2,800	7	1.5	58	16	2,800	7
Ortley member	2,520	7	1.5	58	16	2,520	7
Grouse Creek member (reversed)	2,560	11	4.0	-72	219	2,560	5 (11 max)
Grouse Creek member breccia (rev.)	2,100	11	4.0	-72	219	2,240	6
Wapshilla Ridge member (reversed)	2,590	15	4.0	-72	219	2,650	13 (15 max)
Grande Ronde N1	2,770	6	2.5	58	355	—	—
Grande Ronde R1 (reversed)	2,700–2,780	6	2.5	-66	171	—	—
Eocene/Oligocene sediment	2,450	0	0	—	—	2,420	<1

CROSS SECTION EXPLANATION

Geologic units too thin to show as polygons at the scale of the cross section—Solid where location accurate; long-dashed where approximate; short-dashed where inferred; queried where identity or existence questionable; vertical tics mark separate units.

Arrows show relative fault movement in the plane of the cross section.

Arrow point shows fault movement toward the viewer; arrow feathers show fault movement away from the viewer.

Water well or boring



Lambert conformal conic projection
North American Datum of 1983
Base map from U.S. Geological Survey Kittitas and East Kittitas 7.5-minute quadrangles, 2020
Shaded relief generated from a lidar bare-earth digital elevation model (available from the Washington Geological Survey, https://lidarportal.dnr.wa.gov/)
GIS by Andrew J. Sadowski and Logan R. Wetherell
Digital cartography by Daniel E. Cox and J. Eric Schuster
Editing and production by Susan R. Schuster and Nikolas Mullins

Disclaimer: Neither the State of Washington, nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the State of Washington or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the State of Washington or any agency thereof.

SCALE 1:24,000

1,000 0 1,000 2,000 3,000 4,000 5,000 6,000 7,000 FEET
0.5 0 0.5 1 MILE
1 KILOMETER
contour interval 20 feet

This geologic map was funded in part by the U.S. Geological Survey National Cooperative Geologic Mapping Program under USGS award number G22AC00364.

Research supported by the U.S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award number G22AC00364. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.